

Asymmetrical Reinforcement and *Wolbachia* Infection in *Drosophila*

John Jaenike^{1*}, Kelly A. Dyer^{1,2}, Chad Cornish^{1‡}, Miranda S. Minhas¹

1 Department of Biology, University of Rochester, Rochester, New York, United States of America, **2** Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, United Kingdom

Reinforcement refers to the evolution of increased mating discrimination against heterospecific individuals in zones of geographic overlap and can be considered a final stage in the speciation process. One the factors that may affect reinforcement is the degree to which hybrid matings result in the permanent loss of genes from a species' gene pool. Matings between females of *Drosophila subquinaria* and males of *D. recens* result in high levels of offspring mortality, due to interspecific cytoplasmic incompatibility caused by *Wolbachia* infection of *D. recens*. Such hybrid inviability is not manifested in matings between *D. recens* females and *D. subquinaria* males. Here we ask whether the asymmetrical hybrid inviability is associated with a corresponding asymmetry in the level of reinforcement. The geographic ranges of *D. recens* and *D. subquinaria* were found to overlap across a broad belt of boreal forest in central Canada. Females of *D. subquinaria* from the zone of sympatry exhibit much stronger levels of discrimination against males of *D. recens* than do females from allopatric populations. In contrast, such reproductive character displacement is not evident in *D. recens*, consistent with the expected effects of unidirectional cytoplasmic incompatibility. Furthermore, there is substantial behavioral isolation within *D. subquinaria*, because females from populations sympatric with *D. recens* discriminate against allopatric conspecific males, whereas females from populations allopatric with *D. recens* show no discrimination against any conspecific males. Patterns of general genetic differentiation among populations are not consistent with patterns of behavioral discrimination, which suggests that the behavioral isolation within *D. subquinaria* results from selection against mating with *Wolbachia*-infected *D. recens*. Interspecific cytoplasmic incompatibility may contribute not only to post-mating isolation, an effect already widely recognized, but also to reinforcement, particularly in the uninfected species. The resulting reproductive character displacement not only increases behavioral isolation from the *Wolbachia*-infected species, but may also lead to behavioral isolation between populations of the uninfected species. Given the widespread occurrence of *Wolbachia* among insects, it thus appears that there are multiple ways by which these endosymbionts may directly and indirectly contribute to reproductive isolation and speciation.

Citation: Jaenike J, Dyer KA, Cornish C, Minhas MS (2006) Asymmetrical reinforcement and *Wolbachia* infection in *Drosophila*. PLoS Biol 4(10): e325. DOI: 10.1371/journal.pbio.0040325

Introduction

When incipient species come into secondary contact, natural selection against the production of unfit hybrids may favor the evolution of increased behavioral isolation between them, a process termed reinforcement [1,2]. Serious questions about the reality of reinforcement have been raised [3], but there is now good empirical evidence as well as a convincing theoretical foundation for reinforcement [4,5]. One possible outcome of the process of reinforcement is a pattern of reproductive character displacement, in which behavioral isolation is greater between individuals obtained from sympatric populations of two species than between individuals obtained from allopatric populations [3,6,7]. Reinforcement is typically modeled as acting symmetrically between species with partially overlapping ranges. However, the process may affect the two species differently if the range of one species is completely nested inside that of another [8], if levels of gene flow into the zone of sympatry differ between the species, or if there are differences in the fitness of reciprocal F1 hybrids.

One possible cause of unequal hybrid fitness is unidirectional cytoplasmic incompatibility (CI) induced by *Wolbachia* infection in one of the two species [9]. CI can occur in crosses between *Wolbachia*-infected males and uninfected females and

results in high levels of embryonic mortality among the offspring. In the reciprocal cross between uninfected males and infected females, offspring survival is not affected. Thus, if one species in a zone of sympatry is infected with *Wolbachia* and the other is not, a large difference in the number of surviving offspring produced in reciprocal crosses between species may result.

Several aspects of *Wolbachia* biology suggest that *Wolbachia* could be a common cause of such asymmetry in insects. *Wolbachia* are widely distributed across insect species, where infection is most often manifested as CI [10,11]. Phylogenetic analyses indicate that infections are generally short-lived on

Academic Editor: Mohamed Noor, Duke University, United States of America

Received: May 1, 2006; **Accepted:** August 2, 2006; **Published:** October 10, 2006

DOI: 10.1371/journal.pbio.0040325

Copyright: © 2006 Jaenike et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: CI, cytoplasmic incompatibility; mtDNA, mitochondrial DNA

* To whom correspondence should be addressed. E-mail: joja@mail.rochester.edu

‡ Current address: State University of New York Upstate Medical University, College of Medicine, Syracuse, New York, United States of America

an evolutionary time scale, so that sister insect taxa, including those that are only partially isolated, can differ in their infection status [12–14]. Although *Wolbachia* spread and are maintained as a result of the CI that they cause within species, the same phenomenon can be manifested in matings between species [13,15,16]. Consequently, there may be numerous pairs of partially isolated insect taxa between which there is asymmetrical interspecific CI.

Because CI leads to the death of offspring produced by matings between individuals that differ in infection status, there has been considerable interest in the role that *Wolbachia* may play in speciation [17,18]. Despite the appeal of this apparently simple isolating mechanism, there are substantial difficulties with speciation based solely on *Wolbachia*-induced CI [5,19]. From a theoretical perspective, CI causes unstable equilibrium in the prevalence of *Wolbachia* infection, rather than stable persistence of infected and uninfected populations [20,21]. Even bidirectional incompatibility entails an unstable equilibrium, with only one type expected at equilibrium [22]. Theoretical analyses also indicate that the spread of *Wolbachia* throughout a continuous population will occur much more rapidly than evolution of reproductive isolation [23]. Empirically, there is no evidence that *Wolbachia* contributes to hybrid sterility or to Haldane's rule, two of the most general features of animal speciation [5]. These findings indicate that *Wolbachia* cannot be the primary cause of many speciation events.

Nevertheless, *Wolbachia* may contribute, along with other mechanisms, to overall isolation between populations [13]. In particular, recent theoretical studies have shown that *Wolbachia*-induced CI can work in conjunction with geographically divergent selection on a nuclear locus to bring about reinforcement between populations [24]. Thus, *Wolbachia* may act as an agent for reinforcement of pre-existing isolating mechanisms, such as hybrid male sterility, and thus play a supplementary (not primary) role in the evolution of reproduction isolation between incipient species.

Although the most obvious consequence of reinforcement is decreased mating between individuals of different species, this process can also bring about behavioral isolation between sympatric and allopatric populations of the same species, as postulated theoretically [25] and as found in frogs [26,27] and *Drosophila* [28]. Thus, it is possible that secondary contact between a *Wolbachia*-infected species and a closely related uninfected species may bring about the evolution of behavioral isolation not only between the infected and uninfected species, but also between populations of the uninfected species.

We examined the evolutionary behavioral consequences of secondary contact between *D. recens* and *D. subquinaria*. Like most members of the quinaria species group, these two species are mycophagous, and mushrooms serve not only as larval food resources, but also as sites for adult courtship and mating. *D. recens* is infected with a strain of *Wolbachia* that causes strong intraspecific CI [29]. The prevalence of infection is about 98% among *D. recens* in the wild, which, along with dramatically reduced levels of mitochondrial DNA (mtDNA) diversity [13,30], indicates that the infection has spread to effective fixation. Reduced mtDNA variation is a common consequence of *Wolbachia* infection [31,32]. In contrast, the very closely related *D. subquinaria* is not infected with *Wolbachia*, and its high levels of mtDNA variation

indicate that it has not been infected in the recent evolutionary past [13,32]. *Wolbachia*-induced CI causes a dramatic (~90%) reduction in offspring production in crosses between *D. subquinaria* females and *D. recens* males, but the reciprocal cross yields numerous *Wolbachia*-infected viable hybrid progeny [13]. The F1 hybrid progeny of the latter cross are viable because there is no CI. Although the hybrid males are sterile, the hybrid females are fertile and could thus serve as conduits for introgression of genes between species and for passing genes back into the parental species.

Previous work on this pair of species revealed only moderate behavioral isolation between them, suggesting that hybridization might frequently occur in the wild if these species were to come into contact. However, as pointed out by Rokas [33], the experimental strains of *D. recens* and *D. subquinaria* had been derived from allopatric populations, and thus might not have been selected to discriminate against males of the other species. Sympatric populations of these species, if they exist, might exhibit greater levels of discrimination. *D. subquinaria* is currently known from western North America, whereas *D. recens* is known only from eastern North America [34], although Shoemaker et al. [13] had speculated that the two may be sympatric in Canada, where there is a continuous zone of boreal forest (the preferred habitat of *D. recens* and *D. subquinaria*) that spans the region between the known ranges of these species.

We show here that the ranges of *D. recens* and *D. subquinaria* include a broad region of sympatry in central Canada. Within this zone of sympatry, *D. subquinaria* exhibits much greater reproductive character displacement than *D. recens*, which is consistent with the expected effects of *Wolbachia*-induced CI in hybrid offspring. In accordance with certain models of reinforcement [25,35], sympatric females of *D. subquinaria* not only discriminate against males of *D. recens*, but also against conspecific but allopatric males. Molecular data from mtDNA indicate that *D. recens* and *D. subquinaria* have hybridized in the past, as required by theories of reinforcement. Patterns of variation at the mitochondrial *Cytochrome Oxidase I* (COI) locus indicate that there is little genetic differentiation between sympatric and many allopatric populations within each species, suggesting that reproductive character displacement has evolved in the face of considerable gene flow and is likely the result of selection rather than genetic isolation among populations.

Results

Geographic Range

Our collections revealed a substantial area of sympatry between *D. recens* and *D. subquinaria*, extending, at a minimum, from Jasper, Alberta, to The Pas, Manitoba, a linear extent of some 1,200 km (Figure 1 and Table 1). Both species were collected simultaneously in this area of sympatry, indicating that they are likely to encounter each other at mushrooms in the field. Allopatric populations of *D. recens* occur throughout eastern Canada and the northeastern United States, and extend south through the Appalachians to high-elevation sites in the Smoky Mountains of Tennessee and North Carolina. Allopatric populations of *D. subquinaria* occur over much of western North America. Thus, there are sympatric and allopatric populations for both of these species.




Figure 1. Collection Sites and Relative Abundance of *D. recens* (Gray) and *D. subquinaria* (Black)
 Satellite imagery courtesy of <http://www.GlobeXplorer.com>.
 DOI: 10.1371/journal.pbio.0040325.g001

Table 1. Populations and Strains Used in this Study and Number of Individuals from Each Population Sequenced at *COI*

Species	Region	Population Location ^a	Abbreviation ^b	Year Sampled	<i>n</i> <i>COI</i>
<i>D. subquinaria</i>	Allopatric, coastal	1. Seattle, Washington	S-S/WA	1997	6
		2. Olympic Peninsula, Washington	S-OP/WA	2005	12
		3. Port Hardy, British Columbia	S-PH/BC	1997	13 ^c
		4. Vancouver, British Columbia	S-V/BC ^b	2001	6
	Allopatric, inland	5. Peachland, British Columbia	S-P/BC	1997	15 ^c
		6. Coeur D'Alene, Idaho	S-CD/ID ^b	2001	0
		7. Deary, Idaho	S-D/ID	2000	3
		8. Columbia Falls, Montana	S-CF/MT ^b	2002	6
	Sympatric	9. Big Sky, Montana	S-B5/MT ^b	2002	17
		10. Jasper, Alberta	S-J/AB ^b	2002	18
		11. Edmonton, Alberta	S-E/AB ^b	2002	7 ^c
		12. Winston Churchill, Alberta	S-WC/AB	2002	20
		13. Prince Albert, Saskatchewan	S-PA/SK	2002	2
		14. The Pas, Manitoba	S-P/MB ^b	2002	5
<i>D. recens</i>	Allopatric	15. Charlottetown, Prince Edward Island	R-C/PEI ^b	2003	4
		16. Brunswick Pines, Maine	R-BP/ME	1995	12
		17. Bethlehem, New Hampshire	R-B/NH ^b	2002	7
		18. Eagle Bay, New York	R-EB/NY	1995	13
		19. Great Smoky Mountains, Tennessee	R-SM/TN ^b	2001	17
		20. Rochester, New York	R-R/NY ^b	2002	9
		21. Wawa, Ontario	R-W/ON ^b	2002	0
		22. Munising, Michigan	R-M/MI ^b	2002	6
		23. Bemidji, Minnesota	R-B/MN ^b	2002	6
		24. Minot, North Dakota	R-M/ND ^b	2002	7
	Sympatric	14. The Pas, Manitoba	R-P/MB ^b	2002	9
		13. Prince Albert, Saskatchewan	R-PA/SK	2002	21
		12. Winston Churchill, Alberta	R-WC/AB	2002	16
		11. Edmonton, Alberta	R-E/AB ^b	2002	21
		10. Jasper, Alberta	R-J/AB ^b	2002	8

^aNumbers refer to locations on map in Figure 1.

^bStrains derived from these populations are designated similarly, but in italics (e.g., *S-S/WA*).

^cThese samples each included one individual that carried a *D. recens* mtDNA haplotype.

DOI: 10.1371/journal.pbio.0040325.t001




Figure 2. Proportion of Either Sympatric (sym) or Allopatric (allo) Females of *D. recens* and *D. subquinaria* That Mated with Either *D. recens* or *D. subquinaria* Males in the Mass Population Choice Experiments. For example, within trials, an average of 20 of the 100 allopatric females of *D. subquinaria* mated with *D. recens* males and 38 of them mated with *D. subquinaria*. Means $\pm$ standard error among three replicate trials. DOI: 10.1371/journal.pbio.0040325.g002

Mating Experiments—Mass Population

The proportion of sympatric and allopatric *D. recens* and *D. subquinaria* females that mated with either *D. recens* or *D. subquinaria* males is shown in Figure 2. In each of the three trials, the allopatric females of *D. subquinaria* were significantly more likely to mate with *D. recens* males than were sympatric females, with an across-trials combined probability of $p = 10^{-8}$. In our experiments, sympatric females never mated with *D. recens* males. In contrast, the difference between sympatric and allopatric females of *D. recens* was non-significant within each trial, and the combined across-trials probability was $p = 0.20$. Thus, in the mass mating experiments, *D. subquinaria* females exhibited very strong reproductive character displacement, whereas *D. recens* females showed little.

Mating Experiments—No Choice

Figure 3 presents the numbers of females from sympatric and allopatric populations of *D. recens* and *D. subquinaria* that mated with heterospecific males in no-choice trials. Although contextually very different from the mass population assays, the results are essentially the same. None of the sympatric females of *D. subquinaria* mated with *D. recens* males, whereas a considerable proportion (26% overall) of the allopatric females did. A logistic regression of the probability of mating by females of *D. subquinaria* revealed a highly significant effect of population type (sympatric versus allopatric; Wald $\chi^2 = 17.88$, $p < 0.0001$), but no effect of individual population within population type ($\chi^2 = 1.02$, $p = 0.91$) or trial ($\chi^2 = 2.12$, $p = 0.35$).

As with the mass mating experiment, we found little evidence for reinforcement within *D. recens*. A logistic regression of the probability of female mating revealed nonsignificant effects of population type (sympatric versus allopatric; Wald $\chi^2 = 1.99$, $p = 0.16$) and trial ($\chi^2 = 0.50$, $p = 0.78$). However, we did find a significant effect of individual population with population type ($\chi^2 = 23.93$, $p = 0.0001$), an




Figure 3. Proportion of Females That Mated with Heterospecific Males in the No-Choice Trials

Means $\pm$ standard error among replicate trials. Females are organized by species, population type (sympatric or allopatric within species), and population within type. Population labels from Table 1, with S- or R-prefix deleted. The populations correspond with the sites in Figure 1 as follows: V-BC, 4; CD-ID, 6; BS-MT, 9; P-MB, 14; E-AB, 11; J-AB, 10; R-NY, 20; B-NH, 17; B-MN, 23. DOI: 10.1371/journal.pbio.0040325.g003

effect due primarily to the much greater propensity of females of the allopatric *R-R/NY* strain to mate with *D. subquinaria* males (Figure 3).

Intraspecific Mate Choice—No Choice

We assessed whether there is any geographic differentiation in mate preference within each species. For *D. recens*, a logistic regression indicated that there was no significant effect of trial (Wald $\chi^2 = 0.98$, $p = 0.32$), female population type (sympatric versus allopatric; $\chi^2 = 0.01$, $p = 0.93$), male population type ($\chi^2 = 3.62$, $p = 0.06$), or female population type by male population type interaction ($\chi^2 = 0.002$, $p = 0.96$) on probability of a mating (Figure 4). Thus, there is no evidence of geographic differentiation in mate preferences.

In *D. subquinaria*, there were highly significant effects of female population type (coastal-allopatric, inland-allopatric, and inland-sympatric; $\chi^2 = 88.0$, $p < 0.0001$) and female population type by male population type ($\chi^2 = 62.6$, $p < 0.0001$), and trial ($\chi^2 = 12.3$, $p = 0.006$), whereas the effect of male population type was not significant ($\chi^2 = 1.80$, $p = 0.40$). The significant effects were due to the low probability of mating between sympatric females and allopatric males, an effect that was consistent across trials. Because the significant female-by-male interaction could be due either to the coastal-inland differentiation or to the differences between inland-sympatric and inland-allopatric populations, we also analyzed the data including only the inland populations. In this case, we found significant effects of female population type ($\chi^2 = 77.7$, $p < 0.0001$), male population type ($\chi^2 = 10.0$, $p = 0.0015$), trial ($\chi^2 = 11.6$, $p = 0.01$), and, most importantly, female population type by male population type ($\chi^2 = 62.4$, $p < 0.0001$). Figure 5 illustrates that significant interaction effects




Figure 4. Proportion of *D. recens* Females from Either Sympatric or Allopatric Populations That Mated with Either Sympatric or Allopatric *D. recens* Males in No-Choice Tests

Data from different allopatric populations combined and that from sympatric strains combined. Mean $\pm$ standard error among the two replicate trials.

DOI: 10.1371/journal.pbio.0040325.g004

are due to the reluctance of sympatric females to mate with males from both inland and coastal allopatric populations.

Post-Mating Isolation within *D. recens* and *D. subquinaria*

To examine the extent of differentiation among conspecific populations of *D. subquinaria* and *D. recens*, we determined whether any qualitative post-mating isolating mechanisms operate between our strains. Within both species, all pairwise crosses among strains yielded viable male and female progeny. For each pairwise cross, all three replicate cultures that were set up as crosses between F1 male and F1 female progeny yielded viable male and female F2 progeny. Finally, for each pairwise mating, all three replicate cultures involving crosses between F2 males and females yielded viable male and female progeny. Thus, both male and female F1 and F2 are viable and fertile in all pairwise crosses between geographic strains within species. There is no evidence of hybrid sterility or inviability in any of these intraspecific crosses. Although sympatric females of *D. subquinaria* are reluctant to mate with allopatric males, prolonged confinement with allopatric males did yield matings and offspring. Thus, there is no evidence of qualitative post-mating isolation among any strains of *D. subquinaria*, and likewise for *D. recens*.

Population Structure

Both analyses of molecular variances and F_{ST} -based statistics reveal that there is no overall significant differentiation between sympatric and allopatric groups of populations of either *D. recens* or *D. subquinaria*, although there is significant differentiation among populations within groups in both species (Table 2 and Protocol S1). Among the




Figure 5. Proportion of *D. subquinaria* Females from Different Regions That Mated with *D. subquinaria* Males from These Same Regions in No-Choice Tests

Coastal-allopatric (CA), inland-allopatric (IA), or inland-sympatric (IS). Mean $\pm$ standard error among the four replicate trials.

DOI: 10.1371/journal.pbio.0040325.g005

populations of *D. subquinaria*, there is significant differentiation between the coastal and inland populations (Figure 6 and Protocol S1). These regions are separated by the Pacific Cordillera and the high desert ecosystems of central Oregon, Washington, and British Columbia, suggesting that these areas constitute a substantial barrier to gene flow in *D. subquinaria*, as documented in other species of the Pacific Northwest [36]. Considering only the inland populations, which include all of the sympatric populations, there is no significant differentiation between those that are sympatric with *D. recens* and those that are allopatric ($S_{nn} = 0.52$; $p = 0.69$) (Figure 6). In *D. recens*, there is no significant differentiation between sympatric and allopatric populations (Table 2 and Figure 7), although there is significant within-group differentiation, due largely to the Great Smoky Mountains population, which occurs at the southern periphery of the species range (Figure 7). The Smokies population alone includes mtDNA haplotypes in both of the deepest branches of the mtDNA phylogeny, suggesting that the range of *D. recens* may have expanded out from this general region.

Our samples included three individuals of *D. subquinaria* that carried mtDNA typical of *D. recens* and thus were of hybrid ancestry (Figure 6). These flies included two from allopatric populations—Port Hardy, British Columbia (S-PH/BC-13 [13]), and Peachland, British Columbia (S-P/BC-14)—and one from a sympatric population, Edmonton, Alberta (S-E/AB-8). To verify the hybrid origin of these individuals, each was sequenced at two nuclear loci, *RIB* and *period*. All three of these individuals carried *D. subquinaria*-specific sequences at both loci, and none was infected with *Wolbachia*. These three individuals were excluded from our population genetic analyses. No individuals of *D. recens* (identified by male genitalia and/or PCR assay for *Wolbachia* infection) carried a *D. subquinaria* mtDNA haplotype, and we found no evidence of heteroplasmy (as indicated by multiple peaks on the chromatogram).

Table 2. Molecular differentiation within and among Populations of *D. recens* and *D. subquinaria*.

Species	Source of Variation	Fraction of Variation (%)	F
<i>D. recens</i>	Among groups: allopatric (five) versus sympatric (nine)	–1.04	$\Phi_{CT} = 0.010$ ($p = 0.662$)
	Among populations within groups	13.88	$\Phi_{SC} = 0.137$ ($p < 0.0001$)
	Within populations	87.16	$\Phi_{ST} = 0.128$ ($p < 0.0001$)
<i>D. subquinaria</i>	Among groups: allopatric (eight) versus sympatric (five)	5.02	$\Phi_{CT} = 0.050$ ($p = 0.172$)
	Among populations within groups	22.82	$\Phi_{SC} = 0.240$ ($p < 0.0001$)
	Within populations	72.16	$\Phi_{ST} = 0.278$ ($p < 0.0001$)

Number of populations within each group is shown in parentheses.
DOI: 10.1371/journal.pbio.0040325.t002

Discussion


Our field collections revealed a zone of geographical overlap between *D. recens* and *D. subquinaria* extending ~1,200 km from western Alberta to western Manitoba. Given the geological history of North America and the current ranges of these two species, it is likely that *D. recens* had been confined to eastern North America and *D. subquinaria* to western North America during the Wisconsin Glaciation and came into secondary contact when suitable habitat became connected across central Canada, i.e., within the last 12,000 y [37–39]. This zone of sympatry corresponds to Remington's suture zone 5 [40], which extends east from the eastern slope of the Canadian Rockies.

The principal aims of this study were to determine the following: (1) if sympatric populations of either *D. recens* or *D. subquinaria* exhibited reproductive character displacement; (2) if the species differed in the magnitude of this effect, in particular whether *D. subquinaria* not infected with *Wolbachia* exhibited greater reinforcement; (3) if there were any indirect effects of reinforcement on behavioral isolation within species; and (4) whether behavioral differentiation is simply

an indirect consequence of general genetic differentiation among populations.

Females from sympatric populations of *D. recens* were somewhat (but not significantly) less willing to mate with *D. subquinaria* males than were females from allopatric populations in both the mass matings and the no-choice experiments. However, in the no-choice experiments, the higher level of heterospecific mating by allopatric females of *D. recens* was due entirely to one strain, *R-R/NY* (Figure 3). Thus, although there is evidently genetic variation within *D. recens* in willingness to mate with males of *D. subquinaria*, this variation is only weakly correlated with the sympatric or allopatric status of the *D. recens* populations. Thus, there is little evidence for reinforcement in *D. recens*.

In contrast, we found a very high level of reproductive character displacement in *D. subquinaria*: allopatric females mated quite readily with *D. recens* males, but the sympatric females consistently refused to do so. This dramatic difference between sympatric and allopatric females was seen in two very different behavioral assays: a mass population encounter, as might occur in the vicinity of attractive mushrooms in the wild, and a no-choice assay designed to assess whether females would mate with heterospecific males

**Figure 6.** Minimum-Spanning Tree for *D. subquinaria* COI

Red dots are samples collected from the inland-sympatric populations, black dots represent inland-allopatric samples, and blue represent coastal-allopatric populations. The sizes of the circles indicate number of samples with each haplotype. The individuals identified as hybrids carry mtDNA haplotypes typical of *D. recens* but nuclear genes from *D. subquinaria*. Note the difference in scale between this figure and that for *D. recens* (Figure 7).
DOI: 10.1371/journal.pbio.0040325.g006




Figure 7. Minimum-Spanning Tree for *D. recens* COI

Red dots are samples collected from the sympatric populations, and black dots are samples collected from the allopatric populations. Haplotypes marked with an asterisk were found in the Smokies as well as other areas, and those marked with double asterisks were found only in the Smokies. The areas of the circles are proportional to number of samples with each haplotype. H1 is the most abundant haplotype ($n = 66$). Distance to outgroups can be seen in Figure 6. DOI: 10.1371/journal.pbio.0040325.g007

when no conspecific males were present. This finding is in contrast to the result of Coyne et al. [41], who showed that the degree of behavioral isolation between *D. yakuba* and *D. santomea* depended on the assay conditions, with isolation being significantly greater in choice assays. Our experiments show that the difference in mating behavior between sympatric and allopatric females of *D. subquinaria* is not context-dependent in the laboratory, suggesting that the difference is likely to be manifest under a variety of conditions in the wild.

To explore how *Wolbachia*-induced CI can lead to asymmetrical reinforcement between these two species, consider the following two possible initial crosses between them: (1) *D. subquinaria* female crossed with *D. recens* male and (2) *D. recens* female crossed with *D. subquinaria* male. For simplicity in what follows, assume that CI results in mortality of all offspring, although it is actually closer to 90% [13]. Cross 1 results in CI, causing an immediate loss of all alleles derived from both parental species. Cross 2, however, results in fertile hybrid females (hybrid males are sterile), which can mate with either *D. recens* or *D. subquinaria* males. Because these hybrids are infected with *Wolbachia*, backcross matings between hybrid females and *D. recens* males do not result in CI, thus allowing gene flow from the hybrids into *D. recens*. If hybrid females backcross to *D. subquinaria* males, the female descendants will be infected with *Wolbachia* and can continue to backcross to *D. subquinaria* males, because this is a compatible cross. However, the males produced in this backcross will be infected with *Wolbachia*, and their matings with *D. subquinaria* females (which are uninfected) will result in CI and thus the loss of genes that had been present in the original hybrids. In every backcross generation, half of the remaining nuclear genes that were present in the initial hybrids will segregate into males and thus be lost due to CI. Ultimately, all such nuclear genes will be lost in backcrosses to *D. subquinaria*.

The existence of unidirectional interspecific CI can also be inferred by patterns of mtDNA introgression between species. Because crosses between *D. subquinaria* females and

D. recens males result in CI, introgression of *D. subquinaria* mtDNA into *D. recens* is greatly hindered. In contrast, crosses between *D. recens* females and *D. subquinaria* males do not entail CI, and by backcrossing of the hybrid female to *D. subquinaria*, the mtDNA of *D. recens* can introgress into *D. subquinaria*. Thus, asymmetric introgression of mtDNA between species is an indication of both interspecific hybridization and interspecific CI in the wild. We found that three of 130 *D. subquinaria* individuals carried a *D. recens* mtDNA haplotype, whereas no individuals of *D. recens* have been found with a *D. subquinaria* haplotype ($n = 212$ individuals, including results reported here and those in Shoemaker et al. [13,31]). The difference between *D. recens* and *D. subquinaria* in the proportion of flies carrying heterospecific mtDNA is of borderline significance (Fisher's exact test, two-tailed $p = 0.054$).

One effect of this unidirectional interspecific CI is that nuclear genes can introgress from *D. subquinaria* into *D. recens*, but not vice versa. The model of Telschow et al. [42] can be applied to this situation. The effective rate of nuclear gene flow from an uninfected (*D. subquinaria*) to an infected (*D. recens*) population is reduced by a fraction $L/(1+L)$, where L is the level of CI [42]. Thus, if $L = 1$ (100% offspring mortality), then gene flow is 50% less than it would be in the absence of CI, the reduction being due to cross 1 above between uninfected females (*D. subquinaria*) and infected males (*D. recens*). In contrast, gene flow from the infected to the uninfected population is reduced by L . In other words, with complete CI ($L = 1$), there can be no introgression of genes from the infected into the uninfected species. Thus, the effect of CI is to allow some introgression of nuclear genes from *D. subquinaria* into *D. recens*, but none from *D. recens* into *D. subquinaria*.

How does unidirectional CI affect the evolution of mate choice? Suppose both species vary genetically in their level of discrimination against heterospecific males. Females of *D. subquinaria* that mate with *D. recens* males are likely to carry the less-discriminating alleles. However, because of CI, these alleles are transmitted to inviable progeny and are thereby selected out of the *D. subquinaria* population. In the reciprocal cross between *D. recens* females and *D. subquinaria* males, the less-discriminating alleles from *D. recens* are passed on to viable and fertile hybrid females. As discussed above, these alleles can then be crossed back into *D. recens*, but not into *D. subquinaria*. Thus, there is weaker selection against the less-discriminating alleles in *D. recens* than in *D. subquinaria*, the net effect being greater reinforcement in *D. subquinaria*.

There is an additional effect of the asymmetrical introgression discussed above. When *D. subquinaria* males and *D. recens* females mate and produce hybrid female offspring, these offspring are likely to carry alleles—which are transmitted by *D. subquinaria* males—that confer a female preference for *D. subquinaria* males. This is because the genes that affect mate choice by females may not have strong pleiotropic effects on the tendency of males to court and mate with various types of females, especially because males are the less-choosy sex in *Drosophila* [43]. Note that this is different from the effect of reduced female discrimination discussed above. Because genes can be introgressed from *D. subquinaria* to *D. recens* via these hybrid females, there can be introgression of *D. subquinaria*-preferring alleles into *D. recens*. Such introgression could potentially increase the tendency of

sympatric *D. recens* females to mate with *D. subquinaria* males. In contrast, CI prevents the reciprocal introgression of *D. recens*-preferring alleles into *D. subquinaria*. Thus, *D. subquinaria* experiences greater selection against nondiscriminating alleles (reinforcement) and is less subject to introgression of alleles conferring a preference for heterospecific males. Therefore, the much greater level of reproductive character displacement observed in *D. subquinaria* is consistent with the postulated effects of unidirectional CI imposed by *Wolbachia* infection in *D. recens*.

An alternate explanation for the greater reproductive character displacement exhibited by *D. subquinaria* is that the sympatric populations of this species are less inundated by gene flow from allopatric populations than are those of *D. recens*. Thus, the behavioral difference between sympatric and allopatric populations of *D. subquinaria* might simply be a consequence of general genetic divergence among these populations. However, we found essentially no genetic differentiation between inland-sympatric and inland-allopatric populations of *D. subquinaria*, and females from the inland-allopatric populations from Big Sky, Montana (S-BS/MT), and Coeur D'Alene, Idaho (S-CD/ID), exhibit the same low level of discrimination against *D. recens* males as do females from the allopatric coastal populations (Figure 3). Thus, reproductive character displacement is apparent even between inland-sympatric and inland-allopatric populations, which exhibit little or no mtDNA differentiation. This pattern is consistent with the behavioral patterns being a direct result of natural selection, due to unidirectional CI, rather than a secondary consequence of general genetic differentiation.

The occurrence of *D. subquinaria* individuals carrying mtDNA haplotypes characteristic of *D. recens* indicates that these species were not completely reproductively isolated when they came into secondary contact, as required by models of reinforcement [3]. Three lines of evidence indicate that these introgressions are not recent events. First, two of the three individuals of *D. subquinaria* that carried *D. recens* haplotypes were collected in allopatric populations several hundred kilometers away from the zone of sympatry, a signature of past introgression [44]. Second, none of the three individuals was infected with *Wolbachia*. Loss of infection is expected to result from imperfect maternal transmission, acting over many generations, coupled with a lack of transmission advantage of CI-causing *Wolbachia* when the prevalence of infection is low [45,46]. Finally, one of these haplotypes is identical to the very common *D. recens* haplotype H1, differing only in a synonymous C-to-T substitution at site 7, which may have occurred after introgression of this haplotype into *D. subquinaria*.

Although we have focused on the role that *Wolbachia* may play in the evolution of reinforcement in *D. subquinaria*, an open question is why *D. recens* shows little reproductive character displacement. We suggest three possibilities. First, general theoretical models of reinforcement [47] show that there can be a threshold level of selection against hybrids above which reinforcement occurs and below which it does not. It is thus possible that the extra selective burden imposed by CI results in a surpassing of the threshold for *D. subquinaria* but not for *D. recens*. Second, as discussed above, sympatric populations of *D. recens* may be subject to some introgression of alleles that confer a female mating preference for *D. subquinaria* males, and such introgression would act to reduce

reproductive character displacement. Finally, there may be some reproductive character displacement in *D. recens*, but our studies were not sensitive enough to detect it.

We found no post-mating isolation between the sympatric and allopatric populations of *D. subquinaria*, because both sexes of the F1 and F2 were viable and fertile in crosses between all pairwise combinations of populations. Although not tested yet, it is possible that more subtle forms of isolation operate between these populations, such as differential sperm precedence in multiply-mated females. In contrast, we did find substantial pre-mating behavioral isolation between *D. subquinaria* populations that are either sympatric or allopatric with *D. recens*. In our no-choice assays, we found that females of *D. subquinaria* from sympatric populations very often reject conspecific males from allopatric populations, both coastal and inland. In contrast, females from allopatric populations readily mate with conspecific males from both allopatric and sympatric populations. The finding that sympatric females discriminate against males from a genetically very similar inland-allopatric population suggests that reproductive character displacement in the sympatric populations, rather than general genetic differentiation, is responsible for the behavioral isolation among populations in *D. subquinaria*.

In general, behavioral isolation due to female mate choice can evolve as a result of changes in the mean male phenotype preferred by females or the range of male phenotypes against which females discriminate [25,48–50]. The pattern of behavioral isolation within *D. subquinaria* is not consistent with predictions of either a pure “mean preference” or pure “discrimination” model of behavioral isolation [25]. We propose the following simple model to explain the patterns of behavioral isolation observed in *D. subquinaria*. Suppose males of both *D. recens* and *D. subquinaria* display trait A, where A is a particular courtship song. As a result of selection to discriminate against males of *D. recens*, sympatric females of *D. subquinaria* accept only males that also display trait B, which may be a pheromone blend that distinguishes local (i.e., sympatric) *D. subquinaria* males from *D. recens*. Because there has been no selection for increased discrimination in allopatric populations, the display of male trait A is sufficient for acceptance by females of *D. subquinaria* from allopatric populations. If males from sympatric and allopatric populations of *D. subquinaria* differ in trait B, then these hypothetical mate preferences would yield the patterns of reinforcement and intraspecific behavioral isolation observed in *D. subquinaria*. This scenario is similar to the discrimination model of mate choice, because males displaying traits A and B are a subset of those displaying trait A.

Whatever the proximate cues may be, our data suggest that selection on *D. subquinaria* females to avoid *D. recens* males has indirectly brought about increased reproductive isolation between sympatric and allopatric populations of *D. subquinaria*. Because behavioral isolation is often the first step towards speciation in *Drosophila* [51], our results suggest that the presence of *Wolbachia* in one species may contribute to reproductive isolation within a different, uninfected species.

Materials and Methods

Geographical range and strain establishment. We surveyed many *Drosophila* populations ranging from British Columbia in the west to

Prince Edward Island in the east, including a number of sites where *D. recens* and *D. subquinaria* were hypothesized to be sympatric [13]. Flies were collected by sweep netting over mushrooms from 1995 to 2005, with most collections in 2002 (Table 1). Males were dissected and identified to species by examining their genitalia, which clearly differ between these two species [25]. Females of *D. recens* or *D. subquinaria* were placed in culture to examine their sons for species identification and to establish strains for the mating experiments. Following dissection (males) or culture (female), wild-caught flies were frozen for subsequent population genetic analysis.

Mating experiments—mass population. Genetically heterogeneous strains of each species were created by combining virgin males and females from several isofemale lines that originated from multiple locations within the allopatric and sympatric region. The sympatric populations for both *D. recens* and *D. subquinaria* included strains from Jasper, Edmonton, and The Pas (*R-J/AB*, *R/E-AB*, *R-P/MB*, *S-J/AB*, *S-E/AB*, and *S-P/MB*); the allopatric population of *D. subquinaria* included strains *S-V/BC*, *S-CF/MT*, *S-BS/MT*, and *S-CD/ID*; and the allopatric population for *D. recens* included strains *R-M/ND*, *R-B/MN*, *R-M/MI*, *R-W/ON*, *R-R/NY*, *R-B/NH*, *R-C/PEI*, and *R-SM/TN*.

The experimental flies were reared on Formula 4–24 instant *Drosophila* medium (Carolina Biological Supply, Burlington, North Carolina, United States) plus commercial *Agaricus bisporus* mushroom. Stocks were set up with a consistent starting density of adult flies to ensure a similar rearing density of larvae. Upon emergence as adults, flies were isolated on light CO₂ and kept as virgins on the same food at 22 °C with a 12-h light/dark cycle. The mating experiments were conducted when the males and females were 7 d post-emergence and only with flies with undamaged wings. All behavioral trials were conducted by aspirating flies into mating chambers, without the use of CO₂.

Three mass population trials were carried out, each using a different generation of flies. Each trial included two population cages run simultaneously, one to assess the mating preference of *D. recens* females and one for *D. subquinaria* females. Each trial began by transferring virgin females without anesthesia 1 h after lights-on to a population cage, which contained mushrooms placed on top of moistened paper towel. The mushrooms were intended to provide a rendezvous site for courting flies and mimicked natural situations where scores of flies can be found on and around a single attractive mushroom in the wild.

In the first cage, the females included 100 *D. subquinaria* from allopatric populations and 100 from sympatric populations. In a second cage, the females included 100 *D. recens* from allopatric populations and 100 from sympatric populations. To distinguish the sympatric from allopatric females, one type was marked lightly with fluorescent dust. Sympatric females were marked in trials 1 and 3, whereas the allopatric females were marked in trial 2. Males were added to each cage about 5 min after the females were, also without using anesthesia. Each cage received 100 *D. subquinaria* and 100 *D. recens* males, which included both sympatric and allopatric populations. Males were marked with fluorescent dust to distinguish the species, but not whether they were from allopatric or sympatric populations; *D. subquinaria* males were marked in trials 1 and 3, whereas *D. recens* males were marked in trial 2.

After addition of the males, the cages were observed continuously. Copulating pairs were removed by aspiration and immediately examined under ultraviolet light to identify them. We collected the first 100 mating pairs, or all mating pairs that mated in the first 2 h, whichever came first. All remaining flies were subsequently aspirated from the cage and identified under ultraviolet light.

Mating experiments—no choice. In contrast to the mass population experiment, a no-choice design allows the determination of whether females of one species will mate with males of another species when deprived of an opportunity to mate with their own species. Flies were reared exactly as in the mass population experiment. In the no-choice experiment, females from a genetically mixed strain from a single population were placed individually with males of the other species. For each species, we tested females from three allopatric populations and three sympatric populations. The allopatric populations of *D. subquinaria* used were strains *S-V/BC*, *S-CD/ID*, and *S-BS/MT*; the sympatric populations for both species were the same as those used in the mass population experiment; and the allopatric populations of *D. recens* included strains *R-R/NY*, *R-B/MN*, and *R-B/NH*. *D. subquinaria* females were paired with *D. recens* *R/E-AB* males, and each *D. recens* strain was paired with equal numbers of *D. subquinaria* *S-CD/ID*, *S-E/AB*, and *S-J/AB* males.

To test female mating behavior in a no-choice situation, one each of a 7-d-old female and male and were aspirated into a small vial (1

cm diameter × 3 cm long) containing blended mushroom-agar medium. The pairs consisted of either an allopatric or sympatric female of one species paired with a male of the other species. Three separate trials were conducted, and in each trial, we tested five females from each of three allopatric and three sympatric populations of each species. The number of females that mated within the first 3 h was counted.

Intraspecific mate choice—no choice. To assess whether there is any geographic differentiation in mate preference within species, we used the same methods as for the no-choice experiment, but paired males and females from the same species. For *D. recens*, we used flies from three sympatric (*R-E/AB*, *R-J/AB*, and *R-P/MB*) and three allopatric populations (*R-B/NH*, *R-B/MN*, and *R-R/NY*) in all pairwise combinations in two independent trials. For *D. subquinaria*, in the first two trials we used flies from two inland-sympatric (*S-E/AB* and *S-J/AB*), two inland-allopatric (*S-CD/ID* and *S-BS/MT*), and one coastal-allopatric population (*S-V/BC*). Two additional trials were carried out using three strains of *D. subquinaria*: combined *S-E/AB* and *S-J/AB* = *S-JE/AB* (inland-sympatric), *S-CD/ID* (inland-allopatric), and *S-V/BC* (coastal-allopatric). The use of *D. subquinaria* from these populations enabled us to assess whether intraspecific discrimination was associated with the inland-coastal genetic structure of *D. subquinaria* or the allopatric-sympatric distinction among genetically similar populations. The proportion of females that mated was analyzed via logistic regression as a function of trial, female population type, male population type, and interaction between the female and male population types.

Post-mating isolation within *D. recens* and *D. subquinaria*. Because we found evidence of behavioral isolation within *D. subquinaria*, we wished to determine whether taxonomically conspecific individuals from different populations exhibit post-mating isolation. Using the same genetically variable strains as for the intraspecific mate choice experiment, we did crosses of all reciprocal pairwise combinations (30 combinations within *D. recens* and 20 within *D. subquinaria*), each starting with three males and three females. For each, we then set up three crosses using the F1 offspring and, subsequently, three crosses using the F2 offspring, where each cross included three males and three females. For each of the F1 and F2 crosses, we scored whether the flies produced viable offspring, and in this manner, we were able to assess the viability and fertility of F1 and F2 offspring produced by inter-population crosses.

Population structure. To assay differentiation among populations as well as the frequency of hybrids, we sequenced 1,432 base pairs of the mtDNA *Cytochrome Oxidase I* (*COI*) gene. Although the mtDNA of *D. recens* is known to harbor low polymorphism [31], there is no evidence that *COI* is currently experiencing a selective sweep, and thus the present polymorphism should reflect recent patterns of gene flow among populations. We sequenced *COI* from wild-caught individuals or single flies from isofemale lines established from wild flies of *D. subquinaria* and *D. recens*. Some of our samples were obtained from isofemale lines established from wild-caught females. In these cases, species identifications were made by dissection and examination of male genitalia, which show diagnostic differences between *D. recens* and *D. subquinaria* [35]. Some wild-caught individuals were frozen before dissection; to provisionally identify these flies, we conducted a PCR assay to test for *Wolbachia* infection (using the primers wsp81F and wsp691R from Zhou et al. [52]). Individuals that tested positive for *Wolbachia* were tentatively identified as *D. recens*, and uninfected flies as *D. subquinaria*. Because about 2% of wild-caught *D. recens* are uninfected due to imperfect maternal transmission of *Wolbachia* [13], the few uninfected flies that had a *D. recens* *COI* haplotype were also sequenced at a nuclear marker (as in [31]) to determine whether they were uninfected *D. recens* or a *D. subquinaria* that carried a *D. recens* *COI* haplotype and thus of hybrid origin. Samples which carried a *D. recens*-type *COI* but a *D. subquinaria*-type nuclear marker were sequenced at a second nuclear marker to confirm species identity. Nuclear markers used were *period* and *R1B*. In all three of these cases, both nuclear markers were of *D. subquinaria* type but the individual carried a *D. recens* type mtDNA, indicative of a historical introgression of the mtDNA.

We used standard techniques for all PCR and DNA sequencing (Protocol S1). Sequences are deposited in GenBank and were combined with previously published sequences.

Within each species, we tested for differentiation between sympatric and allopatric regions and for differentiation between all pairs of populations. We used variants of *F_{ST}* statistics, as implemented in DnaSP 4.0 [53] (Protocol S1). In addition, we partitioned variation within and among sympatric and allopatric regions using an analysis of molecular variance, as implemented in Arlequin [54].

To visualize relationships and relative abundances of mtDNA haplotypes within each species, a network was created in SplitsTree4 [55] (available from <http://www.splitsree.org>). A network was created for *D. recens* and *D. subquinaria* separately, using *D. quinaria* and one sample the other species as outgroups. For the networks shown in Figures 6 and 7, we used a neighbor-joining algorithm and Kimura 2-parameter distance estimates.

Supporting Information

Protocol S1. Population Structure: Methods and Additional Results

Found at DOI: 10.1371/journal.pbio.0040325.sd001 (133 KB DOC).

Accession Numbers

The GenBank (<http://www.ncbi.nlm.nih.gov/Genbank>) accession numbers for sequences discussed in the paper are nuclear markers *period* and *R1B* in *D. recens* (AF248076 and DQ494185–DQ494192), nuclear markers *period* and *R1B* in *D. subquinaria* (DQ494178–DQ494184), and mtDNA *Cytochrome Oxidase I (COI)* from *D. recens*, *D. subquinaria*, and *D. quinaria* (AY154400–AY154457 and DQ851591–DQ851807).

References

- Dobzhansky Th (1937) Genetics and the origin of species. New York: Columbia University Press. 364 p.
- Blair WF (1955) Mating calls and stage of speciation in the *Microhylla olivacea* – *M. carolinensis* complex. *Evolution* 9: 469–480.
- Butlin RK (1987) Speciation by reinforcement. *Trends Ecol Evol* 2: 8–13.
- Servedio MR, Noor MAF (2003) The role of reinforcement in speciation: Theory and data. *Ann Rev Ecol Syst* 34: 339–364.
- Coyne JA, Orr HA (2004) Speciation. Sunderland (Massachusetts): Sinauer Associates. 545 p.
- Rundle HD, Schluter D (1998) Reinforcement of stickleback mate preferences: Sympatry breeds contempt. *Evolution* 52: 200–208.
- Geyer LB, Palumbi SR (2003) Reproductive character displacement and the genetics of gamete recognition in tropical sea urchins. *Evolution* 57: 1049–1060.
- Noor MAF (1995) Speciation driven by natural selection in *Drosophila*. *Nature* 375: 674–675.
- Hoffmann AA, Turelli M (1997) Cytoplasmic incompatibility in insects. In: O'Neill SL, Hoffmann AA, Werren JH, editors. *Influential passengers: Inherited microorganisms and arthropod reproduction*. Oxford: Oxford University Press. pp. 42–80.
- Stouthamer R, Breeuwer JAJ, Hurst GDD (1999) *Wolbachia pipiens*: Microbial manipulator of arthropod reproduction. *Ann Rev Micro* 53: 71–102.
- Werren JW, Windsor DM (2000) *Wolbachia* infection frequencies in insects: Evidence of a global equilibrium? *Proc R Soc Lond B Biol Sci* 267: 1277–1285.
- West SA, Cook JM, Werren JH, Godfray HCJ (1998) *Wolbachia* in two insect host–parasitoid communities. *Mol Ecol* 11: 1457–1465.
- Shoemaker DD, Katju V, Jaenike J (1999) *Wolbachia* and the evolution of reproductive isolation between *Drosophila recens* and *Drosophila subquinaria*. *Evolution* 53: 1157–1164.
- Shoemaker DD, Machado CA, Molbo D, Werren JH, Windsor DM, Herre EA (2002) The distribution of *Wolbachia* in fig wasps: Correlations with host phylogeny, ecology and population structure. *Proc R Soc Lond B Biol Sci* 269: 2257–2267.
- Breeuwer JAJ, Werren JH (1990) Microorganisms associated with chromosome destruction and reproductive isolation between two insect species. *Nature* 346: 558–560.
- Bordenstein SR, O'Hara FP, Werren JH (2001) *Wolbachia*-induced incompatibility precedes other hybrid incompatibilities in *Nasonia*. *Nature* 409: 707–710.
- Werren JH (1998) *Wolbachia* and speciation. In: Howard D, Berlocher S, editors. *Endless forms: Species and speciation*. Oxford: Oxford University Press. pp. 245–260.
- Bordenstein SR (2003) Symbiosis and the origin of species. In: Bourtzis K, Miller T, editors. *Insect symbiosis*. New York: CRC Press. pp. 283–304.
- Weeks AR, Reynolds KT, Hoffmann AA (2002) *Wolbachia* dynamics and host effects: What has (and has not) been demonstrated? *Trends Ecol Evol* 17: 257–262.
- Caspari E, Watson GS (1959) On the evolutionary importance of cytoplasmic sterility in mosquitoes. *Evolution* 13: 568–570.
- Turelli M (1994) Evolution of incompatibility-inducing microbes and their hosts. *Evolution* 48: 1500–1513.
- Rousset F, Raymond M, Kjellberg F (1991) Cytoplasmic incompatibilities in the mosquito *Culex pipiens*: How to explain a cytotype polymorphism. *J Evol Biol* 4: 69–81.
- Turelli M, Hoffmann AA (1991) Rapid spread of an inherited incompatibility factor in California *Drosophila*. *Nature* 353: 440–442.

Acknowledgments

We thank Andrea Betancourt, Mohamed Noor, Allen Orr, Howard Rundle, Arndt Telschow, Thomson Webb, Jack Werren, members of the Charlesworth lab group, and two anonymous reviewers for helpful discussions and/or comments on the manuscript. Andy Beckenbach, Paul Gibas, Jessica Savage, and DeWayne Shoemaker assisted with fly collecting and Julie Sullivan and Karin Tetzlaff assisted with the laboratory experiments. *Drosophila* were imported from Canada under United States Department of Agriculture permit 56887 to JJ and were collected in the Smoky Mountains under a National Park collecting permit to KAD.

Author contributions. JJ conceived the project. JJ and KAD designed the experiments. JJ, KAD, CC, and MSM performed the experiments. JJ and KAD analyzed the data. JJ and KAD wrote the paper.

Funding. This research was supported by grants from the US National Science Foundation (DEB-0315521 and EF-0328363) to JJ, a Royal Society USA Fellowship to KAD, and a DeKiewiet fellowship to CC.

Competing interests. The authors have declared that no competing interests exist.

- Telschow A, Hammerstein P, Werren JH (2005) The effect of *Wolbachia* versus genetic incompatibilities on reinforcement and speciation. *Evolution* 59: 1607–1619.
- Kelly JK, Noor MAF (1996) Speciation by reinforcement: A model derived from studies of *Drosophila*. *Genetics* 143: 1485–1497.
- Littlejohn MJ, Loftus-Hills JJ (1968) An experimental evaluation of premating isolation in the *Hyla ewingi* complex (Anura: Hylidae). *Evolution* 22: 659–663.
- Hoskin CJ, Higgie M, McDonald KR, Moritz C (2005) Reinforcement drives rapid allopatric speciation. *Nature* 437: 1353–1356.
- Zouros E, D'Entremont CJ (1980) Sexual isolation among populations of *Drosophila-mojavensis*: Response to pressure from a related species. *Evolution* 34: 421–430.
- Werren JW, Jaenike J (1995) *Wolbachia* and cytoplasmic incompatibility in mycophagous *Drosophila* and their relatives. *Heredity* 75: 320–326.
- Shoemaker DD, Dyer KA, Ahrens M, McAbee K, Jaenike J (2004) Molecular evolutionary effects of *Wolbachia* infections: Decreased diversity but increased substitution rate in host mtDNA. *Genetics* 168: 2049–2058.
- Hurst GDD, Jiggins FM (2005) Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: The effects of inherited symbionts. *Proc R Soc Lond B Biol Sci* 272: 1525–1534.
- Rasgon JL, Corneli AJ, Scott TW (2006). Evolutionary history of a mosquito endosymbiont revealed through mitochondrial hitchhiking. *Proc R Soc Lond B Biol Sci* 273: 1603–1611.
- Rokas A (2000) *Wolbachia* as a speciation agent. *Trends Ecol Evol* 15: 44–45.
- Wheeler MR (1960) New species of the quinaria group of *Drosophila* (Diptera, Drosophilidae). *Southwestern Naturalist* 5: 160–164.
- Pfennig KS, Ryan MJ (2006) Reproductive character displacement generates reproductive isolation among conspecific populations: An artificial neural network study. *Proc R Soc Lond B Biol Sci* 273: 1361–1368.
- Carstens BC, Brunsfeld SJ, Demboski JR, Good JM, Sullivan J (2005) Investigating the evolutionary history of the Pacific Northwest mesic forest ecosystem: Hypothesis testing within a comparative phylogeographic framework. *Evolution* 59: 1639–1652.
- Love D (1959) The postglacial development of the flora of Manitoba: A discussion. *Can J Botany* 37: 547–585.
- Prentice IC, Bartlein PJ, Webb T (1991) Vegetation and climate change in eastern North America since the last glacial maximum. *Ecology* 72: 2038–2056.
- Williams JW, Shuman BN, Webb T, Bartlein PJ, Leduc PL (2004) Late Quaternary vegetation dynamics in North America: Scaling from taxa to biomes. *Ecol Monographs* 74: 309–334.
- Remington CL (1968) Suture-zones of hybrid interaction between recently joined biotas. *Evol Biol* 2: 321–428.
- Coyne JA, Elwyn S, Rolan-Alvarez E (2005) Impact of experimental design on *Drosophila* sexual isolation studies: Direct effects and comparison to field hybridization data. *Evolution* 59: 2588–2601.
- Telschow A, Hammerstein P, Werren JH (2002) The effect of *Wolbachia* on genetic divergence between populations: Models with two-way migration. *Am Nat* 160: S54–S66.
- Bateman AJ (1948) Intra-sexual selection in *Drosophila*. *Heredity* 2: 349–368.
- Barton NH, Hewitt GM (1989) Adaptation, speciation, and hybrid zones. *Nature* 341: 497–503.
- Caspari E, Watson GS (1959) On the evolutionary importance of cytoplasmic sterility in mosquitoes. *Evolution* 13: 568–570.
- Turelli M (1994) Evolution of incompatibility-inducing microbes and their hosts. *Evolution* 48: 1500–1513.
- Servedio MR (2000) Reinforcement and the genetics of nonrandom mating. *Evolution* 54: 21–29.

48. Lande R (1981) Models of speciation by sexual selection on polygenic traits. *Proc Natl Acad Sci U S A* 78: 3721–3725.
49. Kaneshiro KY (1989) The dynamics of sexual selection and founder effects in species formation. In: Giddings VL, Kaneshiro KY, Anderson WW, editors. *Genetics, speciation, and the founder principle*. New York: Oxford University Press. pp 279–296.
50. Butlin RK (1993) The variability of mating signals and preferences in the brown planthopper, *Nilaparvata lugens* (Homoptera, Delphacidae). *J Insect Behavior* 6: 125–140.
51. Coyne JA, Orr HA (1997) “Patterns of speciation in *Drosophila*” revisited. *Evolution* 51: 295–303.
52. Zhou WG, Rousset F, O'Neill S (1998) Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. *Proc R Soc Lond B Biol Sci* 265: 509–515.
53. Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R (2003) DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 19: 2496–2497.
54. Schneider S, Roessli D, Excoffier L (2000) Arlequin version 2000: Software for population genetic analyses. Genetics and Biometry Laboratory, University of Geneva, Switzerland.
55. Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol* 23: 254–267.