

Organism	Body Length* (cm)	SW	Source
<i>Dasyatis sabina</i>	20.9	17.78	[3]
<i>Dasyatis say</i>	23.3	17.68	[3]
<i>Dasyatis americana</i>	27.7	25.09	[3]
<i>Gymnura micrura</i>	16	20.21	[3]
<i>Rhinoptera bonasus</i>	28.7	19.43	[3]
<i>Dasyatis violecea</i>	55	24.13	[3]
<i>Raja eglanteria</i>	22	19.59	[3]
<i>Taenuira lymma</i>	16.4	20.5	[3]
<i>Potamotrygon orbignyi</i>	12.8	24.03	[5]
<i>Gymnarchus niloticus</i>	24.5	19.02	[6, 7]
<i>Apteronotus albifrons</i>	14.2	18.03	[10]
<i>Rhinecanthus aculeatus</i>	9.79	18.1	[8]
<i>Regalecaus glesne</i>	255	22.47	[12]
<i>Apteronotus leptorhynchus</i>	15.5	19.81	[13]
<i>Gymnotus carapo</i>	20	22.45	[13]
<i>Gymnorhamphichthys hypostomus</i>	17.8	22.81	[13]
<i>Eigenmannia virescens</i> [‡]	11.59	17.7	[2]
<i>Xenomystus nigri</i>	15	15.05	[13]
<i>Amia calva</i>	30.1	15.68	[11]
<i>Pseudobiceros bedfordi</i> [†]	8 – 10	15.37	This Study
<i>Pseudobiceros pardalis</i> [†]	2 – 5	15.17	This Study
<i>Sepia officinalis</i> [†]	30 – 45	18.87	This Study

*Disc length in case of batoid fishes.

[†]Approximate range of body length (mantle length in case of *Sepia officinalis*).

[‡]Mean SW of forward and backward counterpropagating waves.