

S2 Table. Relative transcript level of *TOGR1* in 38 rice accessions.

Name	Original producing area	Mean transcript level of <i>TOGR1</i>	S.E.	Mean plant height (cm)	Type
Wuke-4	China	1.00	0.0307	107.5	<i>indica</i>
Hongmangnuo	China	1.13	0.0855	153.7	<i>temperate japonica</i>
Lijianghei	China	0.72	0.038	165	<i>temperate japonica</i>
Maguzhi	China	0.63	0.0261	155	<i>temperate japonica</i>
Nipponbare	Japan	0.59	0.1018	99.8	<i>temperate japonica</i>
Yandao8	China	0.06	0.0013	67.4	<i>temperate japonica</i>
Zhonghua11	China	0.79	0.06	112	<i>temperate japonica</i>
O. rufipogon	China	1.14	0.1198	125	<i>rufipogon</i>
9311	China	0.76	0.0324	110	<i>indica</i>
Baipigu	China	1.13	0.0494	160	<i>indica</i>
Basmati113	India	1.09	0.059	170	<i>indica</i>
Benguzhong	China	0.27	0.0305	125	<i>indica</i>
Changainuo	China	0.23	0.0317	107.4	<i>indica</i>
Dabaihangu	China	1.44	0.0476	178	<i>indica</i>
Dahongjiao-3	China	0.71	0.0129	163	<i>indica</i>
Danuo	China	0.88	0.0567	118	<i>indica</i>
Dayemaozhan	China	0.91	0.0384	92	<i>indica</i>
Erpozao	China	0.76	0.0106	93	<i>indica</i>
Hanzhangu	China	0.31	0.0069	129.6	<i>indica</i>
Hanzhangu	China	0.59	0.0189	129.6	<i>indica</i>
Hongjiaowen	China	0.4	0.0145	154.8	<i>indica</i>
IR24	Philippine	0.97	0.0591	105	<i>indica</i>
Liuyuegu	China	1.04	0.0913	87	<i>indica</i>
Maweibaimizhan	China	0.72	0.0329	179.8	<i>indica</i>
Maweixian	China	0.76	0.1124	143	<i>indica</i>
Nanjing-6	China	0.68	0.0125	151	<i>indica</i>
Nantehao	China	1.17	0.014	160	<i>indica</i>
Xiangzhou-4	China	1.26	0.0538	90	<i>indica</i>
Yangguzi-2	China	0.98	0.0977	156	<i>indica</i>
Yanzhan	China	1.78	0.0549	166	<i>indica</i>
Yanzhanzhong	China	0.78	0.0337	140.5	<i>indica</i>
Yezhugu	China	1.34	0.1526	168	<i>indica</i>
Zaobaigu	China	0.84	0.0718	162.2	<i>indica</i>
Zhefu-802	China	0.74	0.0582	80	<i>indica</i>
Zhenzhunuo	China	1.22	0.0891	133.2	<i>indica</i>
O.granulata I	China	0.027	0.0017	45.9	<i>granulata</i>
O.granulata II	China	0.03	0.0016	79.3	<i>granulata</i>
Dular	India	1.27	0.0312	165	<i>aus</i>