

RESEARCH ARTICLE

# The natural anti-tumor compound Celastrol targets a Myb-C/EBP $\beta$ -p300 transcriptional module implicated in myeloid gene expression

Anna Coulibaly<sup>1‡</sup>, Astrid Haas<sup>1</sup>, Simone Steinmann<sup>1</sup>, Anke Jakobs<sup>1</sup>, Thomas J. Schmidt<sup>2</sup>, Karl-Heinz Klempnauer<sup>1\*</sup>

**1** Institute for Biochemistry, Westfälische-Wilhelms-Universität, D-48149 Münster, Germany, **2** Institute for Pharmaceutical Biology and Phytochemistry, Westfälische-Wilhelms-Universität, D-48149 Münster, Germany

‡ Current address: Universitätsmedizin Mannheim, Medizinische Fakultät Mannheim der Universität Heidelberg, Klinik für Anästhesiologie und Operative Intensivmedizin, Theodor-Kutzer-Ufer 1–3, D-68167 Mannheim, Germany

\* [klempna@uni-muenster.de](mailto:klempna@uni-muenster.de)



**OPEN ACCESS**

**Citation:** Coulibaly A, Haas A, Steinmann S, Jakobs A, Schmidt TJ, Klempnauer K-H (2018) The natural anti-tumor compound Celastrol targets a Myb-C/EBP $\beta$ -p300 transcriptional module implicated in myeloid gene expression. PLoS ONE 13(2): e0190934. <https://doi.org/10.1371/journal.pone.0190934>

**Editor:** Kevin D. Bunting, Emory University, UNITED STATES

**Received:** November 1, 2017

**Accepted:** December 22, 2017

**Published:** February 2, 2018

**Copyright:** © 2018 Coulibaly et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper.

**Funding:** This work was supported by the Deutsche Krebshilfe ([www.krebshilfe.de](http://www.krebshilfe.de)), grant number 109269, Deutsche Forschungsgemeinschaft ([www.dfg.de](http://www.dfg.de)), grant number KL461/17-1, and the Open Access Publication Fund of the University of Münster. The funders had no role in study design, data collection

## Abstract

Myb is a key regulator of hematopoietic progenitor cell proliferation and differentiation and has emerged as a potential target for the treatment of acute leukemia. Using a myeloid cell line with a stably integrated Myb-inducible reporter gene as a screening tool we have previously identified Celastrol, a natural compound with anti-tumor activity, as a potent Myb inhibitor that disrupts the interaction of Myb with the co-activator p300. We showed that Celastrol inhibits the proliferation of acute myeloid leukemia (AML) cells and prolongs the survival of mice in an in vivo model of AML, demonstrating that targeting Myb with a small-molecule inhibitor is feasible and might have potential as a therapeutic approach against AML. Recently we became aware that the reporter system used for Myb inhibitor screening also responds to inhibition of C/EBP $\beta$ , a transcription factor known to cooperate with Myb in myeloid cells. By re-investigating the inhibitory potential of Celastrol we have found that Celastrol also strongly inhibits the activity of C/EBP $\beta$  by disrupting its interaction with the Taz2 domain of p300. Together with previous studies our work reveals that Celastrol independently targets Myb and C/EBP $\beta$  by disrupting the interaction of both transcription factors with p300. Myb, C/EBP $\beta$  and p300 cooperate in myeloid-specific gene expression and, as shown recently, are associated with so-called super-enhancers in AML cells that have been implicated in the maintenance of the leukemia. We hypothesize that the ability of Celastrol to disrupt the activity of a transcriptional Myb-C/EBP $\beta$ -p300 module might explain its promising anti-leukemic activity.

and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

The transcription factor Myb plays a key role as a regulator of proliferation and differentiation of hematopoietic progenitor cells and has been implicated in the development of acute leukemia [1]. Genomic rearrangements of the *MYB* gene and mutations that create de-novo Myb binding sites in the transcriptional control region of the *TAL1* oncogene have been detected in acute lymphoid leukemia [2–4]. Furthermore, acute myeloid leukemia (AML) cells are addicted to high levels of Myb expression, which makes them more vulnerable to Myb inhibition than their normal counterparts [5–7]. Recent genome-wide studies have identified super-enhancers, i.e. cis-regulatory regions that are significantly larger than typical enhancers and that drive the expression of genes involved in cell identity and disease [8–10]. In AML cells, super-enhancers are densely loaded with chromatin regulators such as p300 and BRD4 and hematopoietic transcription factors, such as Myb, C/EBP $\alpha$ , C/EBP $\beta$ , ERG, FLI1, and PU.1 [11]. These studies have confirmed the important role of Myb and C/EBPs for the proliferation of AML cells, suggesting a mechanistic framework for the addiction of AML cells to high levels of Myb expression. Targeting Myb with small-molecule inhibitors therefore offers potential for the development of novel therapeutic strategies for the treatment of AML and other tumors with deregulated Myb, such as adenoid cystic carcinoma or specific subtypes of glioblastoma [12,13].

In myeloid cells Myb cooperates with members of the C/EBP family, basic-region-leucine-zipper (bzip) transcription factors that are involved in the differentiation and proliferation of various cell types [14]. C/EBP activity is controlled by various post-translational modifications [15–20] and by the expression of alternative isoforms based on translation initiation at distinct start codons [21,22].

We have previously established a myeloid reporter cell line as a screening tool to search for low molecular weight Myb inhibitors [23]. These cells carry a stably integrated Myb-responsive reporter gene driven by the promoter and enhancer of the myeloid-specific chicken *mim-1* gene, a Myb target gene whose expression is strongly induced by Myb in myeloid cells. Using these cells we have identified and characterized several compounds as potential Myb inhibitors [24–27]. One of these compounds, the pentacyclic triterpenoid Celastrol, was shown to inhibit Myb by disrupting the interaction of Myb transactivation domain with the Kix domain of the co-activator p300 [25]. Importantly, we also showed that Celastrol inhibits the proliferation of primary mouse AML cells and leukemic cells from AML patients while the proliferation of normal murine or human early hematopoietic progenitor cells was not affected [25]. Furthermore, in vivo experiments showed that Celastrol significantly prolonged the survival of mice in a model of an aggressive AML [25]. Taken together, these findings demonstrated that targeting of Myb by a small-molecule inhibitor is feasible and might have therapeutic potential for the treatment of AML.

We recently became aware that our cell-based screening assay also responds to inhibition of C/EBP $\beta$ , as exemplified by the sesquiterpene lactone helenalin acetate, a compound that inhibits Myb-induced *mim-1* expression but was shown to be a potent inhibitor of C/EBP $\beta$  instead of a Myb inhibitor [28,29]. We showed before that Myb requires C/EBP family members, such as C/EBP $\alpha$  or C/EBP $\beta$ , as essential cooperation partners to stimulate the *mim-1* gene [30,31], explaining why inhibition of C/EBP $\beta$  has a similar effect as the inhibition of Myb in our screening system. We have therefore re-investigated the inhibitory mechanism of Celastrol to know if Celastrol also exerts inhibitory effects on C/EBP $\beta$ . Surprisingly, our data show that Celastrol is also a highly potent inhibitor of C/EBP $\beta$  suggesting that the promising inhibitory effects of Celastrol on AML cells might be due to its ability to simultaneously target Myb and C/EBP $\beta$ .

## Materials and methods

### Cells

The quail fibroblast cell line QT6 was originally obtained from P. Vogt (La Jolla). 3T3-L1 cells were obtained from ATCC 5 year ago and were used within 10 passages from thawing. All cells were tested on a regular basis and were free of mycoplasma contamination. Adipocyte differentiation was induced as published [32]. Celastrol was from Sigma-Aldrich, dissolved in DMSO and stored at 10 mM stock solution at -70°C.

### Transfections

QT6 fibroblasts were transfected by calcium-phosphate co-precipitation, followed by analysis of the transfected cells for mRNA expression, protein interaction or reporter gene activities. A constant amount of the  $\beta$ -galactosidase reporter plasmid pCMV $\beta$  was always included, allowing transfection efficiencies to be normalized according to the  $\beta$ -galactosidase activities. Reporter gene activities were analyzed as described [30,33]. The luciferase reporter genes pG5E4-38Luc and p-240Luc contain 5 tandem copies of Gal4-binding sites or the promoter of the chicken *mim-1* gene, respectively. Luciferase values were divided by the  $\beta$ -galactosidase activity to compensate different transfection efficiencies. All reporter studies were performed in at least 3 independent experiments, with replicate transfections in each experiment. Northern blotting was performed as described [30,33]. Blots were hybridized sequentially with radioactive probes corresponding to the C/EBP $\beta$ -inducible MRP126 mRNA and the ribosomal protein S17 mRNA (as housekeeping control). Radioactive bands were detected and quantified using a phosphor image analyzer. Electrophoretic mobility shift assays (EMSA) were performed as described [34]. Briefly, a pair of complementary single-stranded oligonucleotides containing a consensus C/EBP binding site was annealed and used for gel retardation assays: 5' -TG TAGCTGCAGATTGCGCAATCTGCATCTA-3' and 5' -GTAGATGCAGATTGCGCAATCTGCAGCTACA-3'. After annealing, oligonucleotides were radiolabeled by filling-in the ends using  $\alpha^{32}\text{P}$ -dCTP and Klenow polymerase. Nuclear extract from QT6 cells transfected with C/EBP $\beta$  expression vector and treated with different concentrations of Celastrol was prepared as described [34].

### Expression vectors and transfections

Expression vectors for chicken and mouse C/EBP $\beta$ , a C/EBP $\beta$  mutant lacking all cysteine residues (CallA) and chicken C/EBP $\alpha$  have been described [16,29,30,35]. Expression vectors for full-length human p300 and truncated p300 constructs p300/1751-2370 and p300/1751-1947 have been described [33]. Expression vectors for the p300/1751-1947 mutations C1789A, C1790A and C1789,1790A (CC) were generated by oligonucleotide-directed mutagenesis. Plasmids encoding Gal4-C/EBP $\beta$ , p300-VP16, Gal4-VP16 and C/EBP $\beta$ -YFP fusion proteins were described before [29,33].

### GFP-trap

QT6 fibroblasts expressing YFP or C/EBP $\beta$ -YFP were lysed 16 hours post transfection in ELB buffer containing 120mM sodium chloride, 50mM Tris/HCL, pH 7.4, 20mM sodium fluoride, 1mM EDTA, 6mM EGTA, 15mM sodium pyrophosphate, 1mM Phenylmethylsulfonyl Fluoride (PMSF), 0.5% NP-40 and centrifuged for 20 min at 14,000 x g. An aliquot of the supernatant was retained as input control. The remaining supernatant was incubated with GFP-trap beads (Chromotec, München) for 3 hours at 4°C. Beads were washed three times with ELB

buffer, boiled in sodium dodecyl sulfate (SDS) sample buffer and analyzed together with the input samples by SDS-PAGE and western blotting, using appropriate antibodies.

## Results

### Celastrol is a potent inhibitor of C/EBP $\beta$


To investigate if Celastrol previously identified as a Myb inhibitor also affects the activity of C/EBP $\beta$  we performed luciferase assays with a reporter construct that is driven the promoter of the *mim-1* gene, which contains high affinity C/EBP binding sites and is strongly activated by C/EBP $\beta$  [36]. These experiments clearly demonstrated that Celastrol strongly inhibits the activity of C/EBP $\beta$  (Fig 1A). The amount of C/EBP $\beta$  was not decreased by Celastrol, indicating that the activity and not the expression of C/EBP $\beta$  was inhibited. Importantly, because the reporter assays were performed in fibroblasts that do not express Myb, the inhibition of C/EBP $\beta$  activity by Celastrol is independent of its previously described Myb-inhibitory activity [25]. Hence, we concluded that Celastrol inhibits both Myb and C/EBP $\beta$  independently of each other.

To demonstrate under more physiological conditions that Celastrol inhibits C/EBP $\beta$ , we analyzed its effect on the transcriptional activation of an endogenous C/EBP target gene that is silent in fibroblasts but can be activated by ectopic expression of C/EBP $\beta$  or C/EBP $\alpha$  [15,33], thereby providing a read-out of C/EBP activity at a chromatin-embedded gene. Fig 1B shows that Celastrol strongly inhibited the stimulation of MRP126 expression by C/EBP $\beta$ , confirming that Celastrol is a potent inhibitor of C/EBP $\beta$ . Mouse (Fig 1C) and human C/EBP $\beta$  were also strongly inhibited by Celastrol. Interestingly, while Celastrol erased C/EBP $\beta$  activity virtually completely at the concentrations tested the related transcription factor C/EBP $\alpha$  was inhibited only slightly under the same conditions (Fig 1D).

To demonstrate the inhibitory activity of Celastrol on C/EBP $\beta$  in a different biological context we employed the mouse pre-adipocyte cell-line 3T3-L1, whose differentiation into adipocytes is an established model of a C/EBP $\beta$ -dependent differentiation process. In the 3T3-L1 cells, the initial activation of C/EBP $\beta$  induces a cascade of gene activations ultimately leading to a fat cell phenotype that is characterized by an accumulation of lipid droplets in the cytoplasm of the cells [32]. Because C/EBP $\beta$  plays a key role during the initial phase of adipocyte differentiation we expected Celastrol to inhibit the differentiation of these cells. Fig 1E shows that the massive differentiation observed in the absence of Celastrol was strongly suppressed by Celastrol as demonstrated by microscopic analysis and staining with the lipid dyes Oil Red O and Nile Red. Consistent with previous work [32], as shown by the western blots at the bottom right side of Fig 1E, the amount of C/EBP $\beta$  increased when differentiation was induced, but was not significantly diminished by Celastrol, demonstrating that Celastrol inhibits the activity and not the expression of C/EBP $\beta$ . Together, the data in Fig 1 show that Celastrol is a very potent inhibitor of C/EBP $\beta$ .

### Celastrol disturbs the interaction of C/EBP $\beta$ and p300

To investigate if Celastrol inhibits C/EBP $\beta$  DNA-binding activity we performed electrophoretic mobility shift assays (EMSA) with nuclear extracts from cells transfected with C/EBP $\beta$  expression vector and treated with or without Celastrol. Fig 2A shows that the DNA-binding activity of C/EBP $\beta$  was not significantly affected by Celastrol. As can be seen in the lower part of Fig 2A, complexes of C/EBP $\beta$  bound to the radiolabeled oligonucleotide were not significantly reduced by the treatment of the cells with Celastrol. Similarly, the amount of C/EBP $\beta$  present in the nuclear extract was not significantly affected by the Celastrol treatment (see western blot at the top of Fig 2A). That Celastrol did not inhibit C/EBP $\beta$  activity by interfering




**Fig 1. Celastrol inhibits C/EBP $\beta$  activity.** **A.** QT6 fibroblasts were co-transfected with the luciferase reporter gene p-240luc containing the C/EBP-inducible promoter of the chicken *mim-1* gene, the  $\beta$ -galactosidase expression vector pCMV $\beta$  and an expression vector for C/EBP $\beta$  transfection, the cells were treated with Celastrol as indicated and harvested after 16 hours. The columns show the average luciferase activity normalized to the  $\beta$ -galactosidase activity. Thin lines show standard deviations. Asterisks indicate statistical significance (\*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ , Student's t-test). C/EBP $\beta$  expression is shown at the bottom. **B-D.** QT6 cells were transfected with expression vectors for chicken or mouse full-length C/EBP $\beta$  or chicken C/EBP $\alpha$  and treated with Celastrol as indicated. Cells were harvested after 16 hours and analyzed by northern blotting for expression of *MRP126* and *S17* mRNAs (top and middle panel) and by western blotting for the expression of C/EBP $\beta$  or C/EBP $\alpha$  (bottom panels). **E.** 3T3-L1 cells were induced to undergo adipocyte differentiation for 7 days in the absence or presence of the indicated concentrations of Celastrol. The top and middle panels show microscopic pictures of the cells at low magnification and after staining with the fluorescent lipid dye Nile Red. The bottom panels show staining of the cells with Oil-Red O. The top and middle panels at the right show microscopic pictures of differentiated cells at higher magnification. The bottom panels on the right side show western blots of C/EBP $\beta$  and  $\beta$ -actin expression in undifferentiated (lane 1) and differentiated (lanes 2–4) cells treated without Celastrol (lane 2) or with 0.5  $\mu$ M (lane 3) and 1  $\mu$ M (lane 4) Celastrol.

<https://doi.org/10.1371/journal.pone.0190934.g001>

with the function of its DNA-binding domain was confirmed by comparing the effect of Celastrol on the expression of MRP126 mRNA induced by C/EBP $\beta$  or a Jun-C/EBP $\beta$  hybrid protein containing the DNA-binding domain of C/EBP $\beta$  and the transactivation domain of c-Jun (Fig 2B). While Celastrol strongly inhibited the activity of C/EBP $\beta$  that of Jun-C/EBP $\beta$  was virtually unaffected, clearly indicating that Celastrol does not exert its inhibitory effect via the DNA-binding domain of C/EBP $\beta$ . It is also obvious that the activity of C/EBP $\beta$  strongly exceeded that of Jun-C/EBP $\beta$  (compare first and third lane of Fig 2B). This might be related to the ability of C/EBP $\beta$  to function as a pioneer transcription factor [34].

Our previous work has implicated the co-activator p300 as a key cooperation and interaction partner of C/EBP $\beta$  [33]. Hence, we were interested to know if Celastrol disrupts the cooperation of C/EBP $\beta$  with p300. We first examined if increased expression of p300 overrides the inhibitory effect of Celastrol. We co-transfected fibroblasts with expression vectors for C/EBP $\beta$  and p300 and analyzed the effect of Celastrol on the expression of MRP126 mRNA. As control, the cells were transfected only with C/EBP $\beta$  expression vector. Fig 2C shows that the inhibition of MRP126 expression by Celastrol was less strong when C/EBP $\beta$  was co-expressed with p300 than when it was expressed alone, however, p300 only partially rescued C/EBP $\beta$  activity.

We have previously shown that the interaction of p300 and C/EBP $\beta$  triggers phosphorylation of multiple sites in the C-terminal domain of p300 thereby increasing its co-activator activity [37]. The C/EBP $\beta$ -induced phosphorylation is catalyzed by protein kinase Hipk2, which itself interacts with the amino terminal domain of C/EBP $\beta$  [38]. To confirm that Celastrol disturbs the cooperation of C/EBP $\beta$  and p300 we analyzed its effect on the C/EBP $\beta$ -induced phosphorylation of p300, which can be monitored by the mobility shift of the C-terminal domain of p300 induced by C/EBP $\beta$  [37]. We also used an antiserum against phosphorylated Ser-2280 of p300, one of the sites whose phosphorylation is increased by C/EBP $\beta$  [38]. Fig 2D shows that Celastrol indeed suppressed the C/EBP $\beta$ -induced phosphorylation of p300, as evidenced by the less-pronounced C/EBP $\beta$ -induced mobility shift and a decrease in the phosphorylation of Ser-2280. Because co-immunoprecipitation experiments showed that Celastrol did not interfere with the interaction of C/EBP $\beta$  and Hipk2 we wondered whether Celastrol disrupts the interaction of p300 and C/EBP $\beta$ . To address this we co-transfected fibroblasts with expression vectors for YFP-tagged C/EBP $\beta$  and p300/1751-1947, which harbors the C/EBP $\beta$  binding region of p300 [33]. We then incubated extracts of the transfected cells with GFP-trap beads (which carry a covalently bound high-affinity GFP binding protein) to precipitate YFP-C/EBP $\beta$  together with the bound p300/1751-1947, allowing us to monitor the interaction of both proteins. Fig 3A shows that p300/1751-1947 was co-precipitated with YFP-C/EBP $\beta$  whereas expression of YFP as control did not result in co-precipitation, indicating that the interaction between p300/1751-1947 and YFP-C/EBP $\beta$  observed in this assay is specific. We then performed co-precipitation experiments with extracts from cells that had been treated with Celastrol for different times



**Fig 2. Celastrol does not affect DNA-binding of C/EBP $\beta$  but disrupts its cooperation with p300.** **A.** A radiolabeled double-stranded oligonucleotide with a consensus C/EBP binding site was incubated without nuclear extract, with nuclear extract from cells transfected with C/EBP $\beta$  expression vector and treated with Celastrol as indicated, or with nuclear extract from untransfected cells. Protein-DNA complexes were analyzed by native polyacrylamide gel electrophoresis. The top panel shows a western blot analysis of aliquots of the nuclear extracts stained with antibodies against C/EBP $\beta$ . **B.** QT6 cells transfected with expression vectors for C/EBP $\beta$  or a Jun-C/EBP $\beta$  hybrid protein and incubated with Celastrol were analyzed by northern blotting for expression of *MRP126* and *S17* mRNAs. **C.** QT6 cells transfected with expression vectors for C/EBP $\beta$  and p300 were treated with the indicated concentrations of Celastrol, harvested after 16 hours and analyzed by northern blotting for expression of *MRP126* and *S17* mRNAs. Numbers below the lanes indicate the amounts of *MRP126* mRNA relative to the *S17* mRNA as control determined by quantification with a phosphor image analyzer. The signals for *MRP126* and *S17* mRNAs were obtained by sequential hybridization of the same blot with specific radiolabeled probes. **D.** QT6 cells transfected with expression vectors for C/EBP $\beta$  and p300/1751-2379 and cultivated for 24 hours with or without Celastrol. Aliquots of total cell extracts were then analyzed with antibodies against p300, pp300(Ser2280) and C/EBP $\beta$ . Black and white arrows mark the un-phosphorylated and highly phosphorylated p300.

<https://doi.org/10.1371/journal.pone.0190934.g002>

before harvesting. Fig 3B shows that Celastrol clearly inhibited the co-precipitation of both proteins, demonstrating that it disrupts the C/EBP $\beta$ -p300 interaction.



To substantiate this conclusion we performed a mammalian two-hybrid experiment, using a Gal4-inducible luciferase reporter construct and expression vectors for Gal4-C/EBP $\beta$  and a p300/VP16 construct that contains the Taz2 domain of p300 [33,37]. As expected, p300/VP16 strongly stimulated the activity of the Gal4-dependent reporter gene in the presence but not in the absence of Gal4-C/EBP $\beta$  (Fig 3C). Increasing concentrations of Celastrol gradually reduced the activity of the reporter gene, confirming that Celastrol interferes with the C/EBP $\beta$ -Taz2 interaction. Control transfections showed that Celastrol did not affect the activity of Gal4-VP16. This indicated that Celastrol does not inhibit the Gal4 DNA-binding domain or the VP16 transactivation domain. In summary, our data suggest that Celastrol inhibits the activity of C/EBP $\beta$  by disrupting the interaction of C/EBP $\beta$  with the Taz2 domain of p300.

### Celastrol inhibits the p300-C/EBP $\beta$ interaction by targeting cysteine residues located in the Taz2 domain of p300

Celastrol contains reactive groups that can covalently react with cysteine residues in proteins [39]. To investigate if Celastrol inhibits the C/EBP $\beta$ -p300 interaction by alkylating cysteines we first performed GFP-trap experiments in the presence of the thiol reagent  $\beta$ -mercaptoethanol. Cells were transfected with expression vectors for YFP-C/EBP $\beta$  and p300/1751-1947, treated with or without  $\beta$ -mercaptoethanol and Celastrol before cell extracts were subjected to co-precipitation analysis. Fig 4A shows that  $\beta$ -mercaptoethanol partially relieved the inhibitory effect of Celastrol, suggesting that alkylation of cysteine residues might be involved in the inhibition of the C/EBP $\beta$ -Taz2 interaction by Celastrol. However, this experiment did not reveal whether Celastrol targets cysteine residues of C/EBP $\beta$  or p300 (or both). To address this we first used a cysteine-free mutant of C/EBP $\beta$  which was still active as determined by its ability to stimulate the expression of the endogenous *MRP126* gene. Importantly, the cysteine-free version of C/EBP $\beta$  largely retained its sensitivity towards Celastrol (Fig 4B), indicating that Celastrol does not inhibit C/EBP $\beta$  by alkylating cysteine residues of C/EBP $\beta$ .

The Taz2 domain of p300 also contains cysteine residues, most of which are involved in the coordination of zinc ions to stabilize the three-dimensional structure of the domain [40] (Fig 5A). Because the coordination of zinc ions reduces the nucleophilicity of these cysteine residues they are less likely to react with electrophiles and were not considered as likely targets for Celastrol. However, the Taz2 domain contains two cysteines located immediately adjacent to each other (Cys-1789 and Cys-1790) that are not involved in zinc coordination and, thus, might be potential targets of alkylation by Celastrol. Previous studies have established that C/EBP family members have at least two binding regions (Box A and B, Fig 5E) that mediate direct protein-protein-interactions with the Taz2 domain. According to a recent model based





**Fig 3. Celastrol disrupts the C/EBP $\beta$ -p300 interaction.** A,B. QT6 fibroblasts transfected with the indicated expression vectors were subjected to GFP-trap experiments. Total cell extracts (TCE) and GFP-trap samples (trap) were analyzed by western blotting with antibodies against GFP and p300. In panel B, the cells were also treated with Celastrol. C. QT6 fibroblasts were transfected with the Gal4-dependent reporter gene pG5E4-38luc, the  $\beta$ -galactosidase plasmid pCMV $\beta$  and expression vectors for Gal4-C/EBP $\beta$  and p300-VP19, as indicated. The transfected cells were incubated for 12 hours with or without Celastrol followed by analysis of luciferase activities. The luciferase activity was first normalized to the  $\beta$ -galactosidase activity. The normalized luciferase activity of the Gal4-C/EBP $\beta$  plus p300-VP16 transfected cells or the Gal4-VP16 transfected cells in the absence of Celastrol was then set to 100%. Asterisks indicate statistical significance (\*\*\*)  $p < 0.001$ , Student's t-test).



<https://doi.org/10.1371/journal.pone.0190934.g003>

on the interaction of C/EBP $\epsilon$  with the Taz2 domain [41] Cys-1789 and Cys-1790 of p300 are located close to the binding region on the Taz2 surface (Fig 5F). To investigate if Cys-1789 and Cys-1790 are involved in the disruption of the C/EBP $\beta$ -Taz2 interaction by Celastrol, we mutated them to alanine and examined the ability of the resulting protein to interact with C/EBP $\beta$ . As shown by the GFP-trap experiment illustrated in Fig 5B, the C1789/1790A mutant protein was able to interact with C/EBP $\beta$ , indicating that the amino acid replacement did not result in massive structural changes of the Taz2 domain. Importantly, the ability of Celastrol to disrupt the C/EBP $\beta$ -Taz2 interaction was completely abolished by the mutation. This strongly suggests that one or both of these cysteines are indeed the targets of Celastrol. To investigate if the replacement of a single cysteine residue is sufficient to abolish the inhibitory effect of Celastrol, we also generated mutants in which only Cys-1789 or Cys-1790 were replaced by alanine. Fig 5C and 5D show that the interaction of C/EBP $\beta$  with both mutant proteins was still inhibited by Celastrol. Thus, the presence of either Cys-1789 or Cys-1790 is sufficient for the disruption of the C/EBP $\beta$ -Taz2 interaction by Celastrol.

## Discussion

We have demonstrated that Celastrol, a natural compound previously shown to inhibit the transcriptional activity of Myb, is also a very potent inhibitor of C/EBP $\beta$ . The activity of C/EBP $\beta$  is strongly dependent on p300, which is recruited to C/EBP $\beta$  via direct protein-protein interactions between C/EBP $\beta$  and the Taz2 domain of p300 [33]. Our data show that Celastrol disrupts the C/EBP $\beta$ -Taz2 interaction, most likely by a mechanism that involves the alkylation of cysteine residues by Celastrol. This is supported by the observation that the thiol reagent  $\beta$ -mercaptoethanol reduces the inhibitory effect of Celastrol on the C/EBP $\beta$ -Taz2 interaction. We showed that a mutant of C/EBP $\beta$  devoid of cysteine residues was inhibited by Celastrol as strongly as wild-type C/EBP $\beta$  whereas alanine-substitution of p300 Cys-1789 and Cys-1790, two cysteine residues within the Taz2 domain that are not involved in binding zinc ions, abolished the inhibitory effect of Celastrol on the C/EBP $\beta$ -Taz2 interaction. Overall, these findings strongly suggest that Celastrol inhibits C/EBP $\beta$  activity by alkylating p300 Cys-1789 or Cys-1790 (or both) and thereby prevents the C/EBP $\beta$ -p300 interaction. We have also tried to detect covalent adducts of Celastrol with these cysteines by mass spectrometry. However, because of difficulties in detecting the relevant peptides these experiments were not conclusive. Definitive proof for the suggested mechanism is therefore awaiting more sophisticated mechanistic analyses.

According to a model of the interaction of C/EBP $\epsilon$  with the Taz2 domain [41] binding is mediated by a bi-partite binding site, which is conserved among C/EBP family members and located at amino acids 68–80 and 115–123 of C/EBP $\beta$ . As shown in Fig 5F, the binding region forms two  $\alpha$ -helices that are separated by a turn (or additional sequences in case of C/EBP $\beta$ ) and interact with the Taz2 domain in the vicinity of Cys-1789 and Cys-1790. It is therefore conceivable that the presence of a bulky ligand, such as Celastrol, at one or both of the cysteines compromises the C/EBP $\beta$ -Taz2 interaction. The activity of C/EBP $\alpha$  was only weakly inhibited by Celastrol (Fig 1D), indicating that Celastrol shows a preference for inhibition of




**Fig 4. Inhibition of a cysteine-free mutant of C/EBP $\beta$  by Celastrol.** A. QT6 fibroblasts were transfected with the indicated expression vectors and treated with Celastrol and  $\beta$ -mercaptoethanol. Cell extracts were then subjected to GFP-trap and analyzed as in Fig 3. B. QT6 cells were transfected with expression vectors for wild-type or cysteine-free C/EBP $\beta$  and treated with or without Celastrol. Cells were analyzed by northern blotting for expression of *MRP126* and *S17* mRNAs as in Fig 1 and Fig 2.

<https://doi.org/10.1371/journal.pone.0190934.g004>

C/EBP $\beta$  versus C/EBP $\alpha$ . This is surprising because C/EBP $\alpha$  is also known to recruit p300 as co-activator [16,42]. However, the presumed p300 binding regions of C/EBP $\alpha$  and C/EBP $\beta$  are not identical; in particular, a sequence stretch that separates the bi-partite p300-binding site of C/EBP $\beta$ , is missing in C/EBP $\alpha$  (Fig 5E). This could lead to differences in the binding modes of C/EBP $\beta$  and C/EBP $\alpha$  that might explain the differential effects of Celastrol on the activities of both C/EBP family members. Structural studies of complexes of the Taz2 domain with C/EBP $\beta$  and C/EBP $\alpha$  are currently not available and will be required to address these issues in future studies. In more general terms, the differential effects of Celastrol on C/EBP $\beta$  and C/EBP $\alpha$  clearly illustrate that a compound that acts through reactive groups can nevertheless have quite specific inhibitory activity.

We recently identified Celastrol as a Myb inhibitor using a reporter gene driven by the cis-regulatory elements of the Myb target gene *mim-1* [25]. Mechanistic studies have shown that Celastrol inhibits Myb activity by disrupting the Myb-Kix interaction, thereby depriving Myb of its essential co-activator [25]. The new data presented here provide a more complex view of the inhibitory potential of Celastrol. It appears that the strong inhibitory effect of Celastrol in our screening system was due to the simultaneous suppression of the activities of Myb and C/EBP $\beta$ , both of which cooperate tightly at the cis-regulatory elements of the *mim-1* gene. In both cases affects Celastrol the cooperation with the co-activator p300, which independently interacts with Myb and C/EBP $\beta$  via the Kix and Taz2 domains, as illustrated schematically in Fig 6. Although this might appear surprising at first glance, such a two-tiered inhibitory mechanism can be rationalized because the activation of the *mim-1* gene by Myb is strongly dependent on the cooperation of Myb with C/EBP $\beta$  [30,31]. Furthermore, p300 not only stimulates the activity of Myb and C/EBP $\beta$  individually but also enhances their synergistic behavior [33], suggesting that Myb, C/EBP $\beta$  and p300 constitute a transcriptional module in myeloid cells. In retrospect, it is therefore not too surprising that a compound that was selected for its ability to strongly inhibit the stimulation of the *mim-1* gene by Myb actually inhibits this transcriptional module by two independent mechanisms. We have recently identified the natural compound Withaferin A as another highly active inhibitor of Myb-induced *mim-1* expression. Interestingly, the analysis of the inhibitory mechanism of Withaferin A has shown that it also acts as a dual Myb and C/EBP $\beta$  inhibitor [27].

We have previously shown that Celastrol inhibits the proliferation of leukemic cells from AML patients while not affecting the proliferation of normal hematopoietic progenitors. Furthermore, Celastrol prolonged survival in an in vivo mouse model of an aggressive AML [25]. Recent genome-wide studies have demonstrated that Myb, C/EBP $\beta$  and p300 together with several other hematopoietic transcription factors and the bromo-domain protein BRD4 co-localize at many genomic sites in AML cells to stimulate the activity of so-called “super-enhancers” that are involved in maintaining the proliferation of the leukemic cells [11]. Although we currently have no evidence that Celastrol affects the activity of super enhancers in AML cells it is nevertheless conceivable that the inhibition of the Myb-C/EBP $\beta$ -p300 transcriptional module by Celastrol might diminish the activity of such enhancers, similar to its effect on the *mim-1* enhancer. Thus, our findings suggest the interesting idea that the promising anti-leukemic effects of Celastrol that we have previously described [25] might be due to the abrogation of Myb, C/EBP $\beta$  and p300 function at super-enhancers in AML cells. While this appears to be an interesting possibility, further work is clearly required to validate this




**Fig 5. Disruption of the C/EBP $\beta$ -p300 interaction is dependent on cysteine residues in the Taz2 domain of p300.** A. Sequence comparison of the p300 Taz2 domain from human, mouse and chicken. Cysteine and histidine residues involved in coordination of zinc ions and Cys-1789 and Cys-1790 are highlighted. B-D. QT6 fibroblasts transfected with the indicated expression vectors were subjected to GFP-trap experiments and analyzed as in Fig 3. E. Amino acid sequences of different C/EBP family members implicated in Taz2-binding. Box A and Box B refer to  $\alpha$ -helical regions assumed to directly interact with the Taz2 domain. F. Partial view of the Taz2 domain and its interaction with Box A and Box B sequences of C/EBP $\epsilon$ . The peptide backbones of the Taz2 domain and of CEBP $\epsilon$  are shown in brown and turquoise, respectively. The position of Cys-1790 is shown. The figure was created from PDB entry 3t92.

<https://doi.org/10.1371/journal.pone.0190934.g005>

notion. It should also be kept in mind that Celastrol is an electrophile that can undergo covalent alkylation reactions with cysteine groups in proteins and thereby might affect a panel of “unspecific” targets. Whether or not Celastrol is an interesting lead molecule for further development of therapeutics against leukemia therefore remains to be seen.

Apart from AML, C/EBP $\beta$  shows pro-oncogenic activities in other tumors, such as tumors of the colon, prostate, ovaries, kidneys and the stomach, where C/EBP $\beta$  expression often correlates with increased malignancy and invasive activity of the tumor cells [43–45]. A strong case has been made for glioblastoma, where high C/EBP $\beta$  expression levels correlate with a poor



**Fig 6. Model of the inhibition of a transcriptional Myb-C/EBP $\beta$ -p300 module by Celastrol.**

<https://doi.org/10.1371/journal.pone.0190934.g006>

prognosis for the patient and C/EBP $\beta$  together with STAT3 plays a key role in establishing a mesenchymal gene expression signature that is responsible for the aggressiveness of high-grade glioblastomas [46–48]. In the hematopoietic system C/EBP $\beta$  plays pro-oncogenic roles in multiple myeloma [49] and anaplastic large cell lymphoma (ALCL) [50]. Overall, C/EBP $\beta$  therefore appears to be also an interesting drug target in its own right.

## Acknowledgments

We thank B.Berkenfeld and P.Pieloch for technical assistance and W.Dörner for help with mass spectrometry.

## Author Contributions

**Conceptualization:** Anna Coulibaly, Thomas J. Schmidt, Karl-Heinz Klempnauer.

**Funding acquisition:** Karl-Heinz Klempnauer.

**Investigation:** Anna Coulibaly, Astrid Haas, Simone Steinmann, Anke Jakobs, Karl-Heinz Klempnauer.

**Validation:** Anna Coulibaly, Karl-Heinz Klempnauer.

**Visualization:** Karl-Heinz Klempnauer.

**Writing – original draft:** Karl-Heinz Klempnauer.

**Writing – review & editing:** Anna Coulibaly, Astrid Haas, Simone Steinmann, Thomas J. Schmidt.

## References

1. Ramsay RG, Gonda TJ. Myb function in normal and cancer cells. *Nat Rev Cancer* 2008; 8: 523–34. <https://doi.org/10.1038/nrc2439> PMID: 18574464
2. Lahortiga I, De Keersmaecker K, Van Vlierberghe P, Graux C, Cauwelier B, Lambert F, et al. Duplication of the MYB oncogene in T cell acute lymphoblastic leukemia. *Nat Genet.* 2007; 39: 593–5. <https://doi.org/10.1038/ng2025> PMID: 17435759
3. Clappier E, Cucchini W, Kalota A, Crinquette A, Cayuela JM, Dik WA, et al. The c-MYB locus is involved in chromosomal translocation and genomic duplications in human T-cell acute leukemia (T-ALL), the translocation defining a new T-ALL subtype in very young children. *Blood* 2007; 110: 1251–61. <https://doi.org/10.1182/blood-2006-12-064683> PMID: 17452517
4. Mansour MR, Abraham BJ, Anders L, Berezovskaya A, Gutierrez A, Durbin AD, et al. Oncogene regulation. An oncogenic super-enhancer formed through somatic mutation of a noncoding intergenic element. *Science* 2014; 346: 1373–7. <https://doi.org/10.1126/science.1259037> PMID: 25394790
5. Calabretta B, Sims RB, Valtieri M, Caracciolo D, Szczylik C, Venturelli D, et al. Normal and leukemic hematopoietic cells manifest differential sensitivity to inhibitory effects of Myb antisense oligodeoxynucleotides: an in vitro study relevant to bone marrow purging. *Proc Natl Acad Sci USA* 1991; 88: 2351–5. PMID: 2006173
6. Somervaille TC, Matheny CJ, Spencer GJ, Iwasaki M, Rinn JL, Witten DM, et al. Hierarchical maintenance of MLL myeloid leukemia stem cells employs a transcriptional program shared with embryonic rather than adult stem cells. *Cell Stem Cell* 2009; 4: 129–40. <https://doi.org/10.1016/j.stem.2008.11.015> PMID: 19200802
7. Zuber J, Rappaport AR, Luo W, Wang E, Chen C, Vaseva AV, et al. An integrated approach to dissecting oncogene addiction implicates a Myb-coordinated self-renewal program as essential for leukemia maintenance. *Genes Dev.* 2011; 25: 1628–140. <https://doi.org/10.1101/gad.17269211> PMID: 21828272
8. Lovén J, Hoke HH, Lin CY, Lau A, Orlando D, Vakoc CR, et al. Selective inhibition of tumor oncogenes by disruption of super-enhancers. *Cell* 2013; 153: 320–34. <https://doi.org/10.1016/j.cell.2013.03.036> PMID: 23582323

9. Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, et al. Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* 2013; 153: 307–19. <https://doi.org/10.1016/j.cell.2013.03.035> PMID: 23582322
10. Hnisz D, Abraham BJ, Lee TI, Lau A, Saint-André V, Sigova AA, et al. Super-enhancers in the control of cell identity and disease. *Cell* 2013; 155: 934–47. <https://doi.org/10.1016/j.cell.2013.09.053> PMID: 24119843
11. Roe J-S, Mercan F, Rivera K, Pappin DJ, Vakoc CR. BET bromodomain inhibition suppresses the function of hematopoietic transcription factors in acute myeloid leukemia. *Mol. Cell* 2015; 58: 1028–39. <https://doi.org/10.1016/j.molcel.2015.04.011> PMID: 25982114
12. Persson M, Andrén Y, Mark J, Horlings HM, Persson F, Stenman G. Recurrent fusion of MYB and NFIB transcription factor genes in carcinomas of the breast and head and neck. *Proc Natl Acad Sci USA* 2009; 106: 18740–4. <https://doi.org/10.1073/pnas.0909114106> PMID: 19841262
13. Zhang J, Wu G, Miller CP, Tatevossian RG, Dalton JD, Tang B, et al. Whole-genome sequencing identifies genetic alterations in pediatric low-grade gliomas. *Nat Genet* 2013; 45: 602–12. <https://doi.org/10.1038/ng.2611> PMID: 23583981
14. Nerlov C. The C/EBP family of transcription factors: a paradigm for interaction between gene expression and proliferation control. *Trends Cell. Biol.* 2007; 17: 318–24. <https://doi.org/10.1016/j.tcb.2007.07.004> PMID: 17658261
15. Kowenz-Leutz E, Twamley G, Ansieau S, Leutz A. Novel mechanism of C/EBP beta (NF-M) transcriptional control: activation through derepression. *Genes Dev.* 1994; 8: 2781–91. PMID: 7958933
16. Steinmann S, Coulibaly A, Ohnheiser J, Jakobs A, Klempnauer K-H. Interaction and cooperation of the CCAAT-box enhancer-binding protein  $\beta$  (C/EBP $\beta$ ) with the homeodomain-interacting protein kinase 2 (Hpk2). *J. Biol. Chem.* 2013; 288: 22257–69. <https://doi.org/10.1074/jbc.M113.487769> PMID: 23782693
17. Ceseña TI, Cui TX, Subramanian L, Fulton CT, Iñiguez-Lluhí JA, Kwok RP, et al. Acetylation and deacetylation regulate CCAAT/enhancer binding protein beta at K39 in mediating gene transcription. *Mol. Cell. Endocrinol.* 2008; 289: 94–101. <https://doi.org/10.1016/j.mce.2008.03.009> PMID: 18486321
18. Pless O, Kowenz-Leutz E, Knoblich M, Lausen J, Beyermann M, Walsh MJ, et al. G9a-mediated lysine methylation alters the function of CCAAT/enhancer-binding protein-beta. *J. Biol. Chem.* 2008; 283: 26357–63. <https://doi.org/10.1074/jbc.M802132200> PMID: 18647749
19. Kowenz-Leutz E, Pless O, Dittmar G, Knoblich M, Leutz A. Crosstalk between C/EBPbeta phosphorylation, arginine methylation, and SWI/SNF/Mediator implies an indexing transcription factor code. *EMBO J.* 2010; 29: 1105–15. <https://doi.org/10.1038/emboj.2010.3> PMID: 20111005
20. Eaton EM, Sealy L. Modification of CCAAT/enhancer-binding protein-beta by the small ubiquitin-like modifier (SUMO) family members, SUMO-2 and SUMO-3. *J. Biol. Chem.* 2003; 278: 33416–21. <https://doi.org/10.1074/jbc.M305680200> PMID: 12810706
21. Descombes P, Schibler U. A liver-enriched transcriptional activator protein, LAP, and a transcriptional inhibitory protein, LIP, are translated from the same mRNA. *Cell* 1991; 67: 569–79. PMID: 1934061
22. Calkhoven CF, Müller C, Leutz A. Translational control of C/EBP $\alpha$  and C/EBP $\beta$  isoform expression. *Genes Dev.* 2000; 14: 1920–32. PMID: 10921906
23. Bujnicki T, Wilczek C, Schomburg C, Feldmann F, Schlenke P, Müller-Tidow C, et al. Inhibition of Myb-dependent gene expression by the sesquiterpene lactone mexicanin-I. *Leukemia* 2012; 26: 615–22. <https://doi.org/10.1038/leu.2011.275> PMID: 21986841
24. Uttarkar S, Dukare S, Bopp B, Goblirsch M, Jose J, Klempnauer K-H. Naphthol AS-E phosphate inhibits the activity of the transcription factor Myb by blocking the interaction with the KIX domain of the coactivator p300. *Mol. Cancer Ther.* 2015; 14: 1276–85. <https://doi.org/10.1158/1535-7163.MCT-14-0662> PMID: 25740244
25. Uttarkar S, Dassé E, Coulibaly A, Steinmann S, Jakobs A, Schomburg C, et al. Targeting acute myeloid leukemia with a small molecule inhibitor of the Myb/p300 interaction. *Blood* 2015; 127: 1173–82. <https://doi.org/10.1182/blood-2015-09-668632> PMID: 26631113
26. Uttarkar S, Piontek T, Dukare S, Schomburg C, Schlenke P, Berdel WE, et al. Small-molecule disruption of the Myb/p300 cooperation targets acute myeloid leukemia cells. *Mol. Cancer Ther.* 2016; 15: 2905–15. <https://doi.org/10.1158/1535-7163.MCT-16-0185> PMID: 27707899
27. Falkenberg KD, Jakobs A, Matern JC, Dörner W, Uttarkar S, Trentmann A, et al. Withaferin A, a natural compound with anti-tumor activity, is a potent inhibitor of transcription factor C/EBP $\beta$  *Biochim. Biophys. Acta* 2017; 1864: 1349–58. <https://doi.org/10.1016/j.bbamcr.2017.05.003> PMID: 28476645
28. Jakobs A, Uttarkar S, Schomburg C, Steinmann S, Coulibaly A, Schlenke P, et al. An isoform-specific C/EBP $\beta$  inhibitor targets acute myeloid leukemia cells. *Leukemia* 2016; 30: 1612–5. <https://doi.org/10.1038/leu.2016.16> PMID: 26854027



29. Jakobs A, Steinmann S, Henrich SM, Schmidt TJ, Klempnauer K-H. Helenalin acetate, a natural sesquiterpene lactone with anti-inflammatory and anti-cancer activity, disrupts the cooperation of CCAAT-box/enhancer-binding protein beta (C/EBP $\beta$ ) and co-activator p300. *J. Biol. Chem.* 2016; 291: 26098–108. <https://doi.org/10.1074/jbc.M116.748129> PMID: 27803164
30. Burk O, Mink S, Ringwald M, Klempnauer K-H. Synergistic activation of the chicken *mim-1* gene by *v-myb* and C/EBP transcription factors. *EMBO J.* 1993; 12: 2027–38. PMID: 8491193
31. Ness SA, Kowenz-Leutz E, Casini T, Graf T, Leutz A. Myb and NF-M: combinatorial activators of myeloid genes in heterologous cell types. *Genes Dev.* 1993; 7: 749–59. PMID: 7684005
32. Lane MD, Tang QQ, Jiang MS. Role of the CCAAT enhancer binding proteins (C/EBPs) in adipocyte differentiation. *Biochem. Biophys. Res. Commun.* 1999; 266: 677–83. <https://doi.org/10.1006/bbrc.1999.1885> PMID: 10603305
33. Mink S, Haenig B, Klempnauer K-H. Interaction and functional collaboration of p300 and C/EBP $\beta$ . *Mol. Cell. Biol.* 1997; 17: 6609–17. PMID: 9343424
34. Plachetka A, Chayka O, Wilczek C, Melnik S, Bonifer C, Klempnauer K-H. C/EBP $\beta$  induces chromatin opening at a cell-type-specific enhancer. *Mol. Cell. Biol.* 2008; 28: 2102–12. <https://doi.org/10.1128/MCB.01943-07> PMID: 18195047
35. Cao Z, Umek RM, McKnight SL. Regulated expression of three C/EBP isoforms during adipose conversion of 3T3-L1 cells. *Genes Dev.* 1991; 5: 1538–52. PMID: 1840554
36. Mink S, Kerber U, Klempnauer K-H. Interaction of C/EBPbeta and v-Myb is required for synergistic activation of the *mim-1* gene. *Mol Cell Biol* 1996; 16: 1316–25. PMID: 8657104
37. Schwartz C, Beck K, Mink S, Schmolke M, Budde B, Wenning D, et al. Recruitment of p300 by C/EBP-beta triggers phosphorylation of p300 and modulates coactivator activity. *EMBO J.* 2003; 22: 882–92. <https://doi.org/10.1093/emboj/cdg076> PMID: 12574124
38. Steinmann S, Schulte K, Beck K, Chachra S, Bujnicki T, Klempnauer K-H. v-Myc inhibits C/EBPbeta activity by preventing C/EBPbeta-induced phosphorylation of the co-activator p300. *Oncogene* 2009; 28: 2446–55. <https://doi.org/10.1038/onc.2009.90> PMID: 19448669
39. Salminen A, Lehtonen M, Paimela T, Kaarniranta K. Celastrol: Molecular targets of Thunder God Vine. *Biochem. Biophys. Res. Commun.* 2010; 394: 439–42. <https://doi.org/10.1016/j.bbrc.2010.03.050> PMID: 20226165
40. De Guzman RN, Liu HY, Martinez-Yamout M, Dyson HJ, Wright PE. Solution structure of the TAZ2 (CH3) domain of the transcriptional adaptor protein CBP. *J. Mol. Biol.* 2000; 303: 243–53. <https://doi.org/10.1006/jmbi.2000.4141> PMID: 11023789
41. Bhaumik P, Davis J, Tropea JE, Cherry S, Johnson PF, Miller M. Structural insights into interactions of C/EBP transcriptional activators with the Taz2 domain of p300. *Acta Crystallogr. D Biol. Crystallogr.* 2014; 70: 1914–21.
42. Erickson RL, Hemati N, Ross SE, MacDougald OA. p300 coactivates the adipogenic transcription factor CCAAT/enhancer-binding protein alpha. *J. Biol. Chem.* 2001; 276: 16348–55. PMID: 11340085
43. Rask K, Thörn M, Pontén F, Kraaz W, Sundfeldt K, Hedin L, et al. Increased expression of the transcription factors CCAAT-enhancer binding protein-beta (C/EBPbeta) and C/EBPzeta (CHOP) correlate with invasiveness of human colorectal cancer. *Int. J. Cancer* 2000; 86: 337–43. PMID: 10760820
44. Kim MH, Minton AZ, Agarwal V. C/EBPbeta regulates metastatic gene expression and confers TNF-alpha resistance to prostate cancer cells. *Prostate* 2009; 69: 1435–47. <https://doi.org/10.1002/pros.20993> PMID: 19489038
45. Regalo G, Förster S, Resende C, Bauer B, Fleige B, Kemmner W, et al. C/EBP $\beta$  regulates homeostatic and oncogenic gastric cell proliferation. *J. Mol. Med.* 2016; 94: 1385–95. <https://doi.org/10.1007/s00109-016-1447-7> PMID: 27522676
46. Homma J, Yamanaka R, Yajima N, Tsuchiya N, Genkai N, Sano M, et al. Increased expression of CCAAT/enhancer binding protein beta correlates with prognosis in glioma patients. *Oncol. Rep.* 2006; 15: 595–601. PMID: 16465418
47. Aguilar-Morante D, Morales-Garcia JA, Santos A, Perez-Castillo A. CCAAT/enhancer binding protein beta induces motility and invasion of glioblastoma cells through transcriptional regulation of the calcium binding protein S100A4. *Oncotarget* 2015; 6: 4369–84. <https://doi.org/10.18632/oncotarget.2976> PMID: 25738360
48. Carro MS, Lim WK, Alvarez MJ, Bollo RJ, Zhao X, Snyder EY, et al. The transcriptional network for mesenchymal transformation of brain tumours. *Nature* 2010; 463: 318–27. <https://doi.org/10.1038/nature08712> PMID: 20032975
49. Pal R, Janz M, Galson DL, Gries M, Li S, Jöhrens K, et al. C/EBPbeta regulates transcription factors critical for proliferation and survival of multiple myeloma cells. *Blood* 2009; 114: 3890–8. <https://doi.org/10.1182/blood-2009-01-201111> PMID: 19717648

50. Piva R, Pellegrino E, Mattioli M, Agnelli L, Lombardi L, Boccalatte F, et al. Functional validation of the anaplastic lymphoma kinase signature identifies CEBPB and BCL2A1 as critical target genes. *J. Clin. Invest.* 2006; 116: 3171–82. <https://doi.org/10.1172/JCI29401> PMID: 17111047