

RESEARCH ARTICLE

Chemical Composition and Biological Activities of Trans-Himalayan Alga *Spirogyra porticalis* (Muell.) Cleve

Jatinder Kumar¹, Priyanka Dhar¹, Amol B. Tayade¹, Damodar Gupta², Om P. Chaurasia^{1*}, Dalip K. Upreti⁴, Kiran Toppo⁴, Rajesh Arora^{2,3}, M. R. Suseela⁴, Ravi B. Srivastava¹

1 Defence Institute of High Altitude Research, Defence Research & Development Organisation, Leh-Ladakh, Jammu & Kashmir, India, **2** Medicinal and Aromatic Plants Laboratory, Radiation Biotechnology Group, Institute of Nuclear Medicine and Allied Sciences, Defence Research and Development Organisation, Brig S. K. Mazumdar Marg, Delhi, India, **3** Office of the Director General-Life Sciences, DRDO Bhawan, Rajaji Marg, New Delhi, India, **4** National Botanical Research Institute, Rana Pratap Marg, Lucknow, Uttar Pradesh, India

* dropchaurasiadihardrdo@gmail.com



OPEN ACCESS

Citation: Kumar J, Dhar P, Tayade AB, Gupta D, Chaurasia OP, Upreti DK, et al. (2015) Chemical Composition and Biological Activities of Trans-Himalayan Alga *Spirogyra porticalis* (Muell.) Cleve. PLoS ONE 10(2): e0118255. doi:10.1371/journal.pone.0118255

Academic Editor: Gianfranco Pintus, University of Sassari, ITALY

Received: July 13, 2014

Accepted: December 25, 2014

Published: February 18, 2015

Copyright: © 2015 Kumar et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data are available from the central records keeping centre of Defence Institute of High Altitude Research (DIHAR) for researchers who meet the criteria for access to confidential data. Please direct requests for data to Mr. Jatinder Kumar (Defence Institute of High Altitude Research, DRDO, Leh-Ladakh, J&K, Pin 194101).

Funding: The study was entirely supported by Defence Research & Development Organisation (DRDO), Ministry of Defence, Government of India. The funder had no role in study design, data

Abstract

The freshwater alga *Spirogyra porticalis* (Muell.) Cleve, a filamentous charophyte, collected from the Indian trans-Himalayan cold desert, was identified on the basis of morpho-anatomical characters. Extracts of this alga were made using solvents of varying polarity *viz.* n-hexane, acetonitrile, methanol and water. The antioxidant capacities and phenolic profile of the extracts were estimated. The methanol extract showing highest antioxidant capacity and rich phenolic attributes was further investigated and phytochemical profiling was conducted by gas chromatography-mass spectrometry (GC/MS) hyphenated technique. The cytotoxic activity of methanol extract was evaluated on human hepatocellular carcinoma HepG2 and colon carcinoma RKO cell lines. The anti-hypoxic effect of methanol extract of the alga was tested on *in vivo* animal system to confirm its potential to ameliorate oxidative stress. The antioxidant assays *viz.* ferric reducing antioxidant power (FRAP), 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS), 1,1-diphenyl-2-picrylhydrazyl (DPPH) and nitric oxide (NO) radical scavenging capacities, β -carotene-linoleic acid bleaching property and lipid peroxidation exhibited analogous results, wherein the algal extracts showed significantly high antioxidant potential. The extracts were also found to possess high content of total proanthocyanidin, flavonoid and polyphenol. GC/MS analysis revealed the presence of thirteen chemotypes in the methanol extract representing different phytochemical groups like fatty acid esters, sterols, unsaturated alcohols, alkynes etc. with substantial phyto-pharmaceutical importance. The methanol extract was observed to possess anticancer activity as revealed from studies on HepG2 and RKO cell lines. In the present study, *S. porticalis* methanol extract also provided protection from hypoxia-induced oxidative stress and accelerated the onset of adaptative changes in rats during exposure to hypobaric hypoxia. The bioactive phytochemicals present in this trans-Himalayan alga are of enormous interest and can be utilized sustainably for discovery of novel drugs against oxidative stress.

collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

Natural antioxidants present in plants are known to play a vital role against stress-induced oxidative damage resulting due to the production of reactive oxygen species (ROS) within the cellular milieu, or due to various diseases and ageing processes [1–4]. Consequently, such compounds reflect potential in promoting overall health and well-being, and hold potential for medicinal use, particularly for treating a broad range of ailments and maladies confronting humans. As a consequence, various natural plant forms like micro- and macro-algae, lichens, macromycetes and plants etc. have become the major bioresources for natural antioxidants which find substantial use for preventing and treatment of a number of diseases. Algae are of our prime interest in the search for natural antioxidants as they have been reported to possess significant antioxidant activity and free radical scavenging properties [5,6].

Algae are a genetically diverse group of plants that possess a broad range of physiological and biochemical characteristics. Algae are promising source of novel biochemically active compounds like fatty acids, steroids, carotenoids, polysaccharides, lectins, vitamins and phyco-proteins, amino acids, dietary minerals, halogenated compounds, polyketides, toxins and diverse antioxidants. Numerous extraction and analytical techniques were employed to study the chemical composition of algae from different natural sources like marine ecosystem, freshwater habitat and other environmental conditions. A large number of algal products are also find reliability in the food, cosmetic, biochemical and pharmaceutical industries [7–13].

The Indian trans-Himalayan cold desert region of Ladakh represent a valuable source of large number of natural bio-resources. The diverse floral and faunal composition of this remote Himalayan land provides a vast number of natural products beneficial for the armed forces as well as civil population [14]. This high altitude region is one of the most difficult and hostile terrain for human survival due to the extreme environmental factors and scarcity of fresh foods [15]. It is believed that plants growing in high altitude environment produce natural bio-active substances or secondary metabolites, which help them survive under stressful conditions. The high-altitude plant species of Ladakh region have been reported to possess rich medicinal properties and play major role in the traditional systems of medicine since ancient time [16,17] and could be used as prophylactic and therapeutic agent for high altitude maladies and other stress induced disorders [18–20].

The unicellular microalgae in particular have often been considered as a promising, easy and alternative source of useful natural antioxidants [21–27]. The filamentous freshwater green alga *Spirogyra* spp. (family Zygnemataceae) is consumed as food in many Asian countries. They are valuable source of natural bioactive compounds that are widely exploited for antibiotic, antiviral, antioxidant, anti-inflammatory, cytotoxic and other positive biological activities. In recent time, evaluation of chemical composition and biological activity in natural bio-resources is an urgent task for researchers involved in natural product development and medicinal chemistry. A number of chemometric methods are available to determine the bioactive phytochemotypes and determination of biological activities by *in vitro* and *in vivo* assays which equally produce significant results to assess the pharmaceutical, therapeutic and health promoting properties of natural phytoproducts. These methods have provided an efficient and powerful tool for the quality control, safety and bio-efficacy evaluation of natural products for human consumption [28–37].

Information obtained from different reports revealed that the trans-Himalayan region of Kashmir and Ladakh are a rich source of diverse algal communities [38,39]. We have previously developed a number of herbal products from the high-altitude plants of this region and conducted various studies on medicinal, nutritional and antioxidant potential of these plants and natural products by different biochemical approaches and these products showed prophylactic effects against high-altitude ailments [40–42]. However, there is scarcity of information about

the medicinal and therapeutic potential of algae from this mega diversity hotspot, which prompted us to investigate the antioxidant capacities, phenolic content, GC/MS chemometric profile, cytotoxic effect and anti-hypoxic potential of trans-Himalayan alga *Spirogyra porticalis* (Muell.) Cleve from this cold desert region. The present investigation was undertaken with the objective to estimate the *in vitro* antioxidant capacities and phenolic profile of different solvent extracts of *S. porticalis* from Indian trans-Himalaya. The extract showing highest antioxidant capacity and phenolic content was further investigated to determine the chemometric profile using GC/MS hyphenation and the cytotoxic activity was evaluated on human hepatocellular carcinoma HepG2 and colon carcinoma RKO cells. Finally, the anti-hypoxic effect of *S. porticalis* extract was tested on *in vivo* animal system.

Materials and Methods

Chemicals and reagents

For antioxidant assays, 1,1-diphenyl-2-picrylhydrazyl (DPPH), 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS), 2,4,6-tripyridyl-*s*-triazine (TPTZ), ferrous sulfate ($\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$), aluminium chloride (AlCl_3), sodium acetate ($\text{C}_2\text{H}_3\text{NaO}_2$), sodium carbonate (Na_2CO_3), potassium persulfate ($\text{K}_2\text{S}_2\text{O}_8$), potassium chloride (KCl), ferric chloride (FeCl_3), sodium nitroprusside $\{\text{Na}_2[\text{Fe}(\text{CN})_5\text{NO}] \cdot 2\text{H}_2\text{O}\}$, sulfanilic acid ($\text{C}_6\text{H}_7\text{NO}_3\text{S}$), butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), ascorbic acid, thiobarbituric acid (TBA), quercetin and catechin were purchased from Sigma-Aldrich (St. Louis, MO, USA). Folin Ciocalteu's phenol reagent, vanillin, hydrochloric acid, sulphuric acid, methanol, n-hexane, acetonitrile, chloroform, ethanol and sodium carbonate from Merck Chemical Supplies (Merck KGaA, Darmstadt, Germany) have been utilized. All the other chemicals used, including solvents, were of analytical grade. Absorbance of the test samples and standards were recorded using spectrophotometer (Spectramax M2^e, Molecular Devices, Germany). For GC/MS chemical characterization, CHROMASOLV HPLC grade methanol was used and all other chemicals were of analytical grade and purchased from Sigma-Aldrich. For cytotoxicity assays, RPMI medium 1640, penicillin, streptomycin, trypsin-EDTA, sulphorhodamine B, trichloroacetic acid, acetic acid, Tris base and all other analytical grade chemicals were the products from Sigma-Aldrich. For animal studies, the following chemicals *viz.* heparin, 2',7'-dichlorofluorescein diacetate (DCFHDA), PBS buffer, meta-phosphoric acid, L-glutathione reduced (GSH) and L-glutathione oxidized (GSSG) were also procured from Sigma-Aldrich.

Ethics statement

The necessary permits for the field studies and collection of algae sample were obtained and it was issued by Dr. B. Balaji (IFS), Divisional Forest Officer, Leh Forest Division, Jammu & Kashmir, India. Experimental animals, along with the necessary facilities requisite for the studies were utilized at our sister laboratory Institute of Nuclear Medicine and Allied Sciences (INMAS), DRDO, New Delhi, India. Animal experimental protocols were approved by the INMAS Institutional Animal Ethical Committee (IAEC/2010, extended up to 31st December, 2013) and conformed to national guidelines Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) of Government of India on the care and use of laboratory animals. Proper concern was taken to minimize the sufferings of the studied animals during the experiment.

Algal sample collection

In the fragile microclimate of trans-Himalayan Ladakh region, the average minimum and maximum temperature of Defence Institute of High Altitude Research (DIHAR) campus at 3500 m altitude during 1st March to 30th April, 2011 were -2°C and 11.1°C , which allowed the aggravated growth of *S. porticalis*. The *S. porticalis* sample was collected on 20th April, 2011 from a constructed cemented rectangular pond (size $20 \times 15 \times 2$ m, water pH 7.9) of DIHAR, Leh [altitude 3500 m above sea level (ASL), latitude $34^{\circ}8'16.121''\text{N}$, longitude $77^{\circ}34'19.2219''\text{E}$]. The contamination of other materials like dry waste of some herbs, soil content and earthworms entangled in the algae thallus was carefully avoided from the natural spring habitat (where soil was of sandy desert nature having 56–64% sand content and insignificant amount of organic material) with the help of a small, natural, narrow canal (1 ft depth, 1 ft width and 5–10 cm water level) carrying spring water to the constructed cemented pond. For regular stirring and replacement of old water (except slow moving spring water as inoculum), a fast moving tubewell water was routinely supplied to this pond through the same inlet (cemented canal) and at the same time water is removed by an outlet (outlet cemented canal) daily from 11 a.m. to 4 p.m. The inlet and outlet were entangled with 10 mm and 5 mm sieves respectively to avoid transport of *S. porticalis*. The water depth of the pond was 1–2 ft. Finally, 10 kg wet *S. porticalis* was collected, washed under tap water and air dried in room temperature. The algal material was then preserved in 4% formalin and deposited in the Algal Herbarium, CSIR-National Botanical Research Institute (CSIR-NBRI), Lucknow, India.

Identification of alga

The algal samples were observed under Leica DM500 research microscope (Leica Microsystems, Wetzlar, Germany) and photographs were taken with attached EC3 digital camera. Relevant publication of Prescott (1951) [43] was referred for the identification of algal taxa. For the taxonomic description of taxa, dimensions were given in micrometer (μm). All measuring scales given for algae photographs were equal to 10 μm . The specimens were preserved in the Algal Herbarium, CSIR-NBRI, Lucknow.

Algae materials and extraction

S. porticalis was air dried at room temperature to constant weight. The dried algal material was ground separately to powder. Finely ground dried sample (50 g) was taken for sequential extraction in four solvent systems with increasing polarity *viz.* n-hexane, acetonitrile, methanol and water by Soxhlet apparatus (Borosil GlassWorks Limited, Worli, Mumbai, India) under the following temperatures such as 20°C for hexane, 30°C for acetonitrile, 40°C for methanol and 45°C for water. Then all extracts were concentrated under reduced pressure at 40°C by cold water circulation using thermostat maintained at 4°C for minimizing the degradation of thermo labile compounds. The water extract was concentrated by sublimation in Freeze Dryer (Christ Alpha 1–4 LSC Freeze Dryer, Martin Christ Freeze Dryers GmbH, Osterode am Harz). All extracts were finally stored at -80°C freezer for further analysis.

Antioxidant capacities and free radical scavenging assays

Ferric reducing antioxidant power (FRAP). The FRAP assay was performed using the method depicted by Ikram and co-workers (2009) with slight alteration [44]. A total of 75 μl extract and 225 μl distilled water were added to 2.25 ml of freshly prepared FRAP reagent [10 parts of 300 mM sodium acetate buffer at pH 3.6, 1 part of 10 mM 2,4,6-tri (2-pyridyl)-s-triazine (TPTZ) solution and 1 part of 20 mM $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$]. The content was incubated in the dark for

30 min. The increase in absorbance with the formation of colored product (ferrous tripyridyl-triazine complex) was measured at 593 nm. The antioxidant capacity of the algal extract was determined based on a calibration curve plotted using $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ at a concentration ranging between 0.125 and 2 mM. The calibration equation for $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ was $y = 0.464x - 0.0114$, $R^2 = 0.9996$, where x was absorbance at 593 nm and y was concentration of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (mM). Results were expressed in mM Fe (II)/g of extract.

ABTS radical scavenging activity. ABTS assay was performed according to the preceding protocol [45]. The stock solution was prepared by mixing equal volume of 7 mM ABTS solution and 2.45 mM potassium persulfate solution followed by incubation for 12 h at room temperature in the dark to yield a dark-colored solution containing $\text{ABTS}^{\cdot+}$ radicals. Working solution was prepared freshly before each assay by diluting the stock solution by mixing of stock solution to 50% methanol for an initial absorbance of 0.700 ± 0.02 at 745 nm, with temperature control set at 30°C. Free radical scavenging activity was assessed by mixing 30 μl of different fractions (different concentration in respective solvents) with 300 μl of ABTS working standard. The decrease in absorbance was measured after 6 min at 30°C. Quercetin was used as positive control. The scavenging activity was estimated based on the percentage of ABTS radicals scavenged by the following formula:

$\% \text{ scavenging} = [(A_0 - A_s)/A_0] \times 100$, where A_0 was absorbance of control, A_s was absorbance of extract. The antioxidant value was expressed as the half maximal inhibitory concentration (IC_{50}), the amount of antioxidant required to decrease the initial ABTS concentration by 50%. The IC_{50} value was calculated from the regression equation between sample concentration and rate of inhibition.

Inhibition effect on 1,1-diphenyl-2-picrylhydrazyl radical (DPPH)

The method suggested by previous investigators was adopted for this purpose with minor modification [46,47]. The 0.1 mM solution of DPPH in methanol was prepared and 500 μl of this solution was reacted with 25 μl of algal extract. The control sample was reacted with 25 μl of solvent instead of the extract. The mixture was incubated at room temperature for 30 min before the shrink in absorbance at 517 nm was recorded. Ascorbic acid was used as positive control. Antioxidant value was expressed as the half maximal inhibitory concentration (IC_{50}), the amount of antioxidant required to decrease the initial DPPH concentration by 50%. The IC_{50} values were calculated as described previously.

β -carotene-linoleic acid bleaching assay. The method described by Dapkevicius et al. (1998) [48] was used for β -carotene-linoleic acid bleaching assay. A stock solution of β -carotene and linoleic acid was prepared as follows: 0.5 mg of β -carotene was dissolved in 1 ml of chloroform, and 25 μl of linoleic acid with 200 mg of Tween 20 were added. After chloroform evaporation under vacuum, 100 ml of distilled water was added to the residue. An aliquot of 500 μl of extract (in different concentration) was pipetted into separated test tubes and 5 ml of the previous mixture was added. The test tubes were incubated for 2 h at 50°C together with the control sample. The absorbance was measured at 470 nm at the beginning ($t = 0$ min) and after the experiment ($t = 120$ min). BHT was used as positive control. The antioxidant capacity was calculated as percentage inhibition of oxidation (% AA) using the following equation:

$\% \text{ AA} = [1 - (\text{Abs}_s^0 - \text{Abs}_s^{120}) / (\text{Abs}_c^0 - \text{Abs}_c^{120})] \times 100$, where Abs_s^0 was absorbance of sample at t_0 , Abs_s^{120} was absorbance of sample at t_{120} , Abs_c^0 was absorbance of control at t_0 , Abs_c^{120} was absorbance of control at t_{120} . The IC_{50} values were then calculated by aforementioned method.

Nitric oxide (NO) scavenging activity. The previously published method was used to assay the scavenging activity of extracts on nitric oxide (NO) with minor modification [49]. The

reaction solution (100 μ l) containing 10 mM sodium nitroprusside in phosphate buffer saline (PBS, pH 7.0) was reacted with 10 μ l extract (of different concentration). The reaction mixture was then incubated at 37°C for 1 h. Thereafter 50 μ l aliquot was mixed with 50 μ l of Griess reagent and the absorbance at 540 nm was measured. Percent inhibition of NO produced was calculated by comparing with the absorbance value of the negative control (10 mM sodium nitroprusside and PBS). BHA was used as the positive control. Finally, the IC₅₀ value was calculated.

Lipid peroxidation assay. A modified thiobarbituric acid reactive species (TBARS) assay [50] was employed to measure the lipid peroxide formed using egg yolk homogenate as lipid rich media [51]. Egg homogenate (0.5 ml of 10%, v/v) and 0.1 ml of extract were added to a test tube and made up to 1 ml with distilled water. After that, 0.05 ml of FeSO₄ (0.07 M) was added to induce lipid peroxidation and the mixture was incubated for 30 min. Then, 1.5 ml of 20% acetic acid (pH 3.5) and 1.5 ml of 0.8% (w/v) thiobarbituric acid in 1.1% sodium dodecyl sulphate were added and the resulting mixture was vortexed and then heated at 95°C for 60 min. After cooling, 5 ml of butan-1-ol was added to each tube and centrifuged at 3,000 rpm for 10 min. The absorbance of the organic upper layer was measured at 532 nm. Inhibition of lipid peroxidation (%) by the extract was calculated as $[(1-E)/C] \times 100$, where C was absorbance of the fully oxidized control and E was absorbance in presence of extract. The IC₅₀ value was calculated as depicted previously.

Total proanthocyanidin, flavonoid and polyphenolic content

Total polyphenol content (TPC). TPC was determined by Folin Ciocalteu's colorimetric method as described by previous investigators [52]. Extract solution (10 μ l) was mixed with 20 μ l of 10% Folin Ciocalteu's reagent and 200 μ l of H₂O, following incubation at room temperature for 3 min. After that 100 μ l of sodium carbonate (20%, w/v) was added to the reaction mixture. The content was incubated at room temperature in the dark for 30 min and absorbance was measured at 765 nm. Estimation of total polyphenol was carried out on the basis of calibration curve: $y = 0.0104x - 0.0589$, $R^2 = 0.9978$, where x was absorbance and y was gallic acid equivalent (GAE) at a final concentration of 100 μ g/ml. The results were expressed in GAE (mg/g of extract).

Total flavonoid content (TFC). TFC was analyzed with the method of Ordonez et al. (2006) [53]. Briefly, 100 μ l of sample and 2% AlCl₃ in ethanol solution were allowed to react and the reaction mixture was incubated for 1 h at room temperature and the absorbance was finally measured at 420 nm. Total flavonoid content was calculated as quercetin equivalent (QE) in mg/g of extract using the following equation based on the calibration curve: $y = 0.0235x - 0.1803$, $R^2 = 0.9994$, where x was absorbance and y was QE (mg/g of extract) at final concentration 100 μ g/ml.

Total proanthocyanidins content (TPAC). The TPAC content was determined according to the method of Sun et al. (1998) [54] with slight modification. A volume of 0.5 ml of extract solution was allowed to react with 3 ml of 4% vanillin-methanol solution and 1.5 ml hydrochloric acid. Then the mixture was allowed to stand for 15 min. The absorbance was measured at 500 nm at 30°C. Total proanthocyanidin content was calculated as catechin equivalent (CE) in mg/g of extract using the following equation based on the calibration curve: $y = 0.0015x - 0.0209$, $R^2 = 0.9975$, where x was absorbance and y was CE (mg/g of extract). Extract samples were evaluated at a final concentration of 100 μ g/ml.

GC/MS analysis

Preparation of sample for GC/MS analysis. The concentrated methanol extract (25 mg) was dissolved in 25 ml of the solvent, vortexed properly and filtered through 0.22 μm syringe filter (Millipore Corp., Bedford, MA, USA). For requisite analysis, one microlitre aliquot of the sample solution was then injected into the GC/MS MS system.

Instrumentation and chromatographic conditions. Thermo Finnigan PolarisQ Ion Trap GC/MS MS system comprising of an AS2000 liquid autosampler (Thermo Finnigan, Thermo Electron Corporation, Austin, TX, USA) was used for GC/MS analysis. As per our previous work, the peaks in the chromatogram were identified on the basis of their mass spectra [40,55]. The gas chromatograph was interfaced to a mass spectrometer instrument utilizing the following conditions *viz.* Durabond DB-5 ms column (30 m \times 0.25 mm \times 0.25 μm), operating in electron impact [electron ionisation positive (EI+)] mode at 70 eV, helium (99.999%) as carrier gas at a constant flow of 1 ml/min and an injection volume of 0.5 EI (split ratio of 10:1), injector temperature 280°C and transfer line temperature 300°C. The oven temperature of the system was programmed from 50°C (isothermal for 2 min), with gradual increase in steps of 10°C/min, to 300°C. Mass spectra were taken at 70 eV, a scan interval of 0.5 s and full mass scan range from 25 m/z to 1000 m/z. Finnigan Xcalibur data acquisition and processing software version 2.0 (ThermoQuest, LC and LC/MS Division, San Jose, California, USA) was used for data acquisition.

Identification of components. NIST/EPA/NIH Mass Spectral Database (NIST11), with NIST MS search program v.2.0 g [National Institute Standard and Technology (NIST), Scientific Instrument services, Inc., NJ, USA] was utilized for the interpretation GC/MS mass spectrum. Finally, name, molecular weight and structure of the components of the test material were determined.

Cytotoxic effects of *S. porticalis* methanol extract on carcinoma cells

Cell cultures. Two cancerous cell lines namely human hepatocellular cancer (HepG2) and colon cancer (RKO) cell lines were procured from American Type Cell Culture (ATCC), United States of America (USA). The RPMI medium 1640 (supplemented with 10% heat-inactivated foetal bovine serum, 100 units/ml of penicillin and 100 $\mu\text{g}/\text{ml}$ of streptomycin, pH 7.4) was used to maintain the cells at 37°C in 5% CO₂ and 95–100% humidified air atmosphere.

Morphological determination of cell toxicity. Cells growing logarithmically were seeded in 96 flat-well plate (HepG2 cells 3,000/well and RKO cells 3,000/well) and incubated for 24 h to have a partial monolayer. Cells were treated with 29.26, 43.90, 65.84, 98.77, 148.15, 222.22 and 333.33 $\mu\text{g}/\text{ml}$ of *S. porticalis* methanol extract dissolved in RPMI 1640 medium and subsequently incubated at 37°C in a 5% CO₂ and 95–100% humidified air atmosphere (Touch 190S, LEEC, Nottingham, UK). Microscopic assessment was performed under 10X eyepiece and 10X objective lenses with inverted phase contrast microscope (Dewinter Optical, Inc., Delhi, India) and the changes in cellular morphology following different treatments were recorded at 24, 48 and 72 h of time intervals.

Sulphorhodamine B (SRB) assay for cytotoxicity screening. The Sulphorhodamine B (SRB assay) was used for determining cellular growth based on the SRB uptake by cellular protein. The SRB assay for colorimetric determination of cytotoxicity of *S. porticalis* methanol extract was performed [56,57]. The cells were harvested with 0.5% trypsin-EDTA. Then just before T₀ time (T₀ = zero time after extract inoculation in the cell culture) and just after T_t time (T_t = after 72 h of extract inoculation in cell culture) of plate incubation at 37°C in 5% CO₂, in humidified incubator (95–100% humidity), the SRB assay was performed. The medium was removed at both time points and the cells were fixed with trichloroacetic acid (20%, w/v) at 4°C

for 1 h, stained for 30 min with SRB (0.4%, w/v) dissolved in 1% acetic acid for 30 min, and washed four times with 1% acetic acid. The protein-bound dye was made solubilised with 10 mmol/L tris base, pH 10.5. The absorbance of each well was read at 565 nm using spectrophotometer (Spectramax M2^e, Molecular Devices, Germany).

Percentage growth (PG) at each concentration level was calculated as:

% growth = $[(T_t - T_0)/(C - T_0)] \times 100$ for concentrations in which $T_t \geq T_0$ and

% growth = $[(T_t - T_0)/T_0] \times 100$ for concentrations in which $T_t < T_0$

The 50% growth inhibition concentration (GI₅₀, µg/ml) was determined as the concentration where $\{(T_t - T_0)/(C - T_0)\} \times 100 = 50$. Total growth inhibition concentration (TGI, µg/ml) was measured as the concentration where $\{(T_t - T_0)/(C - T_0)\} \times 100 = 0 = PG$. Lethal concentration that kills 50% of the cells (LC₅₀, µg/ml) was calculated as the concentration level where $\{(T_t - T_0)/T_0\} \times 100 = -50$. These parameters were determined in accordance with previous reports [58–60]. Here, T_0 = growth in the presence of *S. porticalis* extract at time zero, C = control growth and T_t = growth in the presence of *S. porticalis* extract after 72 h.

Effect of *S. porticalis* methanol extract on hypobaric hypoxia induced oxidative stress

Administration of algal extract. Vacuum dried methanol extract was dissolved in distilled water to obtain the desired dose of 200 mg/kg body weight of the animal and administered to rats through oral route in a maximum volume of 1 ml through gastric cannula.

Animals. The animal study experimental protocols were approved by the Institutional Ethics Committee and proper concern was taken to minimize the sufferings of the studied animals during the experiment. Adult male Sprague-Dawley rats weighing 220 ± 10 g were used for the study. The animals were housed in hygienic conditions with day and night cycle of 12 h each with temperature maintained at $25 \pm 2^\circ\text{C}$ and humidity at $65 \pm 5\%$.

Altitude simulation. A specially designed animal decompression chamber was used to expose the animals to simulated altitude of 7,600 m (25,000 ft, 282 mm Hg) for hypobaric hypoxia environment. Here altitude was maintained by reducing the barometric pressure and temperature and humidity were controlled precisely. Continuous hypoxic exposure was provided for the stipulated period with a 10–15 min interval each day for drug administration, food and water replenishment and for cage changing purpose. Continuous flushing of fresh air at a rate of 8 l/min was maintained for preventing the accumulation of carbon dioxide within the chamber. The temperature and humidity in the hypobaric chamber were kept at $32 \pm 2^\circ\text{C}$ and $63 \pm 3\%$ respectively. The ascending and descending rate to hypobaric condition was maintained at 300 m/min. The normoxic animals were kept under similar temperature and humidity conditions in a separate chamber maintained at ambient barometric pressure. A 12 h day and night cycle was maintained in both chambers during the period of exposure.

Experimental design. Male adult Sprague-Dawley rats ($n = 5/\text{group}$) were randomly divided into three groups: *Group I*: served as control group (normoxia group with 0 day exposure), *Group II*: served as hypoxia + vehicle group in which rats were exposed to a simulated altitude of 7,600 m and orally administered with 1 ml of vehicle (distilled water) once in a day for 7 days, *Group III*: served as hypoxia + drug treated group in which rats were exposed to 7,600 m simulated altitude and orally administered with 1 ml of algal extract once in a day for 7 days.

Rats were housed one animal per cage for all study groups for 2 days habituation before the initiation of experimental procedures. The hypoxia groups were allowed to habituate and acclimatize to the experimental conditions for 2 days before the initiation of experimental procedures. Body weight, food consumption and water intake of rats were measured every day and recorded. Food and water were freely available to the animals during the experiment. One

millilitre of distilled water was used as vehicle. The ARRIVE guidelines were followed for reporting the animal research investigation [61]. On completion of the stipulated period of hypoxic exposure, fasting experiments were performed for 24 h and animals were sacrificed at 9.30 a.m. Rats were euthanized by CO₂ asphyxiation followed by cervical dislocation [62]. Then blood samples were obtained from orbital sinus in heparin coated blood collection vial [63].

ROS generation. Following the stipulated period of exposure, serum sample was taken from animals and sample homogenate was prepared in 0.15 M KCl. The free radicals were estimated spectrofluorimetrically with 25 μ l of sample homogenate using DCFHDA as described by LeBel et al. (1990) [64] with slight changes. In brief, 1.494 ml of 0.1 M PBS (pH 7.4) was added to 25 μ l of the sample followed by addition of 6 μ l of 1.25 mM DCFHDA. The sample was then incubated for 15 min at 37°C in dark and the fluorescence was measured at 488 nm excitation and 525 nm emission. The readings thus obtained were converted to fluorescent units/mg protein by calculating the protein present in 25 μ l of each sample from a standard curve.

GSH and GSSG. Reduced glutathione (GSH) and oxidized glutathione (GSSG) levels were measured fluorimetrically from the sample homogenate according to the procedure followed by Hissin and Hilf (1976) [65] with minor modifications. In brief, to 250 μ l of the sample homogenate was added to an equal volume of 10% meta-phosphoric acid, followed by centrifugation at 10,000 g for 30 min at 48°C. The supernatant thus obtained was used for estimation of GSH and GSSG spectrofluorimetrically at 350 nm excitation and 420 nm emission.

Statistical analysis

All experiments were performed in triplicate. The mean values of three individual observations were determined and the standard deviation (SD) was calculated followed by one way analysis of variance (ANOVA) to determine the level of significance among means. The values are represented as mean \pm SD. Statistical difference between groups was calculated using one-way analysis of variance followed by the Neumann-Keuls test for post hoc analysis. The *p* value < 0.05 was considered to be statistically significant.

Results

Morphological identification of alga

The morphology of algal specimen was studied carefully and the specimen was identified as *Spirogyra porticalis* (Muell.) Cleve 1868. The taxonomic description and identification characters were as follows: filament of rather short cells, 40–50 μ m in diameter, 68–200 μ m long, with plane end walls; chloroplast solitary, making 3 to 4 turns; conjugation scalariform by tubes from both gametangia; fertile cells sometimes inflated; zygospores ovate to subglobose-ovate; median spore wall smooth and yellow, 38–50 μ m in diameter, 50–83 μ m long (Fig. 1); common in shallow water of lakes, swamps and roadside ditches [43]. *S. porticalis* showed its appearance in the second week of March, 2011 and found till last week of April to first week of May, 2011 in the spring water and cemented pond in DIHAR.

Antioxidant capacities and phenolic content

In the FRAP assay, methanol extract showed highest (171.49 mM Fe (II)/g extract) and hexane extract exhibited lowest (18.91 mM Fe (II)/g extract) antioxidant capacities. In ABTS assay, methanol extract showed highest antioxidant capacity (IC₅₀ 3.48 mg/ml) and water extract showed lowest antioxidant capacity (IC₅₀ 37.71 mg/ml). In β -carotene-linoleic acid assay, methanol extract again showed highest antioxidant capacity (IC₅₀ 0.20 mg/ml) and acetonitrile




Fig 1. Morphology of *S. porticalis* under Leica DM500 research microscope.

doi:10.1371/journal.pone.0118255.g001

extract showed lowest antioxidant capacity (IC₅₀ 11.23 mg/ml). In DPPH assay, methanol extract showed highest antioxidant capacity with an IC₅₀ value of 0.52 mg/ml and hexane extract showed lowest antioxidant capacity with 10.94 mg/ml IC₅₀ value. In nitric oxide assay, methanol extract exhibited highest antioxidant capacity (IC₅₀ 4.31 mg/ml) and acetonitrile extract showed lowest antioxidant capacity (IC₅₀ 61.11 mg/ml). In Lipid peroxidation assay, methanol extract was found to possess highest antioxidant capacity (IC₅₀ 0.06 mg/ml) and water extract showed lowest antioxidant capacity (IC₅₀ 1.38 mg/ml). In all antioxidant assays, the methanol

Table 1. Antioxidant capacities of *S. porticalis* extracts^a.

Antioxidant capacity	SWE	SME	SAE	SHE
FRAP (mM Fe (II)/g extract)	32.69 ± 3.86	171.49 ± 19.25*	137.70 ± 11.16* [#]	18.91 ± 1.46* [#] [§]
ABTS IC ₅₀ (mg/ml)	37.71 ± 0.09	3.48 ± 0.11*	24.73 ± 0.09* [#]	4.68 ± 0.11* [#] [§]
β-carotene IC ₅₀ (mg/ml)	0.43 ± 0.01	0.20 ± 0.00*	11.23 ± 0.10* [#]	0.33 ± 0.02* [#] [§]
DPPH IC ₅₀ (mg/ml)	8.06 ± 0.10	0.52 ± 0.01*	5.23 ± 0.11* [#]	10.94 ± 0.08* [#] [§]
NO IC ₅₀ (mg/ml)	9.54 ± 0.09	4.31 ± 0.11*	61.11 ± 0.11* [#]	31.47 ± 0.08* [#] [§]
Lipid peroxidation IC ₅₀ (mg/ml)	1.38 ± 0.11	0.06 ± 0.01*	0.42 ± 0.00* [#]	0.33 ± 0.01* [#] [§]

^aMean ± SD of three replicates.

SWE: *S. porticalis* water extract, SME: *S. porticalis* methanol extract, SAE: *S. porticalis* acetonitrile extract, SHE: *S. porticalis* n-hexane extract.

p < 0.05

*compared with SWE

[#]compared with SME

[§]compared with SAE

doi:10.1371/journal.pone.0118255.t001

Table 2. Total phenolic (TPC), flavanoid (TFC) and proanthocyanidin content (TPAC) in *S. porticalis* extracts from Indian trans-Himalaya^a.

Phytochemicals	SWE	SME	SAE	SHE
TPC (mg GAE/ g)	14.90 ± 1.26	96.70 ± 9.64*	21.42 ± 1.07**	23.72 ± 2.5*#
TFC (mg QE/g)	13.82 ± 1.36	37.59 ± 2.80*	34.76 ± 2.99*	35.30 ± 3.02*
TPAC (mg CE/g)	25.51 ± 2.25	122.55 ± 10.12*	78.36 ± 5.78**	94.79 ± 7.74*#§

^aMean ± SD of three replicates.

SWE: *S. porticalis* water extract, SME: *S. porticalis* methanol extract, SAE: *S. porticalis* acetonitrile extract, SHE: *S. porticalis* n-hexane extract.

p < 0.05

*compared with SWE

#compared with SME

§compared with SAE

doi:10.1371/journal.pone.0118255.t002

extract of *S. porticalis* was found to exhibit significantly higher (*p* < 0.05) antioxidant capacities than all other studied extracts (Table 1).

Total phenolic content was predominant in methanol extract (96.70 mg GAE/g extract) and lowest in water extract (14.90 mg GAE/ g extract). Similarly, total flavonoid content rendered highest value in methanol extract (37.59 mg QE/g extract) and lowest in water extract (13.82 mg QE/g extract). Total proanthocyanidin content was also highest in methanol extract (122.55 mg CE/g extract) and minimum in water extract (25.51 mg CE/g extract). The methanol extract of *S. porticalis* was found to exhibit significantly higher (*p* < 0.05) total phenolics than all other extracts (Table 2).

GC/MS chemometric profile of methanol extract

As described previously, in course of investigation for the preliminary screening with respect to antioxidant capacities and total phenolic content of different extracts of *S. porticalis*, the highest




Fig 2. GC/MS chemometric profile of *S. porticalis* methanol extract.

doi:10.1371/journal.pone.0118255.g002

Table 3. Algal chemotypes identified in methanol extract of *S. porticalis* by GC/MS.

S. No.	Peak RT (min)	Peak area	Peak area (%)	Compound detected	Mol. Formula	Mol. Wt.	CAS No	
1	15.78	156723	0.20	Unknown	-	-	-	-
2	17.00	257237	0.34	Dihydroactinidiolide	C ₁₁ H ₁₆ O ₂	180	15356–74–8	Antibacterial; antiproliferative; cytotoxic
3	19.24	335417	0.43	1,5-hexadien-3-ol, 3-methyl-6-(methylthio)-1-(2,6,6-trimethyl-1-cyclohexen-1-yl)	C ₁₇ H ₂₈ OS	280	97369–78–3	-
4	20.53	422013	0.55	Loliolide	C ₁₁ H ₁₆ O ₃	196	5989–02–6	Strong anti-repellent; antioxidant; protective effects against H ₂ O ₂ -induced cell damage
5	21.16	6137430	8.05	8-Hexadecyne	C ₁₆ H ₃₀	222	19781–86–3	Used in the preparation of fatty alcohols with potential for application in commercial products like as bio-diesel, lubricants, greases and cosmetics
6	21.76	7509928	9.82	5,8,11,14,17-Eicosapentaenoic acid, methyl ester, (all-Z)-	C ₂₁ H ₃₂ O ₂	316	2734–47–6	Hypolipidemic action; lipid-lowering effect; enhancing effect on hepatic biliary secretion; anti-arteriosclerotic
7	21.95	791387	1.02	3-Heptadecen-5-yne, (Z)-	C ₁₇ H ₃₀	234	74744–55–1	-
8	22.31	9034647	12.01	Hexadecanoic acid, methyl ester	C ₁₇ H ₃₄ O ₂	270	112–39–0	Antioxidant; nematocide; pesticide; lubricant; antiandrogenic; flavor, hemolytic 5-alpha reductase inhibitor; hypocholesterolemic
9	23.88	5526076	7.26	4,7,10,13,16,19-Docosahexaenoic acid, methyl ester, (all-Z)-	C ₂₃ H ₃₄ O ₂	342	2566–90–7	-
10	24.43	16103379	21.32	Ethyl Linoleolate	C ₂₀ H ₃₆ O ₂	308	544–35–4	Antioxidant; lightens UV-induced skin pigmentation; anti-inflammatory; competitive inhibitor of prostaglandin production; wounds healing properties; effective anti-acne agent
11	24.56	13175492	18.22	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296	102608–53–7	Used in manufacturing synthetic vitamins E and K; an ingredient of fragrances; end applications include soap, detergent, beauty care product, household product; cancer preventive properties
12	26.29	945579	1.23	Ethyl-5,8,11,14,17-icosapentaenoate	C ₂₂ H ₃₄ O ₂	330	84494–70–2	It has lipid-lowering or hypolipidemic effects, provides protection against atherosclerosis and its complications, cardio-protective functions
13	39.67	14709007	19.55	Stigmasta-5,24(28)-dien-3-ol, (3β)-	C ₂₉ H ₄₈ O	412	18472–36–1	Androgenic; angiogenic; anorexic; antiadenomic; antiandrogenic; antibacterial; anticancer; antiedemic; antiestrogenic; antifeedant; antifertility; antigonadotrophic; antihyperlipoproteinaemic; anti-inflammatory, antileukemic

doi:10.1371/journal.pone.0118255.t003

antioxidant capacities as well as the total phenolic content was achieved by the methanol extract. Henceforth, it was further evaluated for GC/MS chemometric profiling. GC/MS chromatograms of methanol extract of *S. porticalis* as per aforementioned experimental procedure showed various peaks indicating the presence of different cluster of phytochemotypes in the extract (Fig. 2).

The methanol extract of *S. porticalis* revealed the presence of thirteen different chemotypes which were characterized and identified by comparison of their mass fragmentation patterns with the similar compounds present in the NIST database. Of these thirteen chemotypes, ethyl linoleolate (21.32%), stigmasta-5,24(28)-dien-3-ol, (3β) (19.55%), 3,7,11,15-tetramethyl-2-hexadecen-1-ol (18.22%), methyl palmitate (hexadecanoic acid, methyl ester) (12.01%),

5,8,11,14,17-eicosapentaenoic acid, methyl ester, (all-Z) (9.82%), 8-hexadecyne (8.05%), and 4,7,10,13,16,19-docosahexaenoic acid, methyl ester, (all-Z) (7.26%) were found to be the major constituents. In addition, ethyl-5,8,11,14,17-icosapentaenoate (1.23%), 3-heptadecen-5-yne, (Z) (1.02%), (-)-loliolide (0.55%), 1,5-hexadien-3-ol, 3-methyl-6-(methylthio)-1-(2,6,6-trimethyl-1-cyclohexen-1-yl) (0.43%), dihydroactinidiolide (0.34%) and an unknown compound (0.20%) were detected in trace amount (Table 3, Fig. 2). The identified compounds were further investigated for their biological activities and most of them were found to possess diverse range of positive pharmacological and therapeutic functions [66–85] (Table 3).

Cytotoxic effects of methanol extract

Cell cytotoxicity by SRB assay. In HepG2 cells, the methanol extract was found to be non-toxic at concentration level 43.90 µg/ml (63.32% growth at 43.90 µg/ml extract concentration after 72 h). The GI₅₀ and TGI of this extract were 86 µg/ml and 183 µg/ml respectively for HepG2 cells. The % growth at highest concentration level (333.33 µg/ml) was recorded as -39.91. In case of RKO cells, the methanol extract was non-toxic at concentration level of 98.77 µg/ml (79.56% growth at 98.77 µg/ml extract concentration after 72 h). The methanol extract of the alga was found to cross the LC₉₉, LC₅₀, TGI and GI₅₀ line at 333, 205, 179 and 154 µg/ml respectively for RKO cells. All these results have been depicted in Fig. 3 and Table 4.

Cell morphology by microscopic analysis. In HepG2 cells, after 24, 48 and 72 h of extract treatment, it was found to be non-toxic at concentration level 148.15, 98.77 and 98.77 µg/ml respectively whereas the extract showed dead cells at 222.22 and 333.33 µg/ml concentration after 24, 48 and 72 h along with stressed condition at concentration 148.15 µg/ml after 48 and 72 h. In morphological visualization after 24, 48 and 72 h of methanol extract treatment, it was observed as non-toxic at concentration level of 148.15, 98.77 and 148.15 µg/ml respectively for RKO cells. The extract showed dead cells at 222.22 and 333.33 µg/ml concentration after 24, 48 and 72 h (for concentration 222.22 µg/ml after 24 h treatment there was partially arrested RKO cells). All these results have been depicted in Fig. 4.

Effect of algae methanol extract following hypoxia induced oxidative stress

GSH, GSSG and GSH/GSSG level. GSH level in rat blood decreased significantly following exposure to hypobaric hypoxia for 7 days compared to the normoxic group. *S. porticalis* methanol extract administration during hypoxic exposure significantly increased GSH level as compared to the vehicle treated animals of 7 days hypoxic group. However, the GSH level in extract treated group was significantly lower when compared with the normoxic group (Fig. 5A).

GSSG level in rat blood increased significantly following 7 days hypoxic exposure in comparison with the normoxic group. Administration of *S. porticalis* methanol extract to animals during hypoxic exposure significantly caused decrease in the GSSG level as compared to 7 days vehicle treated hypoxic group. However, the GSSG level in extract treated group was significantly higher when compared with the normoxic group (Fig. 5B).

From our results it was clearly observed that there was significant decrease in the GSH level with a concomitant increase in GSSG on animals with 7 days of exposure to hypobaric hypoxia when compared to the normoxic group. The GSH/GSSG value also increased significantly in the 7 days hypoxic animals treated with extract in comparison with the vehicle treated animals of 7 days hypoxic group. However, GSH/GSSG values were found to be reduced significantly in both 7 days hypoxic rats treated with vehicle and extract when compared with the normoxic rats (Fig. 5C).




Fig 3. Cytotoxic effect of *S. porticalis* methanol extract on HepG2 and RKO cells.

doi:10.1371/journal.pone.0118255.g003

Table 4. Cytotoxicity of methanol extract of *S. porticalis* against HepG2 and RKO cells.

Concentration (µg/ml)	Control (C)			HepG2 cells			RKO cells		
	After 24 h	After 48 h	After 72 h	After 24 h	After 48 h	After 72 h	After 24 h	After 48 h	After 72 h
333.33	nt	nt	nt	d	d	d	d	d	d
222.22	nt	nt	nt	dfp	dep	dep	gapf	dsp	dfp
148.15	nt	nt	nt	nt	ga	ga	nt	gapf	nt
98.77	nt	nt	nt	nt	nt	nt	nt	nt	nt
65.84	nt	nt	nt	nt	nt	nt	nt	nt	nt
43.90	nt	nt	nt	nt	nt	nt	nt	nt	nt
29.26	nt	nt	nt	nt	nt	nt	nt	nt	nt

nt: non-toxic, ga: growth arrested, gapf: growth arrested 50%, dfp: death 50%, dsp: death 70%, dep: death 80%, d: dead

doi:10.1371/journal.pone.0118255.t004




Fig 4. Microscopic images (10×10X) showing cell growth and morphology after 24 h treatment of *S. porticalis* extract in HepG2 and RKO cancer cells. a. HepG2 cells with good growth, **b.** HepG2 cells with partially arrested or 50% arrested growth, **c.** HepG2 cells with arrested growth, **d.** HepG2 dead cells, **e.** RKO cells with good growth, **f.** RKO cells with partially arrested or 50% arrested growth, **g.** RKO cells with arrested growth, **h.** RKO dead cells.

doi:10.1371/journal.pone.0118255.g004

Estimation of ROS. There was a significant increase in ROS generation in rats following 7 day of hypoxia when compared with the normoxic group. The *S. porticalis* extract treated hypoxic rats showed significant reduction in ROS generation in comparison to the hypoxic vehicle group. However, the free radical generation in *S. porticalis* extract treated hypoxic group was significantly higher when compared with the normoxic group (Fig. 5D).

Discussion

Algae are exposed to very high light intensity and oxygen concentration during their life cycle and growth that support generation of free radicals and other potent oxidizers. The resistance of algal cellular and structural components against this oxidative challenge indicates that the intracellular bioactive antioxidants may be responsible for the preventive role as an intrinsic mechanism for sustenance [86]. In the trans-Himalayan cold desert, high ultraviolet (UV) and infra-red (IR) irradiation in the thin atmosphere, along with freezing temperature (chilling stress) cause unique abiotic stress for algal survival and this atmospheric stressor could be helpful for the production of various antioxidant compounds. Research investigations by Duval et al. (2000) [87] and Kováčik et al. (2010) [88] revealed that the phenolic substances in algal biomass increased upon exposure to UV, which had a positive correlation to its antioxidant properties.

S. porticalis is a group of microalga belonging to the family Zygnemataceae. Based on the diverse target compounds, growth rate, cultivation simplicity, greater biodiversity and other allied factors, the micro-algal community represent a hitherto untapped resource of natural antioxidants [21]. The benefit of antioxidant consumption is generally associated with consuming different plants, fruits, vegetables and other related natural food sources but little emphasis has been put forward towards microalgae consumption for corresponding health effects. Although a number of previous reports are available on the ethnobotany, medicinal properties, antioxidant capacity, bioactive chemical composition, cytotoxicity etc. of trans-Himalayan medicinal plants and lichens [40–42,89–91], no scientific investigation is documented on the algal




Fig 5. Changes in serum GSH, GSSG, GSH/GSSG and ROS level following administration of *S. porticalis* methanol extract during exposure to hypobaric hypoxia. a. Changes in reduced glutathione level, b. Changes in the level of oxidised glutathione, c. Changes in GSH/GSSG level, d. Changes in reactive oxygen species generation. * denotes $p \leq 0.05$ when compared to normoxia, # denotes $p \leq 0.05$ when compared to normoxia + vehicle, \$ denotes $p \leq 0.05$ when compared to hypoxia + vehicle.

doi:10.1371/journal.pone.0118255.g005

species having medicinal and therapeutic effect and bioactive phytochemicals from this region as far as our information goes.

The antioxidant capacity of plant extracts have been recognized by different redox reactions by several workers such as radical scavenging and inhibitory mechanism, prevention of chain initiation, binding of transition metal ion catalysts, reductive ability etc. [92–94] and it also depends on the hydrophilic or lipophilic nature of the components present in the extracts [40]. Therefore, in order to study the antioxidant processes, the present investigation utilized a range of solvents with varied polarity to access and better understand the trend of antioxidant properties of the different solvent extracts in various antioxidant assay system. We have also monitored the total content of different phytochemicals like proanthocyanidin, polyphenol, flavonoid present in the different extracts in order to determine the relation of phenolic content with the antioxidant capacities. Recent reports by other researchers have described a significant positive correlation between the antioxidant capacities with total phenolic content in microalgal species [21,23], which give support to our results.

Polyphenol, flavonoid, flavonol, proanthocyanidin, alkaloid, terpenoid, steroid, sterol etc. are the class of bioactive compounds of botanical origin that possess strong antioxidant and therapeutic potential [95] and these secondary metabolites are the crucial regulators of plant growth and development in stressful and unfavorable environment. The harsh climatic conditions such as severe cold, aridity, water scarcity, UV radiation etc. in the Indian trans-Himalayan cold desert result in the production and accumulation of anti-stress compounds and secondary metabolites that are known for their antioxidant, medicinal and therapeutic properties and these bioactive compounds could be responsible for the biological activities [87,96,97]. Our results were in concordance with previous investigations where *Spirogyra* extracts showed positive biological activities in terms of antioxidant capacity, antimicrobial activity, antihemolytic activity, antihyperglycemic and antihyperlipidemic properties and renoprotective effect and these activities were attributed to diverse group of bioactive phytochemicals present in this alga [36,37,98–101]. The methanol extract exhibited significantly higher antioxidant capacities and total polyphenol content as compared to n-hexane, acetonitrile and water extracts. Henceforth, we aimed towards the chemometric profiling of the methanol extract by GC/MS technique for its detail chemical characterization. In this investigation our primary objective was the screening of *S. porticalis* extracts on the basis of antioxidant capacities, radical scavenging activities and phenolic profile. The extract showing highest antioxidant potential and phenolic attributes were taken for evaluation of biological activity. Therefore, thorough chemometric profiling was necessary to identify the major bioactive phytochemotypes present in the methanol extract possessing antioxidant properties. To validate the traditional uses, pharmacological actions and therapeutic potential of medicinal plants and other botanicals, metabolic profiling by high-throughput technologies have become an important strategy in medicinal chemistry and natural product research [102–104]; however, GC/MS remains a technique of choice for characterization of volatile compounds present in natural products. The GC/MS chemometric fingerprinting of methanol extract of *S. porticalis* showed thirteen bioactive chemotypes having putative phyto-pharmaceutical importance. The major phytochemical groups namely fatty acid esters, sterols, unsaturated alcohols, alkynes etc. were found to be the dominant components and these are well known for their positive biological functions and could be responsible for the high antioxidant capacities observed in methanol extract of *S. porticalis*.

From the present study, it was established that the methanol extract of *S. porticalis* have promising antioxidant potential and diverse bioactive phytochemicals and it can be utilized as a potential natural antioxidant for various purposes. However, the cytoprotective effects of the algal extract in cell culture system is required to be investigated for further validation of the prophylactic and therapeutic potential and *in vitro* antioxidant capacity of this natural biore-source [105]. We therefore extended our investigation towards the evaluation of cytotoxicity of the methanol extract of *S. porticalis* on human hepatocellular carcinoma HepG2 and colon carcinoma RKO cells.

From the cytotoxicity tests in both cell lines, our results have clearly described the fact that GI_{50} of *S. porticalis* methanol extract was 86 $\mu\text{g/ml}$ for HepG2 cells, which was well below 100 $\mu\text{g/ml}$ and according to the NCI (National Cancer Institute) guideline, this extract could act as a potent anticancer drug for these cells. For RKO cell line, the GI_{50} value was measured as 154 $\mu\text{g/ml}$ that was above 100 $\mu\text{g/ml}$, but at 333.33 and 205 $\mu\text{g/ml}$ concentration the extract was found to approach the LC_{100} and LC_{50} . In contrast to this result, the methanol extract never approached the LC_{50} up to concentration level 333.33 $\mu\text{g/ml}$ in HepG2 cells. Hence, it was clear that this extract was acted like an anticancerous agent for both HepG2 and RKO cells under our study.

After confirming the cytotoxic and anticarcinogenic effects of *S. porticalis* methanol extract in HepG2 and RKO cell lines, we further carried out the *in vivo* study to evaluate the biological

activity of the algal extract in animal model. Here, *S. porticalis* methanol extract provided protection from the hypoxia-induced oxidative stress and damage to organ systems. It was also found to accelerate the onset of adaptative changes in rats following hypoxic exposure. The bioactive phytochemicals of the extract could have acted synergistically to produce an optimum protective effect under hypoxic stress. The scarcity of oxygen causes impairment of electron transport chain in mitochondria and incomplete reduction of oxygen in hypoxic condition bring about elevated production of free radicals. Previous reports have shown increased level of oxidative stress during exposure to hypobaric hypoxia [106,107] and depleted endogenous antioxidant enzyme level [108,109]. The results of the present investigation also corroborate with the previous reports, where sharp rise in free radical levels and consequent decline of endogenous antioxidant status was seen in animals following exposure to hypobaric hypoxia for 7 days. Administration of *S. porticalis* methanol extract during exposure to hypobaric hypoxia decreased free radical levels and oxidized glutathione (GSSG) concentration, and also increased concentration of endogenous reduced glutathione (GSH) and GSH/GSSG ratio. The antioxidant action of the algal extract could be attributed to the rich content of polyphenols, flavonoids, proanthocyanidins, fatty acid esters, sterols, unsaturated alcohols and other bioactive components. The algal extract induced augmentation of endogenous antioxidant GSH level and GSH/GSSG ratio in blood proves the efficacy of bioactive components in modulating glutathione biosynthesis and stabilization under hypoxic condition. However, the underlying mechanism of such regulatory effects of *S. porticalis* methanol extract on endogenous glutathione biosynthesis needs further investigation at cellular and molecular level.

As discussed previously, a number of studies have been conducted by several investigators to assess the medicinal and therapeutic effects of freshwater algae and diverse group of bioactive compounds were discovered from this vital natural resource. However, there is limited information available regarding the biological and medicinal potential of *S. porticalis* that limits the scope to compare our results with previous reports specific to this alga [110]. To the best of our knowledge, this is the first report on the medicinal potential of *S. porticalis* from the trans-Himalayan Ladakh region that could lead us to develop novel therapeutic compounds from this hitherto untapped unique bioresource.

Conclusion

The freshwater filamentous alga *Spirogyra porticalis* from the Indian trans-Himalayan cold desert of Ladakh had been identified by morpho-anatomical characteristics and the antioxidant capacities of the algal extracts reported for the first time. The present study revealed high antioxidant capacities and considerable amount of total phenolic compounds in *S. porticalis* extracts which could be used as an efficient natural source of antioxidants. Amongst all tested extracts, the methanol extract was found to possess highest antioxidant capacity and phenolic compounds and this extract was further investigated for GC/MS chemometric profiling, cytotoxic action on human carcinoma cell lines and *in vivo* biological action on animal model. In total, the accession of the study lies within methanol extract of *S. porticalis* which was found to have high antioxidant properties, phenolic content, diverse bioactive phytochemotypes, anti-carcinogenic effects along with *in vivo* anti-hypoxic and anti-stress potential. It is evident from the study that the trans-Himalayan alga *S. porticalis* is a reservoir of novel compounds and could be sustainably utilized as an immense wealth for the discovery of novel drugs against a variety of human ailments, specifically oxidative stress-induced disorders.

Acknowledgments

The authors express their sincere gratefulness to Dr. Manas K. Mandal, Director General-Life Sciences, DRDO Headquarters for his wholehearted support. The authors also acknowledge the contribution of Debasmita Ghosh Dhar, Maharaja Manindra Chandra College, Kolkata for her sincere help in copyediting and proofreading of the manuscript.

Author Contributions

Conceived and designed the experiments: JK PD ABT DG OPC DKU KT RA MRS RBS. Performed the experiments: JK PD ABT DG DKU KT. Analyzed the data: JK PD ABT DG DKU KT. Contributed reagents/materials/analysis tools: JK PD ABT DG DKU KT. Wrote the paper: PD JK ABT. Concise review and laboratory facility: DG OPC DKU RA MRS RBS.

References

1. Prasad TK, Anderson MD, Martin BA, Stewart CR (1994) Evidence for chilling-induced oxidative stress in maize seedlings and regulatory role for hydrogen peroxide. *Plant Cell* 6: 65–74. PMID: [12244221](#)
2. Dai F, Huang Y, Zhou M, Zhang G (2009) The influence of cold acclimation on antioxidative enzymes and antioxidants in sensitive and tolerant barley cultivars. *Biol Plantarum* 53: 257–262.
3. Kohen R, Nyska A (2002) Oxidation of biological systems: oxidative stress phenomena, antioxidants, redox reactions, and method for their quantification. *Toxicol Pathol* 30: 620–650. PMID: [12512863](#)
4. Brieger K, Schiavone S, Miller FJ Jr, Krause KH (2012) Reactive oxygen species: from health to disease. *Swiss Med Wkly* 142: w13659. doi: [10.4414/smw.2012.13659](#) PMID: [22903797](#)
5. Ismail A, Hong TS (2002) Antioxidant activity of selected commercial seaweeds. *Malays J Nutr* 8: 167–177. PMID: [22692475](#)
6. Kumar KS, Ganesan K, Subba Rao PV (2008) Antioxidant potential of solvent extracts of *Kappaphycus alvarezii* (Doty) Doty—An edible seaweed. *Food Chem* 107: 289–295.
7. Cardozo KH, Guaratini T, Barros MP, Falcão VR, Tonon AP, et al. (2007) Metabolites from algae with economical impact. *Comp Biochem Physiol C Toxicol Pharmacol* 146: 60–78. PMID: [16901759](#)
8. Plaza M, Cifuentes A, Ibañez E (2008) In the search of new functional food ingredients from algae. *Food Sci Tech* 19: 31–39.
9. Mungmai L, Jiranusornkul S, Peerapompisal Y, Sirithunyalug B, Leelapompisid P (2014) Extraction, Characterization and Biological Activities of Extracts from Freshwater Macroalga [*Rhizoclonium hieroglyphicum* (C.Agardh) Kützing] Cultivated in Northern Thailand. *Chiang Mai J Sci* 41: 14–26.
10. Freile-Pelegrín Y, Robledo D (2014) Bioactive Phenolic Compounds from Algae. In: Hernández-Ledesma B, Herrero M (editors), *Bioactive Compounds from Marine Foods: Plant and Animal Sources*, First Edition (Feb), John Wiley & Sons, Ltd, The Atrium, West Sussex, UK. pp. 113–129.
11. Frikha F, Kammoun M, Hammami N, Mchirgui RA, Belbahri L, et al. (2011) Chemical composition and some biological activities of marine algae collected in Tunisia. *Ciencias Marinas* 37: 113–124.
12. Shalaby EA (2011) Algae as promising organisms for environment and health. *Plant Signal Behav* 6: 1338–1350. PMID: [21862867](#)
13. Priyadarshani I, Rath B (2012) Commercial and industrial applications of micro algae—A review. *J Algal Biomass Utiln* 3: 89–100.
14. Dhar P (2013) Autonomic responses in high altitude and adaptogenic efficacy of phytococktail from trans-Himalayan plants. PhD thesis, submitted to Bharathiar University, Coimbatore, India.
15. Hota KB, Hota SK, Dhar P, Chaurasia OP, Srivastava RB (2011) Challeages at high altitude: The limits to human performance. In: Srivastava RB, Selvamurthy W (editors), *Innovations in agro animal technologies*. Satish Serial Publishing House, Delhi. pp. 277–299.
16. Singh B, Chaurasia OP (2000) Medicinal flora of Indian cold desert. *Acta Horti (Belgium)* 523: 65–72.
17. Chaurasia OP, Ahmed Z, Ballabh B (2007) *Ethnobotany and Plant of Trans-Himalaya*. Satish Serial Publishing House, Delhi, India. pp. 1–544. doi: [10.1007/s12033-007-0062-9](#) PMID: [23934597](#)
18. Ballabh B, Chaurasia OP (2007) Traditional medicinal plants of cold desert Ladakh-used in treatment of cold, cough and fever. *J Ethnopharmacol* 112: 341–349. PMID: [17459623](#)

19. Ballabh B, Chaurasia OP, Ahmed Z, Singh SB (2008) Traditional medicinal plants of cold desert Ladakh-used against kidney and urinary disorders. *J Ethnopharmacol* 118: 331–339. doi: [10.1016/j.jep.2008.04.022](https://doi.org/10.1016/j.jep.2008.04.022) PMID: [18550306](https://pubmed.ncbi.nlm.nih.gov/18550306/)
20. Ballabh B, Chaurasia OP (2009) Medicinal plants of cold desert Ladakh used in the treatment of stomach disorders. *Indian Journal of Traditional Knowledge* 8: 185–190.
21. Li HB, Cheng KW, Wong CC, Fan KW, Chen F, et al. (2007) Evaluation of antioxidant capacity and total phenolic content of different fractions of selected microalgae. *Food Chem* 102: 771–776.
22. Natrah FMI, Yusoff FM, Shariff M, Abas F, Mariana NS (2007) Screening of Malaysian indigenous microalgae for antioxidant properties and nutritional value. *J Appl Phycol* 19: 711–718.
23. Hajimahmoodi M, Faramarzi MA, Mohammadi N, Soltani N, Oveisi MR, et al. (2010) Evaluation of antioxidant properties and total phenolic contents of some strains of microalgae. *J Appl Phycol* 22: 43–50.
24. Rodriguez-Garcia I, Guil-Guerrero J (2008) Evaluation of the antioxidant activity of three microalgal species for use as dietary supplements and in the preservation of foods. *Food Chem* 108: 1023–1026.
25. Chacón-Lee TL, González-Mariño GE (2010) Microalgae for “healthy” foods-possibilities and challenges. *CRFSFS* 9: 655–675.
26. Lee SH, Lee JB, Lee KW, Jeon YJ (2010) Antioxidant properties tidal pool microalgae, *Halochlorococcum porphyrae* and *Oltamanssiellopsis unicellularis* from Jeju Island, Korea. *Algae* 25: 45–56.
27. Goiris K, Muylaert K, Fraeye I, Foubert I, Brabanter JD, et al. (2012) Antioxidant potential of microalgae in relation to their phenolic and carotenoid content. *J Appl Phycol* 24: 1477–1486.
28. Gad HA, El-Ahmady SH, Abou-Shoer MI, Al-Azizi MM (2013) Application of chemometrics in authentication of herbal medicines: a review. *Phytochem Anal* 24: 1–24. doi: [10.1002/pca.2378](https://doi.org/10.1002/pca.2378) PMID: [22678654](https://pubmed.ncbi.nlm.nih.gov/22678654/)
29. Sarker SD, Nahar L (2012) Hyphenated techniques and their applications in natural products analysis. *Methods Mol Biol* 864: 301–340. doi: [10.1007/978-1-61779-624-1_12](https://doi.org/10.1007/978-1-61779-624-1_12) PMID: [22367902](https://pubmed.ncbi.nlm.nih.gov/22367902/)
30. Lifongo LL, Simoben CV, Ntie-Kang F, Babiaka SB, Judson PN (2014) A bioactivity versus ethnobotanical survey of medicinal plants from Nigeria, west Africa. *Nat Prod Bioprospect* 4: 1–19. doi: [10.1007/s13659-014-0005-7](https://doi.org/10.1007/s13659-014-0005-7) PMID: [24660132](https://pubmed.ncbi.nlm.nih.gov/24660132/)
31. Lagunin AA, Goel RK, Gawande DY, Pahwa P, Glorizova TA et al. (2014) Chemo- and bioinformatics resources for *in silico* drug discovery from medicinal plants beyond their traditional use: a critical review. *Nat Prod Rep* 31: 1585–1611. doi: [10.1039/c4np00068d](https://doi.org/10.1039/c4np00068d) PMID: [25051191](https://pubmed.ncbi.nlm.nih.gov/25051191/)
32. Cragg GM, Grothaus PG, Newman DJ (2014) New horizons for old drugs and drug leads. *J Nat Prod* 77: 703–723. doi: [10.1021/np5000796](https://doi.org/10.1021/np5000796) PMID: [24499205](https://pubmed.ncbi.nlm.nih.gov/24499205/)
33. Rubió L, Motilva MJ, Romero MP (2013) Recent advances in biologically active compounds in herbs and spices: a review of the most effective antioxidant and anti-inflammatory active principles. *Crit Rev Food Sci Nutr* 53: 943–953. doi: [10.1080/10408398.2011.574802](https://doi.org/10.1080/10408398.2011.574802) PMID: [23768186](https://pubmed.ncbi.nlm.nih.gov/23768186/)
34. Balboa EM, Conde E, Moure A, Falqué E, Domínguez H (2013) In vitro antioxidant properties of crude extracts and compounds from brown algae. *Food Chem* 138: 1764–1785. doi: [10.1016/j.foodchem.2012.11.026](https://doi.org/10.1016/j.foodchem.2012.11.026) PMID: [23411309](https://pubmed.ncbi.nlm.nih.gov/23411309/)
35. Lordan S, Ross RP, Stanton C (2011) Marine bioactives as functional food ingredients: potential to reduce the incidence of chronic diseases. *Mar Drugs* 9: 1056–1100. doi: [10.3390/md9061056](https://doi.org/10.3390/md9061056) PMID: [21747748](https://pubmed.ncbi.nlm.nih.gov/21747748/)
36. Thumvijit T, Thuschana W, Amornlerdpison D, Peerapornpisal Y, Wongpoomchai R (2013) Evaluation of hepatic antioxidant capacities of *Spirogyra neglecta* (Hassall) Kützing in rats. *Interdiscip Toxicol* 6:152–156. doi: [10.2478/intox-2013-0024](https://doi.org/10.2478/intox-2013-0024) PMID: [24678253](https://pubmed.ncbi.nlm.nih.gov/24678253/)
37. Ontawong A, Saowakon N, Vivithanapom P, Pongchaidecha A, Lailerd N, et al. (2013) Antioxidant and renoprotective effects of *Spirogyra neglecta* (Hassall) Kützing extract in experimental type 2 diabetic rats. *Biomed Res Int* 2013: 820786. doi: [10.1155/2013/820786](https://doi.org/10.1155/2013/820786) PMID: [23862157](https://pubmed.ncbi.nlm.nih.gov/23862157/)
38. Compère P (1983) Some algae from Kashmir and Ladakh, W. Himalayas. In: *Bulletin de la Société Royale de Botanique de Belgique / Bulletin van de Koninklijke Belgische Botanische Vereniging*. T. 116, Fasc. 2. Royal Botanical Society of Belgium. pp. 141–160.
39. Yoshimura Y, Kohshima S, Ohtani S (1997) A community of snow algae on a Himalayan glacier: Change of algal biomass and community structure with altitude. *Arctic and Alpine Research* Vol. 29, No. 1 (Feb., 1997). INSTAAR, University of Colorado. pp. 126–137.
40. Dhar P, Bajpai PK, Tayade AB, Chaurasia OP, Srivastava RB, et al. (2013a) Chemical composition and antioxidant capacities of phytococktail extracts from trans-Himalayan cold desert. *BMC Complement Altern Med* 13: 259. doi: [10.1186/1472-6882-13-259](https://doi.org/10.1186/1472-6882-13-259) PMID: [24098968](https://pubmed.ncbi.nlm.nih.gov/24098968/)

41. Dhar P, Tayade AB, Kumar J, Chaurasia OP, Srivastava RB, et al. (2013b) Nutritional profiling of phytococktail from Trans-Himalayan plants. PLoS ONE 8: e83008. doi: [10.1371/journal.pone.0083008](https://doi.org/10.1371/journal.pone.0083008) PMID: [24376624](https://pubmed.ncbi.nlm.nih.gov/24376624/)
42. Ballabh B, Chaurasia OP (2007) Traditional medicinal plants of cold desert Ladakh—used in treatment of cold, cough and fever. J Ethnopharmacol 112: 341–349. PMID: [17459623](https://pubmed.ncbi.nlm.nih.gov/17459623/)
43. Prescott GW (1951) Algae of the western great lakes area. Cranbrook Institute of Science, Michigan. pp. 318. Pl, 75, Fig. 10.
44. Ikram EHK, Eng KH, Jalil AMM, Ismail A, Idris S, et al. (2009) Antioxidant capacity and total phenolic content of Malaysian underutilized fruits. J Food Compos Anal 22: 388–393.
45. Re R, Pellegrini N, Proteggente A, Pannala A, Yong M, et al. (1999) Antioxidant activity applying an improved ABTS radical cation decolourisation assay. Free Rad Biol Med 26: 1231–1237. PMID: [10381194](https://pubmed.ncbi.nlm.nih.gov/10381194/)
46. Brand-Williams W, Cuvelier ME, Berset C (1995) Use of a free radical method to evaluate antioxidant activity. LWT-Food Sci Technol 28: 25–30.
47. Zhang D, Hamauz Y (2004) Phenolics, ascorbic acid, carotenoids and antioxidant activity of broccoli and their changes during conventional and microwave cooking. Food Chem 88: 503–509.
48. Dapkevicius A, Venskutonis R, van Beek TA, Linsen JPH (1998) Antioxidant activity of extracts obtained different isolation procedures from some aromatic herbs grown in Lithuania. J Sci Food Agric 77: 140–146.
49. Wang ZH, Hsu CC, Yin MC (2009) Antioxidative characteristics of aqueous and ethanol extracts of glossy privet fruit. Food Chem 112: 914–918.
50. Ohkawa H, Ohishi N, Yagi K (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. Anal Biochem 95: 351–358. PMID: [36810](https://pubmed.ncbi.nlm.nih.gov/36810/)
51. Ruberto G, Baratta MT, Deans SG, Dorman HJ (2000) Antioxidant and antimicrobial activity of *Foeniculum vulgare* and *Crithmum maritimum* essential oils. Planta Med 66: 687–693. PMID: [11199122](https://pubmed.ncbi.nlm.nih.gov/11199122/)
52. Gao X, Ohlander M, Jeppsson N, Björk L, Trajkovski V (2000) Changes in antioxidant effects and their relationship to phytonutrients in fruits of sea buckthorn (*Hippophae rhamnoides* L.) during maturation. J Agric Food Chem 48: 1485–1490. PMID: [10820047](https://pubmed.ncbi.nlm.nih.gov/10820047/)
53. Ordonez AAL, Gomez JD, Vattuone MA, Isla MI (2006) Antioxidant activities of *Sechium edule* (Jacq.) Swart extracts. Food Chem 97: 452–458.
54. Sun JS, Tsuang YW, Chen IJ, Huang WC, Hang YS, et al. (1998) An ultra-weak chemiluminescence study on oxidative stress in rabbits following acute thermal injury. Burns 24: 225–231. PMID: [9677025](https://pubmed.ncbi.nlm.nih.gov/9677025/)
55. Tayade AB, Dhar P, Kumar J, Sharma M, Chauhan RS, et al. (2013) Chemometric profile of root extracts of *Rhodiola imbricata* Edgew. with hyphenated gas chromatography mass spectrometric technique. PLoS ONE 8: e52797. doi: [10.1371/journal.pone.0052797](https://doi.org/10.1371/journal.pone.0052797) PMID: [23326358](https://pubmed.ncbi.nlm.nih.gov/23326358/)
56. Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, et al. (1990) New colorimetric cytotoxicity assay for anticancer-drug screening. J Natl Cancer Inst 82: 1107–1112. PMID: [2359136](https://pubmed.ncbi.nlm.nih.gov/2359136/)
57. Rubinstein LV, Shoemaker RH, Paull KD, Simon RM, Tosini S, et al. (1990) Comparison of *in vitro* anticancer-drug-screening data generated with a tetrazolium assay versus a protein assay against a diverse panel of human tumor cell lines. J Natl Cancer Inst 82: 1113–1118. PMID: [2359137](https://pubmed.ncbi.nlm.nih.gov/2359137/)
58. Monks A, Scudiero D, Skehan P, Shoemaker R, Paull K, et al. (1991) Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human tumor cell lines. J Natl Cancer Inst 83: 757–766. PMID: [2041050](https://pubmed.ncbi.nlm.nih.gov/2041050/)
59. Boyd MR, Paull KD (1995) Some practical considerations and applications of the National Cancer Institute *in vitro* anticancer drug discovery screen. Drug Develop Res 34: 91–109.
60. Boyd MR, Paull KD, Rubinstein L (1992) Data display and analysis strategies for the NCI disease-oriented *in vitro* drug screen. In: Valeriote FA, Corbett T, Baker L (editors), Antitumor drug discovery and development. Amsterdam, The Netherlands: Kluwer Academic Publishers. pp. 11–34.
61. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG (2010) Improving Bioscience Research Reporting: The ARRIVE Guidelines for Reporting Animal Research. PLoS Biol 8: e1000412. doi: [10.1371/journal.pbio.1000412](https://doi.org/10.1371/journal.pbio.1000412) PMID: [20613859](https://pubmed.ncbi.nlm.nih.gov/20613859/)
62. Austin Institutional Animal Care and Use Committee Guidelines (2013) Guidelines #003 for the Use of Carbon Dioxide (CO₂) for Rodent Euthanasia. The University of Texas, Austin, USA. pp. 1–4. doi: [10.1097/ACC.0b013e31829c6877](https://doi.org/10.1097/ACC.0b013e31829c6877) PMID: [25629060](https://pubmed.ncbi.nlm.nih.gov/25629060/)
63. Riley V (1960) Adaptation of orbital bleeding technique to rapid serial blood studies. Proc Soc Exp Biol Med 104: 751–754. PMID: [13741689](https://pubmed.ncbi.nlm.nih.gov/13741689/)

64. LeBel CP, Ali SF, McKee M, Bondy SC (1990) Organometal-induced increases in oxygen reactive species: the potential of 2',7'-dichlorofluorescein diacetate as an index of neurotoxic damage. *Toxicol Appl Pharmacol* 104: 17–24. PMID: [2163122](#)
65. Hissin PJ, Hilf R (1976) A fluorimetric method for determination of oxidized and reduced glutathione in tissue. *Anal Biochem* 74: 214–226. PMID: [962076](#)
66. De Rosa S, Kamenarska Z, Stefanov K, Dimitrova-Konaklieva S, Najdenski C, et al. (2003) Chemical composition of *Corallina mediterranea* Areschoug and *Corallina granifera* Eil. et Soland. *Z Naturforsch C* 58: 325–332. PMID: [12872923](#)
67. Pubchem website. Available: <http://pubchem.ncbi.nlm.nih.gov/compounds/12311356#section=BioAssay-Results>. Accessed 2014 Oct 31.
68. Okunade AL, Wiemer DF (1985) (-) Loliolide, an ant-repellent compound from *Xanthoxylum setulosum*. *J Nat Prod* 48: 472–473.
69. Okada N, Shirata K, Niwano M, Koshino H, Uramoto M (1994) Immuno-suppressive activity of a monoterpene from *Eucommia ulmoides*. *Phytochemistry* 37: 281–282.
70. Kimura J, Maki N (2002) New loliolide derivatives from the brown alga *Undaria pinnatifida*. *J Nat Prod* 65: 57–58. PMID: [11809066](#)
71. Ponnamma SU, Manjunath DK (2012) GC-MS analysis of phytocomponents in the methanolic extract of *Justicia wynaadensis* (NEES) T. ANDERS. *Int J Pharm Bio Sci* 3: 570–576.
72. Kashiwaga T, Mikagia E, Mekuria DB, Borua AD, Tebayashia S, et al. (2005) Ovipositional Deterrent on Mature Stage of Sweet Pepper, *Capsicum annuum*, against *Liriomyza trifolii* (Burgess). *Z Naturforsch* 60: 739–742. PMID: [16320617](#)
73. Mizuguchi K, Yano T, Ishibashi M, Masada A, Mizota M, et al. (1993) Ethylall-cis-5,8,11,14,17-icosapentaenoate modifies the biochemical properties of rat very low-density lipoprotein. *Eur J Pharmacol* 235: 221–227. PMID: [8508903](#)
74. Yano T, Mizuguchi K, Takasugi K, Tanaka Y, Sato M (1995) Effects of ethyl all-cis-5,8,11,14,17-icosapentaenoate on low density lipoprotein in rabbits. *Yakugaku Zasshi* 115: 843–851. PMID: [8531063](#)
75. Yano T, Kawano H, Yamashita M, Mizuguchi K, Mochizuki H, et al. (1997) Effects of ethyl-all-cis-5,8,11,14,17-icosapentaenoate (EPA-E), pravastatin and their combination on serum lipids and intimal thickening of cuff-sheathed carotid artery in rabbits. *Life Sci* 61: 2007–2015. PMID: [9366508](#)
76. Satoh N, Shimatsu A, Kazuhiko K, Sakane N, Yamada K, et al. (2007) Purified Eicosapentaenoic Acid Reduces Small Dense LDL, Remnant Lipoprotein Particles, and C-Reactive Protein in Metabolic Syndrome. *Diabetes care* 30: 144–146. PMID: [17192349](#)
77. Percot A, Yalcin A, Aysel V, Erduğan H, Dural B, et al. (2009) Loliolide in marine algae. *Nat Prod Res* 23: 460–465. doi: [10.1080/14786410802076069](https://doi.org/10.1080/14786410802076069) PMID: [19296390](#)
78. Jelenko C, Wheeler ML, Anderson AP, Callaway BD, McKinley JC (1975) Studies in burns: XIV, Healing in burn wounds treated with Ethyl Linoleate alone or in combination with selected topical antibacterial agents. *Ann Surg* 182: 562–566. PMID: [1190861](#)
79. Hølmer G, Kristensen G, Søndergaard E, Dam H (1962) The influence of ethyl linoleate and ethyl linolenate on polyenoic fatty acids and cholesterol in tissues of chicks. *Zeitschrift für Ernährungswissenschaft* 2: 223–228. doi: [10.1107/S0108767309007235](https://doi.org/10.1107/S0108767309007235) PMID: [19349661](#)
80. Hutsell TC, Quackenbush FW (1967) Reduction of atherogenicity of natural fats by small additions of ethyl linoleate in the diet of the rat. *Lipids* 2: 342–344. PMID: [17805763](#)
81. Charakida A, Charakida M, Chu AC (2007) Double-blind, randomized, placebo-controlled study of a lotion containing triethyl citrate and ethyl linoleate in the treatment of acne vulgaris. *Br J Dermatol* 157: 569–574. PMID: [17635508](#)
82. Downing DT, Stewart ME, Wertz PW, Strauss JS (1986) Essential fatty acids and acne. *J Am Acad Dermatol* 14: 221–225. PMID: [2936775](#)
83. Ando H, Ryu A, Hashimoto A, Oka M, Ichihashi M (1998) Linoleic acid and alpha-linolenic acid lightens ultraviolet-induced hyperpigmentation of the skin. *Arch Dermatol Res* 290: 375–381. PMID: [9749992](#)
84. Ando H, Wen ZM, Kim HY, Valencia JC, Costin GE, et al. (2006) Intracellular composition of fatty acid affects the processing and function of tyrosinase through the ubiquitin-proteasome pathway. *Biochem J* 394: 43–50. PMID: [16232122](#)
85. Ziboh VA, Miller CC, Cho Y (2000) Metabolism of polyunsaturated fatty acids by skin epidermal enzymes: generation of antiinflammatory and antiproliferative metabolites. *Am J Clin Nutr* 71: 361–366. PMID: [10617998](#)
86. Burritt D, Larkindale J, Hurd CL (2002) Antioxidant metabolism in the intertidal red seaweed *Stictosiphonia arbuscula* following desiccation. *Planta* 215: 829–838. PMID: [12244449](#)

87. Duval B, Shetty K, Thomas WH (2000) Phenolic compounds and antioxidant properties in the snow alga *Chlamydomonas nivalis* after exposure to UV light. *J Appl Phycol* 11: 559–566.
88. Kováčik J, Klejduš B, Backor M (2010) Physiological responses of *Scenedesmus quadricauda* (Chlorophyceae) to UV-A and UV-C light. *Photochem Photobiol* 86: 612–616. doi: [10.1111/j.1751-1097.2010.00708.x](https://doi.org/10.1111/j.1751-1097.2010.00708.x) PMID: [20202162](https://pubmed.ncbi.nlm.nih.gov/20202162/)
89. Bajpai PK, Warghat AR, Dhar P, Kant A, Srivastava RB, et al. (2014) Variability and fruit color effect on antioxidant activity, phenolics, flavonoids, anthocyanin and proanthocyanidin content in *Morus alba* L. fruit from the trans-Himalaya, India. *LWT-Food Sci Technol* 59: 981–988.
90. Tayade AB, Dhar P, Sharma M, Chauhan RS, Chaurasia OP, et al. (2013) Antioxidant capacities, phenolic contents, and GC/MS analysis of *Rhodiola imbricata* Edgew. root extracts from Trans-Himalaya. *J Food Sci* 78: 402–410.
91. Korekar G, Dolkar P, Singh H, Srivastava RB, Stobdan T (2014) Variability and the genotypic effect on antioxidant activity, total phenolics, carotenoids and ascorbic acid content in seventeen natural population of Seabuckthorn (*Hippophae rhamnoides* L.) from trans-Himalaya. *LWT-Food Sci Technol* 55: 157–162.
92. Frankel EN, Meyer AS (2000) The problems of using one-dimensional methods to evaluate multifunctional food and biological antioxidants. *J Sci Food Agric* 80: 1925–1941.
93. Zheng W, Wang SY (2001) Antioxidant activity and phenolic compounds in selected herbs. *J Agric Food Chem* 49: 5165–5170. PMID: [11714298](https://pubmed.ncbi.nlm.nih.gov/11714298/)
94. Huang D, Ou B, Prior L (2005) The chemistry behind antioxidant capacity assays. *J Agric Food Chem* 53: 1841–1856. PMID: [15769103](https://pubmed.ncbi.nlm.nih.gov/15769103/)
95. Jeong WY, Jin JS, Cho YA, Lee JH, Park S, et al. (2011) Determination of polyphenols in three *Capsicum annuum* L. (bell pepper) varieties using high-performance liquid chromatography–tandem mass spectrometry: Their contribution to overall antioxidant and anticancer activity. *J Sep Sci* 34: 2967–2974. doi: [10.1002/jssc.201100524](https://doi.org/10.1002/jssc.201100524) PMID: [21898818](https://pubmed.ncbi.nlm.nih.gov/21898818/)
96. Solecka D (1997) Role of phenylpropanoid compounds in plants responses to different stress factors. *Acta Physiol Plant* 19: 257–268.
97. Chandrasekhar MJN, Praveen B, Nanjan MJ, Suresh B (2006) Chemoprotective effect of *Phyllanthus maderaspatensis* in modulating cisplatin-induced nephrotoxicity and enotoxicity. *Pharm Biol* 44: 100–106. PMID: [16689610](https://pubmed.ncbi.nlm.nih.gov/16689610/)
98. Khalid MN, Shameel M (2012) Studies on the phytochemistry and biological activity of *Spirogyra rhizoides* (Chlorophycota). *Pak J Bot* 44: 1815–1820.
99. Naik AA, Hemavani C, Thippeswamy B (2012) Evaluation of antimicrobial property of *Spirogyra* species. *International Multidisciplinary Research Journal* 2: 13–15.
100. Nabavi SF, Nabavi SM, Ebrahimzadeh M, Jafari N, Yazdanpanah S (2012) Biological Activities of Freshwater Algae, *Spirogyra singularis* Nordstedt. *J Aquat Food Prod Technol* 22: 58–65.
101. Junthip R, Amornlerdpison D, Chimsook T (2013) Phytochemical screening, antioxidant activity and total phenolic content of *Spirogyra* spp. *Advanced Materials Research* 699: 693–697.
102. Shellie RA, Poynter SD, Li J, Gathercole JL, Whittock SP, et al. (2009) Varietal characterization of hop (*Humulus lupulus* L.) by GC–MS analysis of hop cone extracts. *J Sep Sci* 32: 3720–3725. doi: [10.1002/jssc.200900422](https://doi.org/10.1002/jssc.200900422) PMID: [20029908](https://pubmed.ncbi.nlm.nih.gov/20029908/)
103. Roessner U, Luedemann A, Brust D, Fiehn O, Linke T, et al. (2001) Metabolic profiling allows comprehensive phenotyping of genetically or environmentally modified plant systems. *Plant Cell* 13: 11–29. PMID: [11158526](https://pubmed.ncbi.nlm.nih.gov/11158526/)
104. Gullberg J, Jonsson P, Nordstrom A, Sjostrom M, Moritz T (2004) Design of experiments: an efficient strategy to identify factors influencing extraction and derivatization of *Arabidopsis thaliana* samples in metabolomic studies with gas chromatography/mass spectrometry. *Anal Biochem* 331: 283–295. PMID: [15265734](https://pubmed.ncbi.nlm.nih.gov/15265734/)
105. Kumar J, Dhar P, Tayade AB, Gupta D, Chaurasia OP, et al. (2014) Antioxidant capacities, phenolic profile and cytotoxic effects of mountaneous saxicolous lichens from Trans-Himalayan cold desert of Ladakh. *PLoS ONE* 9: e98696. doi: [10.1371/journal.pone.0098696](https://doi.org/10.1371/journal.pone.0098696) PMID: [24937759](https://pubmed.ncbi.nlm.nih.gov/24937759/)
106. Nakanishi K, Tajima F, Nakamura A, Yagura S, Ookawara T, et al. (1995) Effects of hypobaric hypoxia on antioxidant enzymes in rats. *J Physiol* 489: 869–876. PMID: [8788950](https://pubmed.ncbi.nlm.nih.gov/8788950/)
107. Maiti P, Singh SB, Sharma AK, Muthuraju S, Banerjee PK, et al. (2006) Hypobaric hypoxia induces oxidative stress in rat brain. *Neurochem Int* 49: 709–716. PMID: [16911847](https://pubmed.ncbi.nlm.nih.gov/16911847/)
108. Sairam M, Sharma SK, Dipti P, Pauline T, Kain AK, et al. (1998) Effect of hypobaric hypoxia on immunofunction in albino rats. *Int J Biometeorol* 42: 55–59. PMID: [9780847](https://pubmed.ncbi.nlm.nih.gov/9780847/)

109. Ilavazhagan G, Sridharan K, Sharma SK, Bansal A, Prasad D, et al. (1996) Role of Vitamin C and E under high altitude combat stress, Defence Institute of Physiology and Allied Sciences, Report No. DIPAS/26/96.
110. Guiry MD (2014) *Spirogyra porticalis* (Müller OF) Dumortier, 1822. In: Guiry MD, Guiry GM. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M.D. Guiry). Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=609396> on 2014-05-22.