

Experimental quantification of long distance dispersal potential of aquatic snails in the gut of migratory birds

Casper H.A. van Leeuwen, Gerard van der Velde, Bart van Lith & Marcel Klaassen

Supporting Information

Table S2: The number of intact, damaged and viable snails retrieved after 24 hours for the two experiments combined.

Species	Snails ingested	Damaged shells (average \pm 95%CI)	Intact shells (average \pm 95%CI)	Viable snails (average \pm 95%CI)
<i>Hydrobia (Peringia) ulvae</i>	12×200 + 14×300 = 6600	11.4 (\pm 7.0): 4.33%	11.2 (\pm 8.5): 4.23%	0.8 (\pm 1.2): 0.32%
<i>Potamopyrgus antipodarum</i>	12×200 + 14×300 = 6600	4.8 (\pm 3.6): 1.88%	13.9 (\pm 10.7): 5.47%	-
<i>Potamopyrgus antipodarum</i> + macrophytes	14×150 = 2100	0.8 (\pm 0.5): 0.52%	5.3 (\pm 3.7): 3.52%	-
<i>Bithynia leachii</i>	12×100 = 1200	-	1.0 (\pm 1.3): 1.0%	-
<i>Bathyomphalus contortus</i>	12×100 = 1200	-	-	-