

<b>Supplemental Table S1. Neutralization sensitivity of pseudovirus constructed with envelopes from subject 107747</b>				
	<b>Neutralization titer (IC<sub>50</sub>) obtained with HIV+ serum</b>			
<b>Clone / Serum</b>	<b>T500105</b>	<b>T500107</b>	<b>T500208</b>	<b>Z23</b>
<b>016</b>	<b>47</b>	<b>4144</b>	<b>417</b>	<100
<b>041</b>	<40	<b>1260</b>	<b>136</b>	<100
<b>048 wtS</b>	<b>80</b>	<b>5734</b>	<b>439</b>	<b>168</b>
<b>066</b>	<b>84</b>	<b>1553</b>	<b>175</b>	<b>111</b>
<b>072</b>	<b>43</b>	<b>1553</b>	<b>217</b>	<100
<b>073</b>	<b>56</b>	<b>4525</b>	<b>431</b>	<100
<b>077</b>	<b>68</b>	<b>4549</b>	<b>387</b>	<b>110</b>
<b>083</b>	<b>67</b>	<b>1617</b>	<b>143</b>	<b>157</b>
<b>091</b>	<b>56</b>	<b>1866</b>	<b>133</b>	<b>109</b>
<b>092 wtR</b>	<b>45</b>	<b>964</b>	<b>120</b>	<100
<b>JRCSE</b>	<40	<b>1490</b>	<40	<b>300</b>
<b>NL43</b>	<b>1597</b>	<b>75</b>	<b>66</b>	<b>4438</b>
<b>aMLV</b>	<40	<40	<40	<100
<p>The neutralizing antibody titer (IC<sub>50</sub>) is defined as the reciprocal of the plasma dilution that produces a 50% inhibition in target cell infection. Values in bold represent significant neutralization titers that are at least three times greater than those observed against the negative control (aMLV). The Envs designated wildtype resistant (wtR) and wildtype sensitive (wtS) are indicated. Envs for both the wtR and wtS isolates were from CCR5-dependent viruses as determined by the Trofile® assay (Monogram Biosciences).</p>				