

**Supporting Information Table S2. Immunogenicity of WHc-Pv-78 VLPs.**

**Antibody Titer (1/dilution)**

Immunogen	Adjuvant	Dose	1°/2°	α-WHc	α-Pv1a	IFA
<b>WHc-Pv-78</b> n=9	Alum	100 µg	1°	80.5K	24.5K	-
		100 µg	2°	9.5x10 <sup>6</sup>	2.2x10 <sup>6</sup>	8.1K
<b>WHc-Pv-78</b> n=3	IFAd	20 µg	1°	250K	250K	-
		10 µg	2°	3x10 <sup>6</sup>	3x10 <sup>6</sup>	-
<b>WHc-Pv-78</b> n=8	IFAd	100 µg	1°	3x10 <sup>6</sup>	9x10 <sup>6</sup>	-
		100 µg	2°	6.5x10 <sup>7</sup>	1.5x10 <sup>8</sup>	72.9K
<b>WHcAg</b> n=10	IFAd	100 µg	1°	6x10 <sup>6</sup>	0	-
		100 µg	2°	2.3x10 <sup>8</sup>	0	0
<b>Mab 2F2</b> ( mg/ml)	-	-	-	0	45x10 <sup>6</sup>	-

Mean endpoint dilution Ab titers shown. 1°, primary; 2°, secondary antisera. IFA assay used Pb/Pv dry sporozoites.