

S1 Table. Comorbidities and additional laboratory analyses in anorexia nervosa patients.

Parameter	Low anxiety (n=32)	High anxiety (n=32)	Missing data	p
<i>Psychiatric/psychological comorbidities</i>			0	
Alcohol abuse	0	2		n.C.
Major depressive disorder	5	7		n.C.
Adjustment disorder	2	1		n.C.
Somatoform disorder	3	0		n.C.
Panic disorder	1	0		n.C.
Obsessive-compulsive disorder	1	3		n.C.
Personality disorder/accentuation	1	2		n.C.
<i>Somatic Comorbidities</i>			0	
Hypothyreosis	0	2		n.C.
Epilepsy	1	1		n.C.
Ascites	0	3		n.C.
Pericardial effusion	3	5		n.C.
Chronic heart failure	1	0		n.C.
Interstitial nephritis	1	0		n.C.
Urinary stasis	0	1		n.C.
Gastritis	3	0		n.C.
Bronchial asthma	1	0		n.C.
Sinusitis	0	1		n.C.
Locomotor system disorders	1	0		n.C.
Osteopenia	1	1		n.C.

<i>Additional laboratory analyses</i>				
TSH (mU/l)	1.7 (1.3/3.0)	2.6 (2.0/3.4)	2	0.03^a
CRP (mg/l)	0.4 (0.2/0.9)	0.2 (0.2/0.7)	2	0.32 ^a
Glucose (mg/dl)	74.6 ± 10.4	70.0 ± 12.1	1	0.12 ^b
Cortisol (nmol/l)	866.4 ± 363.1	845.9 ± 364.4	5	0.83 ^b

Statistical analyses: Normal distribution was determined by Kolmogorov-Smirnov test. Differences between groups: ^a Mann-Whitney-U-test, data expressed as median (25. percentile/75. percentile); ^b t-test, data are expressed as mean ± standard deviation. Significant differences are displayed in bold. Abbreviation: n.a., not applicable; n.c., not calculated.