

| Patient Number | Experiment | Visit Date Sample tested | Months viral load <50 copies/ml | Coinfection history | | | | | | | | | | |
|------------------|-----------------------|--------------------------|---------------------------------|--|--|-----------------------------|-------------------------|--|----------------------|-----------|---|------------------------------|--|------------------------------|
| | | | | Cytomegalovirus (CMV) | Epstein-Barr virus (EBV) | Hepatitis B virus (HBV) | Hepatitis C virus (HCV) | Herpes simplex virus (HSV) | Influenza A | Rotavirus | Respiratory syncytial virus (RSV) | Varicella zoster virus (VZV) | | |
| Figure 4 - Pt 28 | Cross-sectional Study | 2/19/08 | 13 | | PCR + (2/15/2006, 1/27/06), Negative 4/28/05 | | | | | | | | | |
| Figure 4 - Pt 29 | Cross-sectional Study | 8/5/08 | 17 | | | | | | | | Isolate parainfluenza type 1 (7/14/09), EIA + (1/29/08) | | | |
| Figure 4 - Pt 30 | Cross-sectional Study | 1/21/10 | 41 | | | | | | | | | | | |
| Figure 4 - Pt 31 | Cross-sectional Study | 3/22/11 | 37 | | | | | | PCR + (02/16/11) | | | | | |
| Figure 4 - Pt 32 | Cross-sectional Study | 9/21/10 | 8 | | 7447550 copies/mL (6/15/13), 850 copies/mL (8/15/2012), PCR + (10/22/2009) | | | | | | | | | |
| Figure 4 - Pt 33 | Cross-sectional Study | 12/7/10 | 19 | | | | | | | | | | | |
| Figure 4 - Pt 34 | Cross-sectional Study | 6/22/06 | 7 | | PCR + (3/15/06, 11/3/05), PCR - (7/20/05) | | | | Culture + (03/22/06) | | | | | |
| Figure 4 - Pt 35 | Cross-sectional Study | 4/23/03 | 17 | | | | | PCR + (1/4/07), RNA IU: 24900000 IU/mL (10/5/11), 13300000 IU/mL (6/14/11), 64,300 IU/mL (11/4/08), 257000 IU/mL(1/4/07) RNA Quantitative: 67230000 copies/mL (10/5/11), 35910000 copies/mL (6/14/11), 173610 copies/mL (11/4/08), 693900 copies/mL (1/4/07) | | | | | | |
| Figure 4 - Pt 36 | Cross-sectional Study | 5/14/03 | 6 | | | | | | | | Isolate Influenza A 12/26/03 | | | |
| Figure 4 - Pt 37 | Cross-sectional Study | 10/13/09 | 9 | | | | | | | | | | | |
| Figure 4 - Pt 38 | Cross-sectional Study | 2/8/05 | 7 | | | HBs Ag: Reactive (02/08/05) | | | Culture + (7/20/04) | | | | | |
| Figure 4 - Pt 39 | Cross-sectional Study | 4/28/09 | 12 | | | | | | | | | | | |
| Figure 4 - Pt 40 | Cross-sectional Study | 12/16/04 | 8 | | PCR + (7/22/04), ND (8/18/09) PCR + (1/13/10), PCR Low + (11/12/09, 6/3/09, 2/11/09, 7/1/08, 3/13/08, 12/18/07), ND (10/23/08, 9/25/07) | | | | | | | | | |
| Figure 4 - Pt 41 | Cross-sectional Study | 12/18/07 | 6 | PCR Low + (10/23/08, 6/5/07, 5/10/07, 4/11/07, 3/29/07), PCR - (7/1/08, 7/3/07, 2/11/09) PCR + (4/25/07) | | | | | | | | | | PCR + (7/1/08), PCR - 4/4/07 |

| Patient Number | Experiment | Visit Date Sample tested | Months viral load <50 copies/ml | Coinfection history | | | | | | | | | |
|------------------|-----------------------|--------------------------|---------------------------------|-----------------------|-----------------------------------|---|---|----------------------------|------------------------------|-----------|-----------------------------------|------------------------------|--|
| | | | | Cytomegalovirus (CMV) | Epstein-Barr virus (EBV) | Hepatitis B virus (HBV) | Hepatitis C virus (HCV) | Herpes simplex virus (HSV) | Influenza A | Rotavirus | Respiratory syncytial virus (RSV) | Varicella zoster virus (VZV) | |
| Figure 4 - Pt 44 | Cross-sectional Study | 11/8/05 | 16 | | | HBs Ag: Reactive (1/13/09), Repeat reactive (8/16/05, 9/3/04, 2/17/04), Hbe Ag: Reactive 9/6/02, DNA Quantitative: None detected (6/23/05), 927 copies/mL (9/3/04), 37,600,000 diluted 1:1000 copies/mL (2/17/04), 2,500,000,000 copies/mL dilute 1:10,000 (11/24/03), >500,000 copies/mL (9/6/02), 322,000 copies/mL (6/26/02) | | | Culture + (10/15/04) | | | | |
| Figure 4 - Pt 45 | Cross-sectional Study | 5/5/09 | 17 | | PCR + (3/11/10) | | | | | | | | |
| Figure 4 - Pt 46 | Cross-sectional Study | 11/13/08 | 12 | | | | | | | | | | |
| Figure 4 - Pt 47 | Cross-sectional Study | 2/24/10 | 11 | | | | | | | | | | |
| Figure 4 - Pt 48 | Cross-sectional Study | 12/18/08 | 36 | | PCR + (3/15/06, 11/3/05, 7/20/05) | | | | Culture + (3/22/06, 3/15/06) | | | | |
| Figure 4 - Pt 49 | Cross-sectional Study | 7/20/10 | 40 | | | | RNA Quantitative: 143640 copies/mL (10/18/12), 10,530,000 copies/mL (8/23/05), <1,620 copies/mL (6/7/05) None detected (5/3/09), RNA IU: 53200 IU/mL (10/18/12), 3,900,000 IU/mL, <600 IU/mL (6/7/05), none detected (5/3/09) | | PCR + (10/2/12) | | | | |
| Figure 4 - Pt 50 | Cross-sectional Study | 3/29/10 | 6 | | | | | | Culture + (7/24/07) | | | | |
| Figure 4 - Pt 51 | Cross-sectional Study | 4/28/09 | 8 | | | | | | | | | | |
| Figure 4 - Pt 52 | Cross-sectional Study | 12/3/10 | 21 | | PCR Low + (12/4/08) | | | | | | | | |
| Figure 4 - Pt 53 | Cross-sectional Study | 3/29/11 | 113 | | | HBs Ag: + (1/5/10, 7/2/09), Repeat reactive known positive (4/6/04, 2/11/03), DNA Quantitative: None detected (9/24/13, 5/4/10, 2/26/08, 6/26/03, 10/3/02), <169 copies/mL (3/29/11), 58 copies/mL (1/5/10), DNA IU: <29 IU/mL (3/29/11), <10 IU/mL (1/5/10) | | | | | | | |