

Table S1: Distribution of genetic loci in the Genus *Salmonellae*

Species (strain)	<i>srcA</i>^a	SPI-1	SPI-2	<i>pipB2</i>	<i>sseL</i>
<i>S. enterica</i>					
Typhimurium (LT2)	+ (100)	+	+	+ (100)	+ (100)
Typhimurium (SL1344)	+ (100)	+	+	+ (100)	+ (100)
Typhi (Ty2)	+ (97)	+	+	+ (95)	+ (93)
Typhi (CT18)	+ (97)	+	+	+ (95)	+ (93)
Paratyphi A	+ (97)	+	+	+ (98)	+ (93)
Choleraesuis	+ (99)	+	+	+ (98)	+ (93)
Newport (SL254)	+ (100)	+	+	+ (98)	+ (96)
Dublin (CT02021853)	+ (99)	+	+	+ (99)	+ (93)
Heidelberg (SL476)	+ (99)	+	+	+ (99)	+ (93)
Heidelberg (SL486)	+ (99)	+	+	+ (99)	+ (93)
Agona (SL483)	+ (98)	+	+	+ (94)	+ (94)
Schwarzengrund (SL480)	+ (98)	+	+	+ (94)	+ (91)
Schwarzengrund (CVM19633)	+ (98)	+	+	+ (94)	+ (91)
Enteriditis	+ (98)	+	+	+ (99)	+ (94)
Gallinarum	+ (98)	+	+	+ (99)	+ (94)
<i>S. bongori</i>					
	-	+	-	-	-

^a +, present; -, absent (% identity to SL1344 orthologue)