

CORRECTION

Correction: Neurally Encoding Time for Olfactory Navigation

The *PLOS Computational Biology* Staff

An error was introduced during the typesetting process. There are two typographical errors in the x-axis label in [Fig 2](#). The publisher apologizes for these errors.

Please view the correct version of [Fig 2](#) here:



click for updates

OPEN ACCESS

Citation: The *PLOS Computational Biology* Staff (2016) Correction: Neurally Encoding Time for Olfactory Navigation. *PLoS Comput Biol* 12(1): e1004742. doi:10.1371/journal.pcbi.1004742

Published: January 27, 2016

Copyright: © 2016 The PLOS Computational Biology Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 2. Odor plume PLIF videos taken at 15 locations. Instantaneous odor concentration (expressed as % of source concentration) at (A) $x = 50$ cm, (B) $x = 100$ cm, (C) $x = 150$ cm, (D) $x = 200$ cm, (E) $x = 250$ cm from the source along the odor plume centerline, and (F) $y = 5$ cm, (G) $y = 10$ cm from the odor plume centerline at $x = 150$ cm.

doi:10.1371/journal.pcbi.1004742.g001

Reference

1. Park IJ, Hein AM, Bobkov YV, Reidenbach MA, Ache BW, Principe JC (2016) Neurally Encoding Time for Olfactory Navigation. *PLoS Comput Biol* 12(1): e1004682. doi:[10.1371/journal.pcbi.1004682](https://doi.org/10.1371/journal.pcbi.1004682) PMID: [26730727](https://pubmed.ncbi.nlm.nih.gov/26730727/)