CORRECTION

Correction: Exonic Splicing Mutations Are More Prevalent than Currently Estimated and Can Be Predicted by Using *In Silico* Tools

Omar Soukarieh, Pascaline Gaildrat, Mohamad Hamieh, Aurélie Drouet, Stéphanie Baert-Desurmont, Thierry Frébourg, Mario Tosi, Alexandra Martins

There is an error in [Fig 4](#). The symbol for correlation coefficient " ρ " should be replaced with " r ", to match the description in the main text and Figure legend. Please see the correct [Fig 4](#) here.

There is a typographical error in the last sentence of the paragraph following [Fig 4](#) (Results section): ' $\Delta\Delta\text{HZEI}$ ' should be replaced with ' ΔHZEI '. The correct sentence is: "Moreover, given their lack of comprehensiveness and global quantitation, the performance of these two *in silico* tools could not be statistically analyzed, nor properly compared to that of $\Delta\text{tESRseq}$, ΔHZEI , $\Delta\Psi$ or EX-SKIP."

OPEN ACCESS

Citation: Soukarieh O, Gaildrat P, Hamieh M, Drouet A, Baert-Desurmont S, Frébourg T, et al. (2016) Correction: Exonic Splicing Mutations Are More Prevalent than Currently Estimated and Can Be Predicted by Using *In Silico* Tools. PLoS Genet 12(4): e1005971. doi:10.1371/journal.pgen.1005971

Published: April 6, 2016

Copyright: © 2016 Soukarieh et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Fig 4. Correlation analysis between exon 10 inclusion levels and results obtained from new ESR-dedicated bioinformatics tools. (A), (B) and (C) refer to results obtained with $\Delta tESRseq$ -, ΔHZ_{EI} - and $\Delta \Psi$ -based bioinformatics approaches, respectively, as described under Materials and Methods. Only *MLH1* exon 10 variants located outside the sequences that define the reference splice sites were retained for this analysis as already mentioned in Fig 3. The precise correspondence between each Δ value ($\Delta tESRseq$, ΔHZ_{EI} or $\Delta \Psi$), the level of exon inclusion observed in the pSPL3m-M1e10 minigene assay, and the identity of the corresponding *MLH1* exon 10 variant, is indicated on S1 Table. Correlation coefficients (r) and p-values were determined by performing a Pearson correlation analysis, as described under Materials and Methods.

doi:10.1371/journal.pgen.1005971.g001

Reference

1. Soukarieh O, Gaildrat P, Hamieh M, Drouet A, Baert-Desurmont S, Frébourg T, et al. (2016) Exonic Splicing Mutations Are More Prevalent than Currently Estimated and Can Be Predicted by Using In Silico Tools. PLoS Genet 12(1): e1005756. doi: [10.1371/journal.pgen.1005756](https://doi.org/10.1371/journal.pgen.1005756) PMID: [26761715](https://pubmed.ncbi.nlm.nih.gov/26761715/)