

S1 Table. Strains.

	Genotype or description	Source
<u><i>C. albicans</i> strains</u>		
CAI4	<i>ura3Δ::imm434/ura3Δ::imm434</i>	[1]
CAF2-1	<i>ura3Δ::imm434/URA3</i>	[1]
RM1000	<i>ura3::imm434/ura3::imm434 iro1/iro1::imm434 his1::hisG/his1::hisG</i>	[2]
SK47	as CAI4 but <i>dom34Δ::FRT/dom34Δ::FRT</i>	this work
JH47-1 /-2	as CAI4 but <i>dom34Δ ::FRT/dom34Δ ::FRT ura3Δ ::imm434/URA3</i>	this work
SK24	as CAI4 but <i>pmt1Δ::hisG/pmt1Δ::hisG dom34Δ::FRT/dom34Δ::FRT</i>	this work
JH24-4	<i>pmt1Δ::hisG/pmt1Δ::hisG dom34Δ::FRT/ dom34Δ::FRT ura3Δ ::imm434/URA3</i>	this work
JH5-3-1	as SK47 but <i>pmt5Δ-hisG-URA3-hisG/pmt5Δ-hisG</i>	this work
P15-274	as SK47 but <i>pmt1Δ::hisG/pmt1Δ::hisG pmt5Δ-URA3/pmt5Δ-hisG</i>	this work
P15-274-1	as SK47 but <i>pmt1Δ::hisG/pmt1Δ::hisG pmt5Δ-hisG/pmt5Δ-hisG</i>	this work
CIS23	as CAI4 but <i>PMT1/PMT1^{HA}-SAT1</i>	this work
CAP1-3121	as CAI4, but <i>pmt1Δ::hisG /pmt1Δ::hisG</i>	[3]
SPCa2	as CAI4, but <i>pmt1Δ::hisG /pmt1Δ::hisG ura3Δ::imm434/URA3</i>	[4]
SPCa10	as CAI4, but <i>pmt5Δ::hisG/pmt5Δ::hisG ura3Δ::imm434/URA3</i>	[4]
CPP1121	as CAI4, but <i>pmt1Δ::hisG/pmt1Δ::hisG pmt6Δ-hisG/pmt6Δ-hisG</i>	[4]
SPCa8	as CAI4, but <i>pmt6Δ::hisG/pmt6Δ::hisG ura3Δ::imm434/URA3</i>	[4]
SPCa4	as CAI4, but <i>PMT2/pmt2Δ::hisG ura3Δ::imm434/URA3</i>	[4]
SPCa6	as CAI4, but <i>pmt4Δ::hisG/pmt4Δ::hisG ura3Δ::imm434/URA3</i>	[4]
JH-Ca1-1 /-2	as CAI4, but <i>DOM34/DOM34^{HA}-URA3</i>	this work
<u><i>S. cerevisiae</i> strains</u>		
YE449	<i>MATα leu2 ura3-52 prb1-112 pep4-3 cir⁰</i>	[5]
YE449 <i>pmt1</i>	as YE449, but <i>pmt1::URA3</i>	this work
YE449 <i>pmt2</i>	as YE449, but <i>pmt2::LEU2</i>	this work
YE449 <i>pmt1 pmt2</i>	as YE449, but <i>pmt1::URA3 pmt2::LEU2</i>	this work

YE449 <i>dom34</i>	as YE449, but <i>dom34Δ::kanMX4</i>	this work
W21, W32	as YE449, but <i>pmt1Δ::URA3 dom34Δ::kanMX4</i>	this work
W12, W14	as YE449, but <i>pmt1Δ::URA3 yil001wΔ::kanMX4</i>	this work
M577	as YE449, but mutant isolate after EMS mutagenesis	[5]
M577 <i>pmt1</i>	as M577, but <i>pmt1::URA3</i>	this work
M577 <i>pmt2</i>	as M577, but <i>pmt2::LEU2</i>	this work
M577 <i>pmt1 pmt2</i>	as M577, but <i>pmt1::URA3 pmt2::LEU2</i>	this work
RC130	<i>MATa/MATα his1/his1 [KIL-K1] (K⁺ R⁺)</i>	R. Chan, unpublished

References

1. Fonzi W, Irwin Y. Isogenic strain construction and gene mapping in *Candida albicans*. *Genetics*. 1993;134: 717-728.
2. Negrodo A, Monteoliva L, Gil C, Pla J, Nombela C. Cloning, analysis and one-step disruption of the *ARG5,6* gene of *Candida albicans*. *Microbiology*. 1997;143: 297-302.
3. Timpel C., Strahl-Bolsinger S, Ziegelbauer K, Ernst JF. Multiple functions of Pmt1p-mediated protein O-mannosylation in the fungal pathogen *Candida albicans*. *J Biol Chem*. 1998;273: 20837-20846.
4. Prill SK-H, Klinkert B, Timpel C, Gale CA, Schröppel K, Ernst JF. *PMT* family of *Candida albicans*: five protein mannosyltransferase isoforms affect growth, morphogenesis and antifungal resistance. *Mol Microbiol*. 2005;55: 546-560.
5. Finck M, Bergmann N, Jansson B, Ernst JF. Defective threonine-linked glycosylation of human insulin-like growth factor in mutants of the yeast *Saccharomyces cerevisiae*. *Glycobiology*. 1996;6: 313-320.