

Editorial

Translating Translational Research into Global Health Gains

The *PLOS Medicine* Editors*

Last month, the United States Supreme Court struck down a patent claim by Myriad Genetics on the *BRCA1* and *BRCA2* genes [1]. The case, *Association for Molecular Pathology v. Myriad Genetics*, considered whether genes isolated in the lab are “products of nature” that may not be patented or are “human-made inventions” that would be eligible for patent protection. In its decision, the Court disallowed the patenting of human genes, writing that “A naturally occurring DNA segment is a product of nature and not patent eligible merely because it has been isolated” [2]. This ruling may well have considerable implications for the cost and delivery of health care, and leads us to consider just how important translational medicine is becoming to medical practice, and how far it still has to go to benefit the global community.

Translational medicine seeks to build upon basic scientific research and to translate scientific findings into medical advances that will improve clinical practice and public health. In the case above, the discovery of variants in the *BRCA1* and *BRCA2* genes that predispose individuals to breast and ovarian cancer led to the development of medical tests based on gene sequencing. Individuals with a strong family history of disease can now be tested, counseled, and potentially treated to prevent the development of disease. Because of Myriad Genetics’ patent on and exclusive test for the *BRCA1* and *BRCA2* genes, individuals seeking this test were paying well above the current cost for gene sequencing; now other companies will be able to offer competing *BRCA* tests that may drive the costs down [3].

The potential of translational medicine to improve clinical care is reflected in two recent studies in *PLOS Medicine*. Pierre Laurent-Puig and colleagues reported on a transcriptome-based classification of colon cancer that groups colon cancers into six molecular subtypes, using clinicopathological variables and common DNA markers that are detectable due to defects in distinct biological pathways [4]. The authors of the study classified colon cancer into six different subtypes, two of which have worse outcomes overall. This classification is a significant improvement over

the previous three-subtype classification, which had little prognostic value. The hope is that this new classification system can be used to help predict prognosis for stage II–III colon cancers and to identify and develop targeted treatments for the different subtypes of disease.

Another recent study published in the journal, by Nathan Ledebner and colleagues, assessed the performance of a diagnostic platform for detecting Gram-positive bacteria in blood cultures, an important step in the diagnosis and treatment of patients with sepsis [5]. Gram-positive bacteria are the most common cause of sepsis, and rapid detection and diagnosis are critical. Currently, standard treatment of sepsis is hampered by long turnaround times to identify bacteria after culture and determine antibiotic susceptibility. In this study, the authors tested the microarray-based Verigene Gram-Positive Blood Culture (BC-GP) Test for 12 different Gram-positive bacteria and genetic markers of antibiotic resistance and demonstrate that the BC-GP Test identified Gram-positive bacteria directly from positive blood culture broths, with high sensitivity and specificity. In addition, the 2-hour turnaround time to both identify bacteria after culture and determine their antibiotic resistance could help clinicians implement rapid, effective treatment for their patients.

The promise of translational medicine has also attracted increased calls for and funding of more translational research. The UK government, for instance, announced plans to increase funding for

translational medicine by 30% [6], while the US National Institutes of Health (NIH) established the National Center for Advancing Translational Sciences, with a US\$575 million dollar investment, aiming to “transform the translational science process so that new treatments and cures for disease can be delivered to patients faster” [7].

With more funding, the amount of translational research is sure to increase, but to realize the potential of translational medicine, research must be of the highest possible quality. In a recent *PLOS Medicine* Essay, Carljin Hooijmans and colleagues point out several challenges in translating animal research to humans, including the poor methodological quality of some animal experiments, insufficient reporting of animal research, and publication bias. These issues may be at least as big a problem for preclinical research as they are for clinical research, where the requirements for study protocols and prospective trial registration help encourage more standardized and transparent methodology. The authors emphasize the importance of conducting systematic reviews of animal research to better support future animal and human studies and conclude that “Improving the quality and translation of animal research requires co-operation from the wider scientific community, journals, researchers, regulators, funding bodies, peer reviewers, and patients” [8]. We support this call to improve the conduct and reporting of animal research and endorse the ARRIVE guidelines for reporting of animal research

Citation: The *PLOS Medicine* Editors (2013) Translating Translational Research into Global Health Gains. *PLoS Med* 10(7): e1001493. doi:10.1371/journal.pmed.1001493

Published: July 30, 2013

Copyright: © 2013 *PLOS Medicine* Editors. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors are each paid a salary by the Public Library of Science, and they wrote this editorial during their salaried time.

Competing Interests: The authors’ individual competing interests are at <http://www.plosmedicine.org/static/editorsInterests.action>. PLOS is funded partly through manuscript publication charges, but the *PLOS Medicine* Editors are paid a fixed salary (their salary is not linked to the number of papers published in the journal).

Abbreviations: GP-BC, Gram-Positive Blood Culture; LMIC, low- and middle-income country; TB, tuberculosis

* E-mail: medicine_editors@plos.org

The *PLOS Medicine* Editors are Virginia Barbour, Jocalyn Clark, Laureen Connell, Rhona MacDonald, Melissa Norton, Amy Ross, Paul Simpson, and Margaret Winker.

Provenance: Written by editorial staff; not externally peer reviewed.

[9]. Further, *PLOS Medicine* now requires authors to submit the ARRIVE checklist with all manuscripts using animal models.

Methodological issues in preclinical research are also addressed in a recent systematic review by Valerie Henderson and colleagues [10]. The authors analyzed existing guidelines for the design of preclinical research, developed an evidence-based list of methodological issues across various disciplines, and provide a researcher-friendly tool to aid in planning research. Implementing such tools can only help improve the state of the science, and ultimately perhaps a single evidence-based tool to aid researchers in the planning, conduct, and reporting of research can be developed.

The potential of translational medicine is also reduced by prohibitive expense, as exemplified in the case of the recently overturned *BRCA1* and *BRCA2* gene patents. If such testing is considered expensive in high-income countries, the tests are out of reach for most individuals in low- and middle-income countries (LMICs). Why then does *PLOS Medicine*, with our focus on global health, include a focus on translational medicine? The potential impact of translational research on global health is not a new concept. In a report from the World Health Organiza-

tion on genomics and global health in 2002, commissioned after the sequencing of the human genome, the authors write, “it is clear that the science of genomics holds tremendous potential for improving health globally” [11]. If anything, LMICs have more to gain from translational medicine, where the lack of state-of-the-art laboratories make the potential for rapid, simple, low-cost diagnostics all the more compelling.

There have been some translational medicine successes for global health recently, one example of which is Xpert MTB/RIF. A PCR-based test built on the GeneXpert platform [12] and contained within a desktop machine that rapidly detects *Mycobacterium tuberculosis*, Xpert MTB/RIF can identify drug resistance in 2 hours [13]. Tuberculosis is one of the world’s most serious infections, and TB cases and drug resistance are both increasing [14]. Effective diagnostic tools for TB or for detecting resistant strains had previously not been available in many highly affected areas, so Xpert MTB/RIF has significantly changed the landscape of TB diagnosis, especially for patients with HIV [15]. Xpert MTB/RIF does not require specialized laboratories or skills, and the rapid diagnosis (about 2 hours vs. weeks to test for drug-resistant

TB with previous tests) is well-suited for endemic settings. However, even after the price was negotiated for high-burden countries, the cost of each machine (about USD\$17,000, with extra costs for the cartridges) remains an impediment to rollout [16].

While GeneXpert and Xpert MTB/RIF are a step toward realizing the potential for translational medicine to improve global health, these advances must be affordable to realistically tackle diseases on a global scale. Governments and funding bodies have invested heavily in global public-private partnerships in the hope that vaccines, diagnostics, and treatments might be developed to benefit the global community. However, these advances must be accessible to all countries regardless of income if translational medicine is to truly improve global health and not simply widen the current inequalities in global health.

Author Contributions

Wrote the first draft of the manuscript: LC MN. Contributed to the writing of the manuscript: VB JC LC RM MN AR PS MW. ICMJE criteria for authorship read and met: VB JC LC RM MN AR PS MW. Agree with manuscript results and conclusions: VB JC LC RM MN AR PS MW.

References

1. Liptak A (June 13, 2013) Justices, 9-0, bar patenting human genes. New York Times online. Available: <http://www.nytimes.com/2013/06/14/us/supreme-court-rules-human-genes-may-not-be-patented.html?hpw>. Accessed 09 July 13.
2. Supreme Court of the United States (2013) Syllabus: Association for Molecular Pathology et al. v. Myriad Genetics, Inc., et al. Certiorari to the United States Court of Appeals for the Federal Circuit. No. 12–398. Available: http://www.supremecourt.gov/opinions/12pdf/12-398_1b7d.pdf. Accessed 09 July 13.
3. Munro D (June 16, 2013) Supreme Court breaks Myriad monopoly. Forbes online. Available: <http://www.forbes.com/sites/danmunro/2013/06/16/supreme-court-breaks-myriad-monopoly/>. Accessed 09 July 13.
4. Marisa L, de Reyniès A, Duval A, Selves J, Gaub MP, et al. (2013) Gene Expression Classification of Colon Cancer into Molecular Subtypes: Characterization, Validation, and Prognostic Value. *PLoS Med* 10(5): e1001453. doi:10.1371/journal.pmed.1001453
5. Buchan BW, Ginocchio CC, Manii R, Cavagnolo R, Pancholi P, et al. (2013) Multiplex Identification of Gram-Positive Bacteria and Resistance Determinants Directly from Positive Blood Culture Broths: Evaluation of an Automated Microarray-Based Nucleic Acid Test. *PLoS Med* 10(7): e1001478. doi:10.1371/journal.pmed.1001478
6. Perks B (2011) UK plans 30% funding bump for translational research. *Nat Med* 17: 394. doi:10.1038/nm0411-394b
7. National Center for Advancing Translational Sciences. Homepage. Available: <http://www.ncats.nih.gov/about/about.html>. Accessed 09 July 13.
8. Hooijmans CR, Ritskes-Hoitinga M (2013) Progress in Using Systematic Reviews of Animal Studies to Improve Translational Research. *PLoS Med* 10(7): e1001482. doi:10.1371/journal.pmed.1001482
9. Kilkenny C, Browne WJ, Cuthill IC, Emerson M, Altman DG (2010) Improving Bioscience Research Reporting: The ARRIVE Guidelines for Reporting Animal Research. *J Pharmacol Pharmacother* 1: 94–99. doi: 10.4103/0976-500X.72351
10. Henderson VC, Kimmelman J, Fergusson D, Grimshaw JM, Hackam DG (2013) Threats to Validity in the Design and Conduct of Preclinical Efficacy Studies: A Systematic Review of Guidelines for In Vivo Animal Experiments. *PLoS Med* 10(7): e1001489. doi:10.1371/journal.pmed.1001489
11. WHO (2002) Genomics and World Health. Report of the Advisory Committee on Health Research. Available: <http://whqlibdoc.who.int/hq/2002/a74580.pdf>. Accessed 09 July 13.
12. Raja S, Ching J, Xi L, Hughes SJ, Chang R, et al. (2005) Technology for automated, rapid, and quantitative PCR or reverse transcription-PCR clinical testing. *Clin Chem* 51: 882–890.
13. Helb D, Jones M, Story E, Boehme C, Wallace E, et al. (2010) Rapid Detection of *Mycobacterium tuberculosis* and Rifampin Resistance by Use of On-Demand, Near-Patient Technology. *J Clin Microbiol* 48: 229–237. doi:10.1128/JCM.01463-09
14. Evans CA (2011) GeneXpert—A Game-Changer for Tuberculosis Control? *PLoS Med* 8(7): e1001064. doi:10.1371/journal.pmed.1001064.
15. Lawn SD, Brooks SV, Kranzer K, Nicol MP, Whitelaw A, et al. (2011) Screening for HIV-Associated Tuberculosis and Rifampicin Resistance before Antiretroviral Therapy Using the Xpert MTB/RIF Assay: A Prospective Study. *PLoS Med* 8(7): e1001067. doi:10.1371/journal.pmed.1001067
16. Vassall A, van Kampen S, Sohn H, Michael JS, John KR, et al. (2011) Rapid Diagnosis of Tuberculosis with the Xpert MTB/RIF Assay in High Burden Countries: A Cost-Effectiveness Analysis. *PLoS Med* 8(11): e1001120. doi:10.1371/journal.pmed.1001120