S2 Table. Risk score and mortality data

Number of survivors, patients dying, observed and predicted mortality risk for **A**) 'full' mortality predictor score (based on the regression coefficients) and **B**) simplified mortality risk score.

Risk score	Proportion			Observed	Predicted
category	of patients	Survived	Died	mortality	Mortality
0-5	1.3	4	0	0.0	7.7
6-10	14.0	38	6	13.6	12.1
11-15	27.3	68	18	20.9	18.4
16-20	17.8	44	12	21.4	27.0
21-25	17.8	36	20	35.7	37.8
26-30	17.8	27	29	51.8	49.9
31-35	3.2	3	7	70.0	62.0
>35	1.0	1	2	66.7	72.8
Low	15.2	42	6	12.5	9.7
Medium	45.1	112	30	21.1	23.0
High	39.7	67	58	46.4	45.3

A)

Mortality risk groups were defined as low risk (10 points or fewer), medium risk (11 to 20 points) or high risk (more than 20 points)

B)

Risk score or risk					
score	% of			Observed	Predicted
category	patients	Survived	Died	mortality	Mortality
0	1.3	4	0	0.0	5.6
1	14.0	38	6	13.6	10.7
2	29.2	74	18	19.6	19.5
3	33.7	74	32	30.2	32.7
4	17.8	27	29	51.8	49.5
5	3.8	4	8	66.7	66.3
6	0.3	0	1	100.0	79.9
Low	15.2	42	6	12.5	10.5
Medium	29.2	74	18	19.6	21.7
High	55.6	105	70	40.0	39.5

Mortality risk groups were defined as low risk (1 point or fewer), medium risk (2 points) or high risk (more than 2 points)