

RESEARCH ARTICLE

# Endoparasites and vector-borne pathogens in dogs from Greek islands: Pathogen distribution and zoonotic implications

Anastasia Diakou<sup>1</sup>, Angela Di Cesare<sup>2</sup>, Simone Morelli<sup>2</sup>, Mariasole Colombo<sup>2</sup>, Lenaig Halos<sup>3</sup>, Giulia Simonato<sup>4</sup>, Androniki Tamvakis<sup>5</sup>, Frederic Beugnet<sup>3</sup>, Barbara Paoletti<sup>2</sup>, Donato Traversa<sup>2\*</sup>

**1** Laboratory of Parasitology and Parasitic Diseases, School of Veterinary Medicine, Faculty of Health Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece, **2** Faculty of Veterinary Medicine, University of Teramo, Località Piano D'Accio snc., Teramo, Italy, **3** Boehringer-Ingelheim Animal Health, Lyon, France, **4** Department of Animal Medicine, Production and Health, University of Padua, Viale dell'Università, Legnaro, Padua, Italy, **5** Laboratory of Ecology and System Dynamics, Department of Marine Sciences, University of the Aegean Mytilene, Lesvos, Greece

\* [dtraversa@unite.it](mailto:dtraversa@unite.it)



**OPEN ACCESS**

**Citation:** Diakou A, Di Cesare A, Morelli S, Colombo M, Halos L, Simonato G, et al. (2019) Endoparasites and vector-borne pathogens in dogs from Greek islands: Pathogen distribution and zoonotic implications. *PLoS Negl Trop Dis* 13(5): e0007003. <https://doi.org/10.1371/journal.pntd.0007003>

**Editor:** Hans-Peter Fuehrer, Vienna, AUSTRIA

**Received:** November 13, 2018

**Accepted:** March 28, 2019

**Published:** May 8, 2019

**Copyright:** © 2019 Diakou et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant results generated in the study are described in the article. The data underlying the results presented in the study are available from the corresponding author.

**Funding:** The execution of the present study has been supported by Merial S.A.S (now part of Boehringer-Ingelheim), 29 Av Tony Garnier, 69007, Lyon, France. The funders had no role in study design, data collection and analysis. The decision to publish has been taken into accordance with the funders, who have contributed in the preparation of

## Abstract

The present study investigated the presence of endo- and ecto-parasites, and vector-borne pathogens, in dogs from four islands of Greece. A total of 200 (123 owned and 77 sheltered) dogs were examined with different microscopic, serological and molecular methods.

Of the examined dogs, 130 (65%) were positive for one or more parasites and/or vector-borne pathogens. The most common zoonotic intestinal helminths recorded were *Ancylostomatidae* (12.5%) and *Toxocara canis* (3.5%). Ninety-three dogs (46.5%) seroreacted to *Rickettsia conorii*. Twenty-two (11%) of them were also PCR positive and 7 (3.5%) showed corpuscles suggestive of *Rickettsia* spp. on the blood smears. Nineteen dogs (9.5%) were seropositive for *Ehrlichia canis*, three of them being also PCR positive. Dogs positive for *Anaplasma phagocytophilum*-*Anaplasma platys* (1%), *Dirofilaria immitis* (0.5%) and *Babesia canis* (0.5%) were also found. Fleas and ticks were recorded in 53 (26.5%) and 50 (25%) dogs, respectively, and all specimens were identified as *Ctenocephalides felis felis* and *Rhipicephalus sanguineus sensu lato*. Binary multiple univariate Generalized Linear Models were used to investigate factors and clinical signs related to the recorded positivity, while the association of specific signs with the pathogens was evaluated using tests of independence. Knowledge of occurrence and impact of zoonotic parasites and vector-borne pathogens in dog populations is crucial to prevent the infection in animals and people, and to control the risk of spreading of these pathogens in endemic and non-endemic areas.

## Author summary

Both owned and sheltered dogs can harbor a variety of intestinal and extra-intestinal endoparasites, as well as vector-borne pathogens and ectoparasites, of zoonotic concern. Dog shelters and stray dogs are present in several touristic areas of Greece, including

the manuscript under a mere scientific point of view. There are no financial, personal, or professional interests that could be construed to have influenced the work. Lénaïg Halos and Frederic Beugnet are employers of Merial S.A.S (now part of Boehringer-Ingelheim), 29 Av Tony Garnier, 69007, Lyon, France

**Competing interests:** The corresponding author declares, on behalf of Anastasia Diakou, Angela Di Cesare, Simone Morelli, Mariasole Colombo, Lenaig Halos, Antonio Frangipane di Regalbono, Androniki Tamvakis, Frederic Beugnet, Barbara Paoletti and Donato Traversa that there are no financial, personal, or professional interests that could be construed to have influenced the work. Lénaïg Halos and Frederic Beugnet are employers of Merial S.A.S (now part of Boehringer-Ingelheim), 29 Av Tony Garnier, 69007, Lyon, France.

Sporades and Cyclades islands, where tourists often bring their pets with them, likely travelling from non-endemic to endemic areas. The present study has been carried out with the aim to evaluate the occurrence of the aforementioned pathogens. Data obtained showed that they are present in canine populations of Greece, with possibilities of infection for travelling dogs, which can also contribute to the spreading of zoonotic vector-borne diseases, introducing new pathogens in previously non-endemic areas. For these reasons, a constant monitoring of the epidemiological situation, improving control measures and correct diagnostic approaches are of primary importance for the prevention of canine and human infections, decreasing the spreading of potentially deadly pathogens.

## Introduction

Several parasitoses (e.g. internal helminthoses) and vector-borne diseases (VBDs) of veterinary importance represent a serious hazard for human health, particularly when transmission pressure and circulation of zoonotic infections are difficult to control. Because of a lifestyle that implies a low-grade of sanitary management, stray and free-roaming dogs are at high risk of becoming infected with a wide range of pathogens. Consequently, they act as a permanent source of infection for vectors, other animals and humans [1, 2].


Although dogs are efficient sentinels for investigating the occurrence and the epidemiological impact of zoonoses [2, 3] and a constant sanitary monitoring of canine populations is crucial, data on the simultaneous occurrence of endo/ecto- parasites and VBDs in several Mediterranean areas are still limited to specific narrow areas or to selected pathogens [4–7].

The number of stray dogs in most Greek Islands is low but many privately owned animals have a free-roaming lifestyle due to the local rural territory. Moreover, some islands have shelters for stray and abandoned animals, where prevention and treatment regimens are not regularly applied. At the same time, the number of dogs travelling to and from the insular Greece is increasing in proportion to tourism [8] with the realistic risk that these pets may acquire or introduce pathogens. Interestingly, a recent study conducted in Greece has shown that cats may be infected and/or exposed to a number of intestinal parasites and agents transmitted by arthropods, several of them of zoonotic importance and potentially shared between cats and dogs [9]. Therefore, the present study aimed to investigate the simultaneous occurrence of intestinal parasites, vector-borne pathogens and ectoparasites, in different canine populations, including stray and owned dogs in four regions of insular Greece.

## Methods

### Study areas and sampling

The study was conducted in four islands of Greece, located in the Aegean Sea (Fig 1). From north to south these islands were Skiathos (39.1704°N, 23.4506°E), Tinos (37.5968°N, 25.1216°E), Ios (36.7152°N, 25.3294°E) and Santorini (36.4199°N, 25.4316°E). The islands of the Aegean Sea have Mediterranean climate, with mild and moderately rainy winters, warm and sunny summers and frequent, often strong, winds. The vegetation is generally poor, especially in the islands of the Cyclades Archipelagos (Tinos, Ios and Santorini). The selection of these islands was based on their high tourism, prior collaboration with local vets and information they provided on availability of free-roaming and sheltered dogs. Authorizations to sample and examine the dogs were obtained case-by-case from local authorities, animal responsables/owners and animal rights organisations. Available data on sex, breed, living



**Fig 1.** A map of Greece showing the 4 study sites (islands Skiathos, Tinos, Ios and Santorini), of an investigation on the occurrence of zoonotic intestinal parasites and vector-borne pathogens in dogs.

<https://doi.org/10.1371/journal.pntd.0007003.g001>

conditions, age and presence or absence of clinical signs compatible with parasitoses were registered for each animal. Both faecal and blood samples were collected from dogs, that were also examined for fleas, ticks, ear mites and, in the presence of compatible skin lesions, body manges.

Ethics approval is not applicable as all activities carried out on dogs in the present study have been performed by routine diagnostics procedures performed by veterinarians working on each study site. Consent to examine and sample the animals have been obtained, case by case, from local authorities, animal owners and animal right organisations. Therefore, no official ethics permission or further authorization was required.

### Copromicroscopic examinations

Faecal samples were examined using standard zinc sulphate flotation and merthiolate iodine formaldehyde (MIF)—ether sedimentation methods [10, 11].

### Blood analysis

**Microscopic examinations.** Giemsa stained blood smears were performed to evaluate by light microscopy at 1000× magnification the presence of *Babesia* spp., *Rickettsia* spp., *Ehrlichia* spp. and *Anaplasma* spp. elements, based on morphology, size and cellular tropism [12, 13].

A modified Knott's technique was performed to detect and identify circulating microfilariae under light microscope (100×, 200× and 400× magnifications) [12]. If present, microfilariae were identified on the basis of differential morphometric (i.e. length and width) and morphological (i.e. anterior and posterior end) characteristics [14, 15].

**Serological examinations.** Sera obtained from blood samples were subjected to the following serological examinations, according to the manufacturers’ instructions.

- SNAP 4Dx Plus test (IDEXX Laboratories, Inc. USA) for the detection of *Dirofilaria immitis* circulating antigens and of antibodies against *Anaplasma phagocytophilum*/*Anaplasma platys*, *Ehrlichia canis* and *Borrelia burgdorferi*;
- *Leishmania* IC test (Agrolabo diagnostics, Italy) for the qualitative detection of anti-*Leishmania infantum* antibodies;
- indirect immunofluorescence antibody assay kits (IFAT) “Mega FLUO BABESIA *canis*” (Megacor Diagnostik GmbH, Austria) and “Mega FLUO RICKETTSIA *conorii*” (Megacor Diagnostik GmbH, Austria) for the detection of anti-*Babesia canis*-IgG and anti-*Rickettsia conorii*-IgG, with a screening dilution of 1:128 and 1:64 for *B. canis* and *R. conorii* respectively.

Positive and negative control sera were included in each test series.

**Molecular detection.** Genomic DNA was extracted from blood samples using a commercial kit (QIAamp DNA blood Mini kit—Qiagen GmbH, Hilden, Germany), and PCR positivity to Anaplasmataceae, *Rickettsia* spp., *Babesia* spp. and *Leishmania* spp. DNA was tested by different protocols (Table 1) [16, 17].

PCR products were individually sequenced directly in an automated sequencer. Sequences were determined in both directions, aligned using ClustalX software and analyzed with sequences available in GenBank using Nucleotide Basic Local Alignment Search Tool (BLASTN) [18].

### Ectoparasites

**Fleas and ticks.** The entire body of each dog was examined with an extra-fine flea comb. Fleas seen on the comb were removed using forceps and placed in a 2 ml microtube containing 70% ethanol for storage. If fleas were not present, any debris found on the comb was transferred to a piece of moist white paper, and animals were considered infected if the debris dissolved into a red color.

Ticks were individuated by thumb-counting, removed using forceps and placed in a 2 ml microtube containing 70% ethanol pending identification.

Collected fleas and ticks were identified by standard morphologic and morphometric keys [11, 19, 20].

**Table 1. Target genes amplified for the PCR analyses and primers used.**

Pathogen	Gene	Primers (5'-3')	Product size (bp)	Reference
<i>Anaplasma</i> spp./ <i>Ehrlichia</i> spp.	16S	EHR16SD: GGTACCYACAGAAGAAGTCC	345	[16]
	rRNA	EHR16SR: TAGCACTCATCGTTTACAGC		
<i>Babesia</i> spp.	18S rRNA	PIRO-A: AATACCCAATCCTGACACAGGG	400	[16]
		PIRO-B: TTAAATACGAATGCCCAAC		
<i>Rickettsia</i> spp.	Citrate synthase	Rsf877: GGGGGCCTGCTCACGGCGG	381	[17]
		Rsf1258: ATTGCAAAAAGTACAGTGAACA		
<i>Leishmania</i> spp.	Small subunit rRNA	Outer primers:	603	[16]
		R221: GGTCCTTTCCTGATTACG		
		R332: GGCCGGTAAAGGCCGAATAG		
		Inner primers:	358	
		R223: TCCCATCGCAACCTCGGTT		
		R333: AAAGCGGGCGGGTGCTG		

<https://doi.org/10.1371/journal.pntd.0007003.t001>

**Mites.** All dogs were subjected to an otoscopic examination by ear swabs to detect the presence of any sign (e.g. erythema, black/brown waxy discharge) caused by the ear mite *Otodectes cynotis*. Each ear swab was smeared with the addition of small quantity of mineral oil onto a glass slide, and examined under a microscope to identify mites by standard morphological keys [11, 21]. In the presence of compatible clinical signs, e.g. skin areas with loss of hair, itching, reddened rash, yellowish crusts, dogs were examined for manges as deemed appropriate. The collected material was examined under a microscope to identify mites by standard morphological keys [11, 21].

### Statistical analysis

The selected islands are small both in area (48 to 195 km<sup>2</sup>) and population (2,030 to 15,250 inhabitants), therefore the dog populations were assessed to be also rather small. Thus, the minimum number of dogs to be examined in each island was decided to be thirty, taking into consideration at the same time the regarding number of inhabitants (Santorini's and Tinos's samples are bigger in size than those of Skiathos and Ios) according to the principles of stratified sampling. Finally, the total sample size (n = 200) along with the subsample sizes of each island are compatible with the requirements of the statistical methods used in the present study.

Statistical analysis was performed to evaluate the association of five main factors (study site, age, sex, lifestyle and travel history) with infections and parasitoses detected in the examined dogs, especially with zoonotic potential. Binary multiple univariate Generalized Linear Models (GLM) were used to test the above mentioned factors with the presence of endoparasites (intestinal helminths and filariae), exposure/positivity for VBDs, positivity to zoonotic pathogens (*R. conorii*, *Anaplasma* spp., *L. infantum*, *Dirofilaria* spp., Ancylostomatidae and *T. canis*) and presence of ectoparasites [22]. In order to assess the GLM tests' importance, the ability in both detecting infected (sensitivity) and non-infected dogs (specificity) were determined. Furthermore, the determined odds ratio (OR) was used to measure the strength of association between the values of each factor to the presence of each infection. The corresponding confidence interval (CI) of OR was also presented to indicate the degree of uncertainty associated with the corresponding ratio. Thus, a wide CI implies that the number of tested animals is rather small, whereas if the CI includes number 1 then the calculated OR is not statistically significant [23]. The same analysis was applied to test whether the occurrence of four major clinical signs, i.e. dermatological signs (hair loss, exfoliative dermatitis, hyperkeratosis, crusts, ulceration, nodules), ocular manifestations (periocular alopecia, conjunctivitis, keratitis, uveitis, panophthalmitis) weight loss and pale mucosae was related to the detection of a VBD.

Finally, the association of gastrointestinal signs with intestinal parasitism and ectoparasitoses as evidence of exposure to VBDs was statistically tested using Fisher's exact test of independence [24]. The statistical analysis was implemented using the R package version 3.2.2 (R Development Core Team, 2006).

## Results

### Enrolment and geographical distribution

Overall 200 dogs were included in the survey, i.e. 66 from Santorini, 50 from Tinos, 43 from Ios and 41 from Skiathos islands. Table 2 reports age, sex and lifestyle of study dogs. In particular, 87 were privately owned with no travel history, 36 were pets that had travelled with their owners at least once (i.e. 32 across Greece, 2 in Bulgaria, 1 in Italy and 1 in different European countries), and 77 were sheltered animals. Both owned and sheltered dogs were highly distributed within different islands and age groups, with the exception of stray animals less than

Table 2. Distribution of the dogs examined in the different islands, per lifestyle, age group and sex.

Island	stray dogs				♂/♀	owned dogs				♂/♀	total				♂/♀
	age in years					age in years					age in years				
	<1	1-7	>7	total		<1	1-7	>7	total		<1	1-7	>7	total	
Santorini	13	19	2	34	19/15	6*	18*	8*	32	18/14	19	37	10	66	37/29
Skiathos	0	25	2	27	14/13	1	10*	3	14	8/6	1	35	5	41	22/19
Ios	0	2	0	2	0/2	5*	27*	9*	41	21/20	5	29	9	43	21/22
Tinos	0	7	7	14	8/6	3	24*	9*	36	17/19	3	31	16	50	25/25
Total	13	53	11	77	41/36	15*	79*	29*	123	64/59	28	132	40	200	105/95

\*group that includes dogs that have travelled in the country or abroad

<https://doi.org/10.1371/journal.pntd.0007003.t002>

1-year-old that were found only in Santorini (n = 13). Although the number of stray dogs examined in Ios seems to be rather small (n = 2), it can be considered as a significant part of the corresponding small population as Ios is a small barren island, with high elevation and only 2,030 inhabitants. Although the dataset can be considered sufficiently large (n = 200) so that the heuristic rule (i.e. a few of the expected cells counts less than five is not violated) provides safe statistical results [25], a high degree of uncertainty regarding prevalences in stray dogs of Ios (n = 2) is expected.

### Exposure to parasites, vector borne diseases and prevalence of zoonotic agents

Overall 130 dogs (65%) were positive for at least one intestinal parasite or VBD by any of the tests applied. In particular, 96 (48%) showed a monospecific infection and 34 (17%) scored positive for mixed infections by endoparasites and VBD. Two animals (1%) were infected by more than 1 endoparasite, 16 (8%) by more than one VBD and 16 (8%) were infected by both intestinal parasites and VBDs. Various zoonotic agents were found in 128 (64%) dogs. Fleas and ticks were recorded in 53 (26.5%) and 50 (25%) dogs, respectively.

**Endoparasites.** Overall thirty-six (18%) dogs were positive for at least one endoparasite at copromicroscopic examinations. Ancylostomatidae (12.5%, 25/200) and *Toxocara canis* (i.e. 3.5%, 7/200) were the most common zoonotic helminths. Non-zoonotic infections by *Trichuris vulpis* (3.5%) and coccidia (1%) were also found (Table 3). Eggs of the zoonotic rat tapeworm *Hymenolepis diminuta* were detected in the faeces of one dog, while eggs of the trematode *Dicrocoelium dendriticum* were found in the faeces of another animal.

The relative prevalence of intestinal parasites in the study animals was related to the island of residence, the age and the lifestyle of the animal (Table 4). In particular, the odd of intestinal parasites occurrence was 5.49, 11.43 and 23.86 times higher in animals residing in Santorini, compared to those that reside in Tinos, Ios and Skiathos, respectively. Animals ageing 1 to 7 years were found to be 4 times (i.e. 1/0.25) more likely to be infected compared to young animals up to 1 year of age. Furthermore, a higher risk of intestinal parasites prevalence was found in stray dogs (odds ratio = 11.37) compared to owned dogs. Some of the computed CIs of statistical significant OR (e.g. CI related to the intestinal parasites between Santorini and Ios or the CI concerning dogs aged up to 1 yr and dogs aged 1 to 7 yrs) are rather wide, implying high uncertainty in the computed value of OR originated by the small number of sampled dogs in some levels of the corresponding factors (Table 2). However, the good values of both sensitivity and specificity show that the three significant factors (island, age and lifestyle) affect the occurrence of the intestinal parasites. Nonetheless, the overall occurrence of intestinal parasites was not related to the presence of clinical signs (Fisher's exact test,  $p = 0.3283$ ).

Table 3. Prevalence of intestinal parasites (as determined by standard copromicroscopic examination) in the study dog population.

Pathogen	n. positive (%) owned dogs	n. positive (%) sheltered dogs	n. positive (%) total
Nematodes			
Ancylostomatidae	3 (2.4)	22 (28.6)	25 (12.5)
<i>Toxocara canis</i>	0 (0)	7 (9.1)	7 (3.5)
<i>Trichuris vulpis</i>	4 (3.2)	3 (3.9)	7 (3.5)
Protozoan			
<i>Cystoisospora</i> spp.	0 (0)	2 (2.6)	2 (1)
Trematodes			
<i>Dicrocoelium dendriticum</i>	0 (0)	1 (1.3)	1 (0.5)
Cestodes			
<i>Hymenolepis diminuta</i>	0 (0)	1 (1.3)	1 (0.5)
<b>Total number of positive dogs*</b>	<b>7 (5.7)</b>	<b>29 (37.7)</b>	<b>36 (18)</b>

\*dogs with mixed infections included.

<https://doi.org/10.1371/journal.pntd.0007003.t003>

**Ectoparasites.** Fleas were recorded in 26.5% of the dogs (53/200), of which 24.2% (16/66) from Santorini, 58% (29/50) from Tinos, 9.3% (4/43) from Ios and 9.8% (4/41) from Skiathos. All collected specimens were identified as *Ctenocephalides felis felis*. Ticks were present in 25% of the dogs (50/200), namely in 40.9% (27/66), 38% (19/50) and 9.3% (4/43) from Santorini, Tinos and Ios respectively. All ticks were *Rhipicephalus sanguineus sensu lato*. Mites were not found in any of the investigated dogs.

Mixed infections by ticks and transmitted diseases were recorded in 28 (14%) dogs while no dogs showed mixed infection by fleas and transmitted pathogens.

The occurrence of fleas and ticks was statistically associated with study site, lifestyle and travelling. Specifically, a higher risk of ectoparasitosis was recorded in Santorini against Ios (odds ratio = 4.51) and Skiathos (odds ratio = 19.28), while the same risk was found in Tinos. Stray dogs increased the odds of ectoparasitosis (odds ratio = 3.17), while travelling decreased the odds (odds ratio = 0.33) (Table 4). Moreover, the high values of sensitivity and specificity confirm the effect of the above factors to the ectoparasites occurrence.

### Blood examination (Table 5)

*Rickettsia conorii*: Overall, 93 dogs (46.5%) seroreacted to *R. conorii*. Twenty-two (11%) were also PCR positive and, at the blood smear examination, the samples of seven (3.5%) of them were positive for corpuscles suggestive of *Rickettsia* spp. in the monocytes. Sequences from the 22 PCR products showed 99% identity with *R. conorii* GenBank Accession number U59730.1. All dogs negative by serological analyses were also negative for *Rickettsia* spp. upon PCRs. Of the 93 *R. conorii*-seropositive dogs, 19 had a tick infestation and 6 scored also PCR-positive for *R. conorii*. The prevalence of *R. conorii* in the animals studied was found statistically related to the presence of lymphadenopathy (Fisher's exact test,  $p = 0.0177$ ) whilst no any relation was detected with signs like weight loss, pale mucous membrane, gastrointestinal disorders, conjunctivitis or dermatological manifestations (Fisher's exact test,  $p \geq 0.05$ ).

Anaplasmataceae: Nineteen dogs (9.5%) were seropositive for *E. canis*, three of them being also positive at the PCR (100% homology with Genbank Accession number LC018188.1), while the others were PCR-negative. Two dogs (1%) seroreacted for *A. phagocytophilum/A. platys*. One of them showed corpuscles suggestive of *Anaplasma* spp. in the neutrophils at the Giemsa staining and scored PCR-positive for *A. phagocytophilum* (homology of 100% with Genbank Accession number KY114936.1), while the other was microscopically and PCR-negative.

**Table 4. Statistical analysis evaluating various factors (i.e. island where the dogs lived, age, sex, lifestyle and traveling history) in relation to the different infections detected in the study animals.**

Variable	Positive for intestinal parasites	Positive for VBDs	Positive for filariae	Positive for ectoparasites	Zoonotic infections
	Odds ratio	Odds ratio	Odds ratio	Odds ratio	Odds ratio
	(95% CI)	(95% CI)	(95% CI)	(95% CI)	(95% CI)
	GLM P-value	GLM P-value	GLM P-value	GLM P-value	GLM P-value
<b>Island</b>					
Santorini vs Tinos	5.49 (1.60–18.83) <b>0.007**</b>	0.30 (0.13–0.67) <b>0.005**</b>	N/A	0.58 (0.24–1.37) <i>0.211</i>	0.66 (0.28–1.59) <i>0.357</i>
Santorini vs Ios	11.43 (1.31–99.56) <b>0.027*</b>	0.76 (0.31–1.85) 0.548	N/A	4.51 (1.52–13.36) <b>0.007**</b>	1.73 (0.70–4.30) 0.233
Santorini vs Skiathos	23.86 (5.51–103.33) <b>&lt;0.001***</b>	0.24 (0.10–0.60) <b>0.002**</b>	N/A	19.28 (5.51–67.41) (5.51–67.41)	0.77 (0.29–2.07) 0.613
<b>Age</b>					
Up to 1 yr vs 1 to 7 yrs	0.25 (0.06–0.97) <b>0.046*</b>	0.36 (0.14–0.96) <b>0.040*</b>	N/A	0.67 (0.24–1.89) <i>0.458</i>	0.26 (0.10–0.69) <b>0.007**</b>
Up to 1 yr vs more than 7 yrs	0.26 (0.05–1.35) <i>0.108</i>	0.54 (0.18–1.64) <i>0.277</i>	N/A	0.59 (0.17–1.97) <i>0.392</i>	0.51 (0.17–1.54) <i>0.229</i>
<b>Sex</b>					
Male vs Female	1.36 (0.54–3.45) <i>0.518</i>	2.33 (0.24–4.37) <b>0.008**</b>	0.61 (0.07–5.45) <i>0.656</i>	0.72 (0.36–1.45) <i>0.356</i>	2.13 (1.11–4.10) <b>0.023*</b>
<b>Lifestyle</b>					
Stray vs Owned	11.37 (3.73–34.72) <b>&lt;0.001***</b>	1.15 (0.55–2.40) <i>0.707</i>	0.29 (0.03–2.98) <i>0.300</i>	3.17 (1.37–7.34) <b>0.007**</b>	3.00 (1.37–6.60) <b>0.006**</b>
<b>Travel</b>					
Yes vs No	0.49 (0.05–4.48) <i>0.526</i>	1.34 (0.58–3.11) <i>0.498</i>	N/A	0.33 (0.11–0.97) <b>0.043*</b>	1.10 (0.48–2.53) <i>0.819</i>
Sensitivity	0.67	0.68	0.02	0.71	0.73
Specificity	0.90	0.66	0.00	0.83	0.64

\*Statistical significant result at the 0.05 level

\*\*Statistical significant result at the 0.01 level

\*\*\* Statistical significant result at the 0.001 level. N/A: Not applicable

<https://doi.org/10.1371/journal.pntd.0007003.t004>

Ticks were present in 15/19 dogs seropositive for *E. canis* at the SNAP 4Dx Plus and 3 of them were among those that scored positive at the PCR.

*Babesia* spp.: One dog (0.5%) was serologically and PCR positive for *B. canis* showing homology of 99% with GenBank Accession number: KJ696714.1, while all other dogs were negative.

*Leishmania infantum*: A total of 13 (6.5%) samples seroreacted for *L. infantum* and all of them were also PCR-sequencing positive, with 99–100% identity with *L. infantum* GenBank Accession number HM807524.1. All seronegative dogs were also negative upon PCR.

**Table 5. Observed prevalence of vector borne pathogens (as determined by PCR, serology and blood microscopy) in the study dog population.**

Pathogen	PCR	Serological detection	Smear or other direct parasitological examination	Positive by at least one test
	n. positive (%)	n. positive (%)	n. positive (%)	n. positive (%)
Anaplasmataceae				
<i>Ehrlichia canis</i>	3 (1.5)	19 (9.5)	0	19 (9.5)
<i>Anaplasma</i> spp.	1 (0.5)	2 (1)	1 (0.5)	2 (1)
	<i>A. ph.</i>	<i>A. ph./A. pl</i>	<i>Anaplasma</i> spp.	<i>A. ph./A. pl.</i>
<i>Leishmania infantum</i>	13 (6.5)	13 (6.5)	ND	13 (6.5)
<i>Rickettsia conorii</i>	22 (11)	93 (46.5)	7 (3.5)	93 (46.5)
	<i>R. c.</i>	<i>R. c.</i>	<i>Rickettsia</i> spp.	<i>R. c.</i>
<i>Babesia canis</i>	1 (0.5)	1 (0.5)	0 (0)	1 (0.5)
<i>Dirofilaria</i> spp.			1 (0.5)	1 (0.5)
	ND	1 (0.5)	<i>D. i.</i>	<i>D. i.</i>
		<i>D. i.</i>	3 (1.5)	3 (1.5)
			<i>D. r.</i>	<i>D. r.</i>
<i>Borrelia burgdorferi</i>	ND	0 (0)	ND	0 (0)

*A. ph.*: *Anaplasma phagocytophilum*; *A. pl.*: *Anaplasma platys*; *L. i.*: *Leishmania infantum*; *R. c.*: *Rickettsia conorii*; *B. c.*: *Babesia canis*; *D. i.*: *Dirofilaria immitis*; *D. r.*: *Dirofilaria repens*. ND: Not Done.

<https://doi.org/10.1371/journal.pntd.0007003.t005>

*Dirofilaria* spp.: One (0.5%) of the examined dogs was positive for *D. immitis*, at both Knott’s test and serology. Moreover, in three dogs (1.5%) microfilariae of *Dirofilaria repens* were found in the Knott’s test.

*Borrelia burgdorferi*: None of the 200 sera examined for *B. burgdorferi* was found positive.

**Risk factor for exposure to parasites and VBDs.** The percentage of dogs positive for at least one intestinal parasites and/or VBDs was similar between travelling (i.e. 58.3%) and non-travelling dogs (66.5%) while the highest number of infected animals was found in sheltered animals (83.1%).

The exposure to at least one VBD was statistically associated to the study site, the sex and the age. In particular, animals residing in Skiathos were 4.17 (i.e. 1/0.24) and 3.17 (i.e. 0.76/0.24) times more likely to be infected than those in Santorini and Ios, respectively, while no difference was found compared to dogs from Tinos. Furthermore, VBD prevalence in the study animals was associated with sex and age, as males increased the odds (odds ratio = 2.33) and dogs ageing up to 1 year decreased the odds (odds ratio = 0.36) against those ageing 1–7 years. These findings are confirmed by the values of both sensitivity and specificity.

The diagnosis of VBDs was associated with the presence of various skin lesions (GLM,  $p < 0.05$ ) in infected animals. In fact, animals with skin lesion were 3.2 times more likely to be seropositive compared to those with no dermatological manifestations. Similarly, weight loss was associated with 5.43-fold increased odds (Table 6). Nonetheless, positivity to a VBD was not related to the presence of ectoparasites (Fisher’s exact test,  $p = 0.6623$ ), and clinical signs (i.e. lymphadenopathy, pale mucous membrane or conjunctivitis) were not statistically significant factors (GLM,  $p > 0.05$ ).

### Geographic distribution of parasites

In the dog population studied herein, animals from Santorini showed the highest prevalence of parasite infection (65.2%) and the highest percentage of Ancylostomatidae occurrence (27.3%) (Fisher’s exact test,  $p < 0.001$ ), while Skiathos had the highest rate of dogs that seroreacted to *R. conorii* (73.2%) (Fisher’s exact test,  $p < 0.001$ ) and of dogs infected by *D. immitis* (2.4%) and *D. repens* (7.3%) (Fisher’s exact test,  $p = 0.008$ ). Skiathos also showed the highest percentage of positivity to *L. infantum* (17.1%).

Table 6. Statistical analysis to evaluate the presence of clinical signs in relation to the different infections detected in the examined dogs.

Variable	Positive for VBD		Positive for <i>L. infantum</i>	
	Odds ratio (95% CI)	GLM P-Value	Odds ratio (95% CI)	GLM P-Value
<i>Lymphadenopathy</i>	0.66	0.335	0.82	0.815
Yes vs No	(0.28–1.54)		(0.16–4.30)	
<i>Dermatological manifestations</i>	3.20	<b>0.027*</b>	2.62	0.205
Yes vs No	(1.14–8.97)		(0.59–11.59)	
<i>Weight loss</i>	5.43	<b>0.033*</b>	2.87	0.210
Yes vs No	(1.15–25.65)		(0.55–14.90)	
<i>Pale mucous membrane</i>	0.48	0.242	0.62	0.679
Yes vs No	(0.14–1.65)		(0.06–6.06)	
<i>Conjunctivitis/Ocular manifestations</i>	4.86	0.154	N/A	0.992
Yes vs No	(0.55–42.54)			

N/A: Not applicable

<https://doi.org/10.1371/journal.pntd.0007003.t006>

### Zoonotic risk

The risk to carry zoonotic pathogens (i.e. *R. conorii*, *Anaplasma* spp., *L. infantum*, *Dirofilaria* spp., Ancylostomatidae and *T. canis*) was associated to animal age, as dogs ageing 1–7 years increased the odds of infections by 3.85-fold (i.e. 1/0.26) in comparison to younger animals. Furthermore, the sex and lifestyle were also found statistically related with zoonotic infections with male and stray dogs being 2.13 and 3 times more likely affected than female and owned dogs, respectively. These results are consistent with the values of sensitivity and specificity.

### Discussion

In the present study 65% of the dogs showed exposure to or were carrying pathogens, and 64% harboured a pathogen with a zoonotic potential. Among them, canine geohelminths have an important health impact for both animals and humans. Thus, dogs here found infected by *T. canis* or hookworms represent a potential health risk especially because free-roaming animals are a source of environmental contamination [26]. Human infection by *T. canis* may lead either to subclinical infections or to different larva migrans syndromes i.e. visceral, ocular and neural, which may have serious clinical manifestations [27, 28]. *Ancylostoma caninum* larvae may penetrate the human skin and cause follicular, papular/pustule and ephemeral lesions, muscular damages and, seldom, eosinophilic enteritis. The risk of human infection is enhanced by walking barefoot or lying on grass and soil in contaminated areas [28].

Eggs of the tapeworm *H. diminuta* were found in one dog. Although the main hosts of this parasite are rodents [29, 30], in rare cases dogs and other mammals, including humans, may also be infected. Similarly, eggs of *D. dentriticum* a trematode commonly found in the liver of ruminants, were found in one dog. While in some cases this parasite can infect other mammals, including dogs and humans [31, 32], it is plausible that pseudoparasitism is the cause of this finding [33].

The high rate of exposure of the here studied dogs to several pathogens transmitted by arthropods, is of importance because some VBDs are shared between companion animals and people. Moreover, to the best of the authors' knowledge, this is the first report of seroprevalence and molecular detection of *Babesia* spp. and *R. conorii* in dogs in Greece and of *E. canis* and *Anaplasma* spp. in dogs from the herein study areas.

The high seroprevalence for *R. conorii*, i.e. the main aetiological agent of the Mediterranean spotted fever in Europe [34], suggests a frequent exposure and/or persistent low-grade

infections. This is not surprising if one considers the geographical location of the study sites, i.e. regions in Southern Europe with favorable environments for ticks and local circulation of transmitted pathogens. It is worth noting that the DNA of this bacterium was found in 11% of examined dogs, thus further corroborating recent findings that have indicate dogs as important reservoirs of this pathogen and source of infection for ticks [35–37]. The role of dogs as reservoirs is further supported by the absence of clinical signs or by the presence of mild aspecific alterations, as in the present study (e.g. lymphadenopathy). Under an epizootiological standpoint it is remarkable that all ticks collected from positive animals were *R. sanguineus*, i.e. the main vector of *R. conorii* [34]. It is also interesting that a past survey carried out in the island of Crete showed a human seroprevalence rate of 7.6% for *R. conorii* [38]. Although no ticks from Crete were PCR-positive for *R. conorii* [38], the bacterium was isolated from *R. sanguineus* and humans in other areas of Greece with high seroprevalence in people [39, 40]. More importantly, several cases of human Mediterranean spotted fever have been reported in Greece [41, 42].

*Ehrlichia canis* causes the canine monocytic ehrlichiosis, a severe and potentially life-threatening illness in dogs transmitted in Europe by *R. sanguineus* [43–45]. Although an *E. canis*-like organism has been described in humans in Venezuela [46], to date this VBD is not considered zoonotic. The infection rate by *E. canis*, much lower than that recorded for *R. conorii*, could appear unexpected, but it is similar with that recorded in a recent study from southern Italy [47]. It should be noted that different lineages of *R. sanguineus* have a low capacity to transmit *E. canis* [48], thus explaining why in some cases the prevalence of *E. canis* infection could result lower than expected [49, 50].

It could be hypothesized that some *R. sanguineus* lineages may have opposite ability to transmit either *R. conorii* or *Ehrlichia/Anaplasma*. Further studies to investigate this issue are thus warranted.

*Anaplasma phagocytophilum* was detected in one dog. It is an emerging vector-borne pathogen transmitted by *Ixodes ricinus* ticks [51]. This tick species was not found on the dogs examined in the present study. However, it is widespread in continental Greece [50, 52, 53] and further investigations to evaluate its presence in insular Greece could be thus useful. The absence or limited occurrence of *A. platys*, the agent of infectious cyclic thrombocytopenia, is more surprising as this bacterium is probably transmitted by *R. sanguineus* ticks and common in the Mediterranean regions [44, 54]. Again, this epidemiological feature could be due to the presence of different lineages of the brown dog tick.

Although the most common canine haemoprotozoan in Mediterranean countries is *Babesia vogeli*, *B. canis* is also enzootic in Europe [3], including countries neighbouring to Greece and the Balkan Peninsula [55, 56]. The presence of *B. canis* in insular regions of Greece is an unexpected result because its primary vector, *Dermacentor reticulatus*, generally lives in cool and wet climates. As *D. reticulatus* can also be found in warm and temperate areas [57], its sporadic presence in Greece, especially in continental regions, needs to be taken into account. The owner of the dog declared that the dog had travelled in continental Greece but further details were not provided. Thus, the origin of this result could be likely due to an infection acquired in other areas of the country, although further information was not available.

The presence of several *L. infantum*-positive dogs in all study sites is explained by its wide distribution in Southern Europe [58], including Greece where mean seropositivity is around 20% [5, 6]. This high infection pressure is shown also by the recent records of infection in cats from Crete and Athens [9] and by the more than 300 autochthonous human cases reported in 2005–2010 according the Hellenic Center for Disease Control and Prevention [6]. The zoonotic potential of *L. infantum* in humans is of great importance, as the disease may be severe and presents with visceral, cutaneous and mucocutaneous signs [59].

Dogs living in Skiathos proved to be at risk of dirofilariosis. Though *Dirofilaria* spp. have been here diagnosed with a low prevalence, their ability in causing disease in humans should be taken into account. The number of human cases of *D. repens* infection in Europe is currently a public health concern and the lack of awareness regarding diagnosis and control in microfilariaemic dogs could lead to lack of vigilance and underestimation for this parasite [60]. *Dirofilaria* infections were here found only in Skiathos (where endemic infections by *D. immitis* are absent) in a single dog that was imported from an area of Central Greece where the prevalence of canine dirofilariosis is about 7% [61]. This confirms that undiagnosed and untreated microfilariaemic dogs are an important source of infection for mosquitoes and may also introduce these pathogens in other regions.

The association of the study sites with given infections may be due to the different island characteristics. Skiathos, where dogs resulted more likely infected with VBDs and ectoparasites than in other sites, is a part of the Vories Sporades, an island formation with a rich vegetation that fosters the maintenance of ectoparasites and vectors more efficiently. Indeed, the dry environment of Santorini, Ios and Tinos, belonging to different island formations (i.e. Cyclades) is characterized by poor vegetation.

The higher occurrence of intestinal parasites in Santorini rather than in the other islands is difficult to explain. These results may harbour high uncertainty due to the small number of stray dogs sampled mainly in Ios and also in Tinos. Although these findings may be attributed to chance, it could also be possible that differences in regular epizootiological vigilance and appropriate veterinary care in terms of prevention, diagnosis and treatment have a certain role, although further investigations are warranted to clarify these issues.

The higher occurrence of intestinal nematodes in dogs ageing 1–7 years should be interpreted with caution given the lack of precise information provided in some cases by the owners about anthelmintic treatments. Owners could be less willing to engage in preventative or therapeutic practices for dogs ageing more than 1 year as they are erroneously considered at less risk of intestinal parasitoses. These data further indicate that a high level of vigilance comprising routine faecal examinations and appropriate anthelmintic treatments are still indicated in adult dogs and should be encouraged among owners. It is worth mentioning that in a recent similar study carried out in 4 localities of Greece, free-roaming cats were found to be exposed to an important number of pathogens (endoparasites and VBP), some of which zoonotic and also capable to infect/infest dogs (e.g. *Dirofilaria* spp., *Leishmania* spp., *Rickettsia* spp.) [9]. These data underline the importance of focused control measures towards the prevention of such parasitic and vector borne pathogens in both cat and dog populations.

In conclusion, it is here shown that regular investigations should be encouraged in areas where data on parasite occurrence is still limited, and that prompt diagnosis and infection control are a priority. This is especially relevant for Mediterranean countries where epizootiological and biological conditions favour the occurrence and circulation of several parasitoses and VBDs in animal populations due to goods and animal trade, dog adoptions and holiday trips of people [37, 44, 62–64]. As visitors and tourists often bring their pets during holidays, introduction of parasites from endemic to free areas can occur. At the same time there is the risk that these dogs acquire pathogens in a new, enzootic environment and introduce them in free areas when they go home. In this view, timely use of dewormers (e.g. macrocyclic lactones) secures treatment of zoonotic geohelminthoses and at the same time provide efficacious prevention for spreading (e.g. *D. immitis*) or largely distributed (e.g. roundworms) parasites. Also, the appropriate use of acaricides, insecticides and repellents are reliable ways to improve dog health and to control environmental infection by vectors, preventing the spread of potentially deadly diseases.

## Acknowledgments

The authors thank Carmine Merola and Raffaella Iorio for the participation in this study. Special acknowledgments are due to the veterinarians in the Islands of the study: Maria Danezi, Kalliopi Tzagari and Manina Vlastaridou as well as to the Greek Action of Volunteer Veterinarians (GAVV) for their valuable help and excellent collaboration.

## Author Contributions

**Conceptualization:** Anastasia Diakou, Lenaig Halos, Donato Traversa.

**Data curation:** Anastasia Diakou, Angela Di Cesare, Giulia Simonato, Androniki Tamvakis, Donato Traversa.

**Formal analysis:** Anastasia Diakou, Angela Di Cesare, Giulia Simonato, Barbara Paoletti, Donato Traversa.

**Funding acquisition:** Donato Traversa.

**Investigation:** Anastasia Diakou, Angela Di Cesare, Simone Morelli, Mariasole Colombo.

**Methodology:** Anastasia Diakou, Angela Di Cesare, Simone Morelli, Mariasole Colombo, Barbara Paoletti.

**Project administration:** Donato Traversa.

**Resources:** Lenaig Halos, Frederic Beugnet.

**Supervision:** Angela Di Cesare, Donato Traversa.

**Validation:** Mariasole Colombo.

**Writing – original draft:** Anastasia Diakou, Angela Di Cesare, Lenaig Halos, Donato Traversa.

**Writing – review & editing:** Anastasia Diakou, Angela Di Cesare, Simone Morelli, Lenaig Halos, Giulia Simonato, Androniki Tamvakis, Frederic Beugnet, Barbara Paoletti, Donato Traversa.

## References

1. Deplazes P, van Knapen F, Schweiger A, Overgaaauw PA. Role of pet dogs and cats in the transmission of helminthic zoonoses in Europe, with a focus on echinococcosis and toxocarosis. *Vet Parasitol.* 2011; 182: 41–53. <https://doi.org/10.1016/j.vetpar.2011.07.014> PMID: 21813243
2. Baneth G, Thamsborg SM, Otranto D, Guillot J, Blaga R, Deplazes P, et al. Major parasitic zoonoses associated with dogs and cats in Europe. *J Comp Pathol.* 2016; 155(1 Suppl 1): S54–74. <https://doi.org/10.1016/j.jcpa.2015.10.179> PMID: 26687277
3. Solano-Gallego L, Sainz Á, Roura X, Estrada-Peña A, Miró G. A review of canine babesiosis: the European perspective. *Parasit Vectors.* 2016; 9: 336. <https://doi.org/10.1186/s13071-016-1596-0> PMID: 27289223
4. Tennant KV, Barker EN, Polizopoulou Z, Helps CR, Tasker S. Real-time quantitative polymerase chain reaction detection of haemoplasmas in healthy and unhealthy dogs from Central Macedonia, Greece. *J Small Anim Pract.* 2011; 52: 645–649. <https://doi.org/10.1111/j.1748-5827.2011.01126.x> PMID: 22017540
5. Athanasiou LV, Kontos VI, Saridomichelakis MN, Rallis TS, Diakou A. A cross-sectional sero-epidemiological study of canine leishmaniasis in Greek mainland. *Acta Trop.* 2012; 122: 291–295. <https://doi.org/10.1016/j.actatropica.2012.02.003> PMID: 22366671
6. Ntais P, Sifaki-Pistola D, Christodoulou V, Messaritakis I, Pratloug F, Poupalos G, et al. Leishmaniases in Greece. *Am J Trop Med Hyg.* 2013; 89: 906–915. <https://doi.org/10.4269/ajtmh.13-0070> PMID: 24062479

7. Kostopoulou D, Claerebout E, Arvanitis D, Ligda P, Voutzourakis N, Casaert S, et al. Abundance, zoonotic potential and risk factors of intestinal parasitism amongst dog and cat populations: The scenario of crete, Greece. *Parasit Vectors*. 2017; 10: 43–55. <https://doi.org/10.1186/s13071-017-1989-8> PMID: [28122583](https://pubmed.ncbi.nlm.nih.gov/28122583/)
8. Chatzidakis A. Development analysis of tourist movement. Greek Tourism Organisation, The National Printing House. Athens, Greece; 2011.
9. Diakou A, Di Cesare A, Accettura PM, Barros L, Iorio R, Paoletti B, et al. Intestinal parasites and vector-borne pathogens in stray and free-roaming cats living in continental and insular Greece. *PLoS Negl Trop Dis*. 2017; 11: e0005335. <https://doi.org/10.1371/journal.pntd.0005335> PMID: [28141857](https://pubmed.ncbi.nlm.nih.gov/28141857/)
10. MAFF Ministry of Agriculture, Fisheries and Food. Manual of Veterinary Parasitological Laboratory Techniques; 1986.
11. Taylor MA, Coop RL, Wall RL. *Veterinary Parasitology*. 3rd ed. Blackwell, Oxford; 2007.
12. Sloss MW, Kemp RL, Zajac AM. *Veterinary Clinical Parasitology*. 6th ed. Iowa State University Press, USA; 1994.
13. Harvey JW, Simpson CF, Gaskin JM. Cyclic thrombocytopenia induced by a *Rickettsia*-like agent in dogs. *J Infect Dis*. 1978; 137: 182–188. PMID: [627738](https://pubmed.ncbi.nlm.nih.gov/627738/)
14. Magnis J, Lorentz S, Guardone L, Grimm F, Magi M, Naucke TJ, et al. Morphometric analyses of canine blood microfilariae isolated by the Knott's test enables *Dirofilaria immitis* and *D. repens* species-specific and *Acanthocheilonema* (syn. *Dipetalonema*) genus-specific diagnosis. *Parasit Vectors*. 2013; 6: 48. <https://doi.org/10.1186/1756-3305-6-48> PMID: [23442771](https://pubmed.ncbi.nlm.nih.gov/23442771/)
15. Euzeby J. Diagnostic experimental des helminthoses animales. Travaux pratiques d'helminthologie veterinaire, Livre 1. Paris, France: Informations Techniques des Services Veterinaires, Ministry of Agriculture; 1981.
16. Maia C, Ramos C, Coimbra M, Bastos F, Martins A, Pinto P, et al. Bacterial and protozoal agents of feline vector-borne diseases in domestic and stray cats from southern Portugal. *Parasit Vectors*. 2014a; 7: 115–123.
17. Maia C, Ferreira A, Nunes M, Vieira ML, Campino L, Cardoso L. Molecular detection of bacterial and parasitic pathogens in hard ticks from Portugal. *Ticks Tick Borne Dis*. 2014b; 5: 409–414.
18. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*. 1997; 25: 3389–3402. <https://doi.org/10.1093/nar/25.17.3389> PMID: [9254694](https://pubmed.ncbi.nlm.nih.gov/9254694/)
19. Manilla G, Iori A. Illustrated key to the ticks of Italy. I. Larval stages of the species of the Ixodinae subfamily (Acari, Ixodoidea, Ixodidae). *Parassitologia*. 1992. 34: 83–95. PMID: [1339981](https://pubmed.ncbi.nlm.nih.gov/1339981/)
20. Linardi PM, Santos JL. *Ctenocephalides felis felis* vs. *Ctenocephalides canis* (Siphonaptera: Pulicidae): some issues in correctly identify these species. *Rev Bras Parasitol Vet*. 2012; 21:345–354. PMID: [23295817](https://pubmed.ncbi.nlm.nih.gov/23295817/)
21. Varma MGR. Ticks and mites. In: Lane RP, Crossey RW, editors. *Medical Insects and Arachnids*. Chapman and Hall, London; 1993.
22. Vittinghoff E, Glidden DV, Shiboski SC, McCulloch CE. *Regression methods in biostatistics: Linear, logistic, survival and repeated measures models*. 2nd ed. Springer, New York; 2012.
23. Bland M. *An Introduction to Medical Statistics*, 3rd ed., Oxford University Press; 2000.
24. Zar JH. *Biostatistical analysis*. 4th ed. New Jersey: Prentice-Hall; 1998.
25. McCullagh P, Nelder JA. *Generalized Linear Models*. 2nd ed. Chapman and Hall, London; 1989.
26. Traversa D. Pet roundworms and hookworms: a continuing need for global worming. *Parasit Vectors*. 2012; 5: 91. <https://doi.org/10.1186/1756-3305-5-91> PMID: [22574783](https://pubmed.ncbi.nlm.nih.gov/22574783/)
27. Despommier D. Toxocarasis: clinical aspects, epidemiology, medical ecology, and molecular aspects. *Clin Microbiol Rev*. 2003; 16: 265–272. <https://doi.org/10.1128/CMR.16.2.265-272.2003> PMID: [12692098](https://pubmed.ncbi.nlm.nih.gov/12692098/)
28. Lee AC, Schantz PM, Kazacos KR, Montgomery SP, Bowman DD. Epidemiologic and zoonotic aspects of ascarid infections in dogs and cats. *Trends Parasitol*. 2010; 26: 155–161. <https://doi.org/10.1016/j.pt.2010.01.002> PMID: [20172762](https://pubmed.ncbi.nlm.nih.gov/20172762/)
29. Guardone L, Macchioni F, Torracca B, Gabrielli S, Magi M. *Hymenolepis diminuta* (rat tapeworm) infection in a dog in Liguria, northwest Italy. *Parassitologia*. 2010; 52: 244.
30. Gupta P, Gupta P, Bhakri BK, Kaistha N, Omar BJ. *Hymenolepis diminuta* infection in a school going child: first case report from Uttarakhand. *J Clin Diagn Res*. 2016; 10: 4–5.
31. Cabeza-Barrera I, Cabezas-Fernández T, Salas Coronas J, Vázquez Villegas J, Cobo F. *Dicrocoelium dendriticum*: an emerging spurious infection in a geographic area with a high level of immigration. *Ann*

- Trop Med Parasitol. 2011; 105: 403–406. <https://doi.org/10.1179/1364859411Y.0000000029> PMID: 21929882
32. Azmoudeh-Ardalan F, Soleimani V, Jahanbin B. *Dicrocoelium dentriticum* in explanted liver: report of an unusual finding. *Exp Clin Transplant*. 2017; 15(Suppl 1): 178–181. <https://doi.org/10.6002/ect.mesot2016.P62> PMID: 28260462
  33. Jayathilakan K, Sultana K, Radhakrishna K, Bawa AS. Utilization of byproducts and waste materials from meat, poultry and fish processing industries: a review. *J Food Sci Technol*. 2012; 49: 278–293. <https://doi.org/10.1007/s13197-011-0290-7> PMID: 23729848
  34. Portillo A, Santibáñez S, García-Álvarez L, Palomar AM, Oteo JA. Rickettsioses in Europe. *Microbes Infect*. 2015; 17: 834–838. <https://doi.org/10.1016/j.micinf.2015.09.009> PMID: 26384814
  35. Levin ML, Killmaster LF, Zemtsova GE. Domestic dogs (*Canis familiaris*) as reservoir hosts for *Rickettsia conorii*. *Vector Borne Zoonotic Dis*. 2012; 12: 28–33. <https://doi.org/10.1089/vbz.2011.0684> PMID: 21923270
  36. Levin ML, Zemtsova GE, Montgomery M, Killmaster LF. Effects of homologous and heterologous immunization on the reservoir competence of domestic dogs for *Rickettsia conorii* (*israelensis*). *Ticks Tick Borne Dis*. 2014; 5: 33–40. <https://doi.org/10.1016/j.ttbdis.2013.07.010> PMID: 24201056
  37. Traversa D, Di Cesare A, Simonato G, Cassini R, Merola C, Diakou A, et al. Zoonotic intestinal parasites and vector-borne pathogens in Italian shelter and kennel dogs. *Comp Immunol Microbiol Infect Dis*. 2017; 51: 69–75. <https://doi.org/10.1016/j.cimid.2017.04.003> PMID: 28504099
  38. Antoniou M, Economou I, Wang X, Psaroulaki A, Spyridaki I, Papadopoulos B, et al. Fourteen-year seroepidemiological study of zoonoses in a Greek village. *Am J Trop Med Hyg*. 2002; 66: 80–85. PMID: 12135274
  39. Psaroulaki A, Spyridaki I, Ioannidis A, Babalis T, Gikas A, Tselentis Y. First isolation and identification of *Rickettsia conorii* from ticks collected in the region of Fokida in Central Greece. *J Clin Microbiol*. 2003; 41: 3317–3319. <https://doi.org/10.1128/JCM.41.7.3317-3319.2003> PMID: 12843081
  40. Psaroulaki A, Germanakis A, Gikas A, Scoulica E, Tselentis Y. First isolation and genotypic identification of *Rickettsia conorii* Malish 7 from a patient in Greece. *Eur J Clin Microbiol Infect Dis*. 2005; 24: 297–298. <https://doi.org/10.1007/s10096-005-1304-z> PMID: 15902541
  41. Germanakis A, Psaroulaki A, Gikas A, Tselentis Y. Mediterranean spotted fever in crete, Greece: clinical and therapeutic data of 15 consecutive patients. *Ann N Y Acad Sci*. 2006; 1078: 263–269. <https://doi.org/10.1196/annals.1374.049> PMID: 17114720
  42. Papa A, Dalla V, Petala A, Maltezou HC, Maltezos E. Fatal Mediterranean spotted fever in Greece. *Clin Microbiol Infect*. 2010. 16: 589–592. <https://doi.org/10.1111/j.1469-0691.2009.02910.x> PMID: 19673967
  43. Little SE. Ehrlichiosis and anaplasmosis in dogs and cats. *Vet Clin North Am Small Anim Pract*. 2010; 40: 1121–1140. <https://doi.org/10.1016/j.cvsm.2010.07.004> PMID: 20933140
  44. René-Martellet M, Lebert I, Chêne J, Massot R, Leon M, Leal A, et al. Diagnosis and incidence risk of clinical canine monocytic ehrlichiosis under field conditions in Southern Europe. *Parasit Vectors*. 2015; 8: 3–12. <https://doi.org/10.1186/s13071-014-0613-4> PMID: 25561342
  45. Parashar R, Sudan V, Jaiswal AK, Srivastava A, Shanker D. Evaluation of clinical, biochemical and haematological markers in natural infection of canine monocytic ehrlichiosis. *J Parasit Dis*. 2016; 40: 1351–1354. <https://doi.org/10.1007/s12639-015-0688-7> PMID: 27876946
  46. Fishbein DB, Dawson JE, Robinson LE. Human ehrlichiosis in the United States, 1985 to 1990. *Ann Intern Med*. 1994; 120: 736–743. PMID: 8147546
  47. Piantadosi D, Neola B, D'Alessio N, Di Prisco F, Santoro M, Pacifico L, et al. Seroprevalence and risk factors associated with *Ehrlichia canis*, *Anaplasma* spp., *Borrelia burgdorferi* sensu lato, and *D. immitis* in hunting dogs from southern Italy. *Parasitol Res*. 2017; 116: 2651–2660. <https://doi.org/10.1007/s00436-017-5574-z> PMID: 28776227
  48. Moraes-Filho J, Krawczak FS, Costa FB, Soares JF, Labruna MB. Comparative evaluation of the vector competence of four South American populations of the *Rhipicephalus sanguineus* group for the bacterium *Ehrlichia canis*, the agent of canine monocytic ehrlichiosis. *PLoS One*. 2015; 10: e0139386. <https://doi.org/10.1371/journal.pone.0139386> PMID: 26414283
  49. Estrada-Peña A, Roura X, Sainz A, Miró G, Solano-Gallego L. Species of ticks and carried pathogens in owned dogs in Spain: results of a one-year national survey. *Ticks Tick Borne Dis*. 2017; 8: 443–452. <https://doi.org/10.1016/j.ttbdis.2017.02.001> PMID: 28188085
  50. Latrofa MS, Angelou A, Giannelli A, Annoscia G, Ravagnan S, Dantas-Torres F, et al. Ticks and associated pathogens in dogs from Greece. *Parasit Vectors*. 2017; 10: 301–307. <https://doi.org/10.1186/s13071-017-2225-2> PMID: 28645329

51. Kachrimanidou M, Papa A, Chochlakis D, Pavlidou V, Psaroulaki A. Molecular evidence for *Anaplasma phagocytophilum* in *Ixodes ricinus* ticks from Greece. *Vector Borne Zoonotic Dis.* 2011; 11: 1391–1393. <https://doi.org/10.1089/vbz.2010.0251> PMID: 21923272
52. Papadopoulos B, Morel PC, Aeschlimann A. Ticks of domestic animals in the Macedonia region of Greece. *Vet Parasitol.* 1996; 63: 25–40. PMID: 8792578
53. Pavlidou V, Gerou S, Kachrimanidou M, Papa A. Ticks infesting domestic animals in northern Greece. *Exp Appl Acarol.* 2008; 45(3–4): 195–198. <https://doi.org/10.1007/s10493-008-9167-5> PMID: 18563592
54. Sainz Á, Roura X, Miró G, Estrada-Peña A, Kohn B, Harrus S, et al. Guideline for veterinary practitioners on canine ehrlichiosis and anaplasmosis in Europe. *Parasit Vectors.* 2015; 8: 75–95. <https://doi.org/10.1186/s13071-015-0649-0> PMID: 25649069
55. Hamel D, Silaghi C, Knaus M, Visser M, Kusi I, Rapti D, et al. Detection of *Babesia canis* subspecies and other arthropod-borne diseases in dogs from Tirana, Albania. *Wien Klin Wochenschr.* 2009; 121 Suppl 3: 42–45.
56. Davitkov D, Vucicevic M, Stevanovic J, Krstic V, Tomanovic S, Glavinic U, et al. Clinical babesiosis and molecular identification of *Babesia canis* and *Babesia gibsoni* infections in dogs from Serbia. *Acta Vet Hung.* 2015; 63: 199–208. <https://doi.org/10.1556/AVet.2015.017> PMID: 26051258
57. Rubel F, Brugger K, Pfeffer M, Chitimia-Dobler L, Didyk YM, Leverenz S, et al. Geographical distribution of *Dermacentor marginatus* and *Dermacentor reticulatus* in Europe. *Ticks Tick Borne Dis.* 2016; 7: 224–233. <https://doi.org/10.1016/j.ttbdis.2015.10.015> PMID: 26552893
58. Gramiccia M, Gradoni L. The current status of zoonotic leishmaniases and approaches to disease control. *Int J Parasitol.* 2015; 35: 1169–1180.
59. Strazzulla A, Cocuzza S, Pinzone MR, Postorino MC, Cosentino S, Serra A, et al. Mucosal leishmaniasis: an underestimated presentation of a neglected disease. *Biomed Res Int.* 2013; 2013: 805108. <https://doi.org/10.1155/2013/805108> PMID: 23853773
60. Genchi C, Kramer L. Subcutaneous dirofilariosis (*Dirofilaria repens*): an infection spreading throughout the old world. *Parasit Vectors.* 2017; 10(Suppl 2): 517. <https://doi.org/10.1186/s13071-017-2434-8> PMID: 29143643
61. Diakou A, Kapantaidakis E, Tamvakis A, Giannakis V, Strus N. *Dirofilaria* infections in dogs in different areas of Greece. *Parasit Vectors.* 2016; 9: 508–514. <https://doi.org/10.1186/s13071-016-1797-6> PMID: 27646111
62. Miró G, Montoya A, Roura X, Gálvez R, Sainz A. Seropositivity rates for agents of canine vector-borne diseases in Spain: a multicentre study. *Parasit Vectors.* 2013; 6: 117. <https://doi.org/10.1186/1756-3305-6-117> PMID: 23607428
63. Alho AM, Pita J, Amaro A, Amaro F, Schnyder M, Grimm F, et al. Seroprevalence of vector-borne pathogens and molecular detection of *Borrelia afzelii* in military dogs from Portugal. *Parasit Vectors.* 2016; 9 (1): 225. <https://doi.org/10.1186/s13071-016-1509-2> PMID: 27160284
64. Vascellari M, Ravagnan S, Carminato A, Cazzin S, Carli E, Da Rold G, et al. Exposure to vector-borne pathogens in candidate blood donor and free-roaming dogs of northeast Italy. *Parasit Vectors.* 2016; 9: 369. <https://doi.org/10.1186/s13071-016-1639-6> PMID: 27357128