

Table S1. Knowledge and practices about chikungunya in the ≥ 15 years study population

	N (%)	Seroprevalence (95%CI)
Considers him/herself informed about chikungunya		
<i>Yes</i>	793 (93.5)	13.2 (10.9, 15.6)
<i>No</i>	55 (6.5)	10.9 (2.4, 19.4)
Main source of information about chikungunya		
<i>TV</i>	613 (77.3)	
<i>Radio</i>	93 (11.7)	
<i>Community health workers</i>	45 (5.7)	
<i>Talks at health center/post</i>	23 (2.9)	
<i>Newspapers</i>	16 (2.0)	
<i>Other</i>	3 (0.4)	
Correctly identifies mosquitoes as vector of transmission		
<i>Yes</i>	766 (90.3)	13.6 (11.2, 16.2)
<i>No</i>	82 (9.7)	8.5 (3.5, 16.8)
Implements at least one measure of chikungunya control		
<i>Yes</i>	758 (95.4)	13.3 (10.9, 15.9)
<i>No</i>	90 (10.6)	11.1 (5.5, 19.5)
Practices of chikungunya prevention*		
<i>Elimination of mosquito breeding sites</i>	620 (73.1)	
<i>Covering water containers</i>	482 (56.8)	
<i>Mosquito nets</i>	411 (48.5)	
<i>Repellent</i>	374 (44.1)	
<i>Cleaning water containers</i>	325 (38.3)	
<i>Window screens</i>	159 (18.8)	

* Multiple choices possible.