

Muscles in “Concert”: Study of Primary Motor Cortex Upper Limb Functional Topography

Jean-Marc Melgari^{1*}, Patrizio Pasqualetti^{2,3}, Flavia Pauri⁴, Paolo Maria Rossini^{1,2,3}

1 Neurology, University Campus Bio-Medico, Rome, Italy, **2** Medical Statistics & Information Technology, AFaR, Fatebenefratelli Hospital, Isola Tiberina, Rome, **3** Casa di Cura SAN RAFFAELE Cassino and IRCCS SAN RAFFAELE, Pisana, Rome, Italy, **4** Department of Neurology and ORL, Università La Sapienza, Rome, Italy

Abstract

Background: Previous studies with Transcranial Magnetic Stimulation (TMS) have focused on the cortical representation of limited group of muscles. No attempts have been carried out so far to get simultaneous recordings from hand, forearm and arm with TMS in order to disentangle a ‘functional’ map providing information on the rules orchestrating muscle coupling and overlap. The aim of the present study is to disentangle functional associations between 12 upper limb muscles using two measures: cortical overlapping and cortical covariation of each pair of muscles. Interhemispheric differences and the influence of posture were evaluated as well.

Methodology/Principal Findings: TMS mapping studies of 12 muscles belonging to hand, forearm and arm were performed. Findings demonstrate significant differences between the 66 pairs of muscles in terms of cortical overlapping: extremely high for hand-forearm muscles and very low for arm vs hand/forearm muscles. When right and left hemispheres were compared, overlapping between all possible pairs of muscles in the left hemisphere (62.5%) was significantly higher than in the right one (53.5%). The arm/hand posture influenced both measures of cortical association, the effect of Position being significant [$p = .021$] on overlapping, resulting in 59.5% with prone vs 53.2% with supine hand, but only for pairs of muscles belonging to hand and forearm, while no changes occurred in the overlapping of proximal muscles with those of more distal districts.

Conclusions/Significance: Larger overlapping in the left hemisphere could be related to its lifetime higher training of all twelve muscles studied with respect to the right hemisphere, resulting in larger intra-cortical connectivity within primary motor cortex. Altogether, findings with prone hand might be ascribed to mechanisms facilitating coupling of muscles for object grasping and lifting –with more proximal involvement for joint stabilization– compared to supine hand facilitating actions like catching. TMS multiple-muscle mapping studies permit a better understanding of motor control and ‘plastic’ reorganization of motor system.

Citation: Melgari J-M, Pasqualetti P, Pauri F, Rossini PM (2008) Muscles in “Concert”: Study of Primary Motor Cortex Upper Limb Functional Topography. PLoS ONE 3(8): e3069. doi:10.1371/journal.pone.0003069

Editor: Brian D. McCabe, Columbia University, United States of America

Received: January 26, 2008; **Accepted:** July 28, 2008; **Published:** August 27, 2008

Copyright: © 2008 Melgari et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: AFaR, Fatebenefratelli Hospital-Isola Tiberina, Roma

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: j.melgari@unicampus.it

Introduction

About 30 years ago a revolutionary technique was introduced which allowed stimulation of the human brain through the skull with non-invasive, high-intensity, extremely brief electric pulses [1–5], followed –few years later– by a new device that employs strong and brief, time-varying magnetic fields able to induce electric currents flowing within the brain without any discomfort [6]; for a review see [7]. Since then, focal types of stimulating coils able to activate relatively small cortical areas allowing cortical mapping were developed [8]. Therefore, focal TMS to different cells of a grid overlying the primary motor cortex (M1) while recording electromyographic (EMG) responses from the contralateral ‘target’ muscles, made it possible to obtain reliable maps of cortical motor output in awake and cooperative subjects [8–21]. Due to technical limitations (i.e. number of recording amplifiers) and to the burden of post-hoc analysis, all the reports regard one (the vast majority) to a maximum of 4 upper limb individual muscles simultaneously analysed [10,22,23]; in this way a roughly

“somatotopic gradient” similar to *penfieldian* organization –in which wide overlapping was observed– was consistently reported.

Different studies demonstrated both in primates and in humans that several loci along the pre-central motor strip contain separate representations of the same muscle (convergence). Meanwhile, it was shown that the same cortical site when focally stimulated can dispatch outgoing impulses to different muscles (divergence) and that muscles acting at different upper limb joints are represented at different cortical foci where they act in concert with different companion muscles. It has been shown that such parameters are highly influenced by limb posture [24–28]. Convergence, divergence, multiple muscle cortical representations and the influence of posture seem to play a pivotal role also in humans, orchestrating several muscles ‘in concert’ during motor actions, motor learning and post-lesional ‘plastic’ brain reorganization [29–32]. Even if invasive stimulation/recording procedures allowing a direct evaluation of these phenomena cannot be used in humans, TMS offers a valuable probe for a better knowledge of the dynamics of the cortical topography of individual muscles

when they are treated as an *ensemble* and not as individual actors.

The aim of the present study was to investigate overlap and covariation of cortical motor output during TMS mapping, and simultaneous recordings from 12 individual upper limb muscles in order to measure cortical overlapping and covariation of pairs of muscles, interhemispheric differences and influence of arm/hand posture. Two measures were chosen to indicate overlap and covariation: 1) cortical overlapping of each pair of muscles, defined as the percentage of grid positions where TMS elicited a MEP response in both muscles respect to the total number of grid positions where TMS elicited a MEP response in at least one of the two, and 2) covariation of each pair of muscles, measured by means of Pearson's correlation. These measures make it possible to disentangle two neurophysiological aspects of functional connectivity: first, overlapping is related to the co-activation of the target muscles and indicates how much pairs of muscles are simultaneously represented in the motor cortex; second the covariation could be an index of the intensity and the direction of the co-activation.

Results

The experimental setting and electrode positioning for simultaneous MEP recordings from the examined muscles is represented in Fig. 1, while traces of the original MEPs obtained from 12 muscles of subject 2 are shown in Fig. 2.

Right hemisphere, left prone hand (10 subjects)

The 66 pairs of muscles showed significant differences for clusters of pairs sharing similar levels of cortical overlapping [$F(65, 590) = 14.221, p < .001$] as shown in Table 1 by Tukey's homogenous subsets (p value higher than 0.90).

A first subset is characterised by a low overlapping level (range: 12% for EIP-TRI, 29% for TRI-BIC) and is clearly identified by all pairs including at least one of the three arm muscles. A second group of pairs is characterized by a high overlapping level (range: 64% for EIP-EUC, 96% for EUC-OPP) which comprises almost all hand-hand, hand-forearm and forearm-forearm pairs. Between these two subsets, there is not a unique and well defined subset, but it can be noted that all pairs are characterised by the presence of EIP muscle: the cortical representation of this muscle overlaps with the other forearm and hand muscles in a range between 56% and 67%, reaching the maximum with FDI (72%). The number of co-activated muscles for each scalp position in one paradigmatic subject is represented in Fig. 3.

Similar analysis was applied to the other measure of association. In this case, we also tested the significance of the correlation index for each pair of muscles with respect to the lack of correlation (0 value). When the 95% confidence intervals did not include the 0 reference line, a correlation (negative or positive) can be documented (with α error at 5%). The size of the correlation index changed according to which specific pair of muscles was considered [ANOVA $F(65, 367) = 2.205, p < .001$]. Looking at Tukey's homogenous subsets, two




Figure 1. Experimental setting and electrode positioning for simultaneous MEP recordings from the examined muscles.

doi:10.1371/journal.pone.0003069.g001




Figure 2. Original MEPs obtained from 12 muscles simultaneously at a two different stimulation sites, corresponding to: left panel, hot-spot of FDI (with respect to Cz, x=3 cm, y=2 cm); right panel, hot-spot of Deltoid (with respect to Cz, x=1 cm, y=2 cm); time and amplitude calibrations are 20 msec and 200 microV respectively.
doi:10.1371/journal.pone.0003069.g002

main subsets were identifiable: 1) pairs including at least one arm muscle; 2) pairs including hand and forearm muscles. However, the standard error of the mean for this index is much larger than for the overlapping one and it was more difficult to assess whether some pairs were statistically more correlated than others. Meanwhile, graphical representation with 95% error bars (Fig. 4) showed that in the first subset (pairs including at least one arm muscle) correlation always crossed the 0 reference line (lack of correlation), while the second subset was consistently characterised by high correlation levels up to the most correlated pair, i.e. EUC-OPP (untransformed $r = 0.786$).

The patterns observed for the two measures of cortical association in the right hemisphere were used as baseline *vs* the left hemisphere and when the influence of the hand position (prone *vs* supine) was analysed.

Comparison between right and left hemispheres (right prone hand: 6 subjects)

When right and left hemispheres were compared in terms of activation—even if with a quite similar pattern of muscle associations (Fig. 5, right panel)—a significant *Hemisphere* effect was found [$F(1, 659) = 21.199, p < .001$], indicating that the overlapping between all possible pairs of muscles was significantly higher in the left (mean

overlapping 62.5%; 95% CI=59.8–65.3) than in the right hemisphere (mean overlapping 53.5; 95% CI=50.8–56.2) (Fig. 5). No interaction *Hemisphere*×*Muscle_pairs* was observed [$F(65, 659) = 0.536, p = .999$] also when pairs were grouped according to the upper limb district [$F(5,779) = 0.817, p = .538$].

A different pattern was observed when the covariation measure was analysed. In this case, no evidence of higher association in left *vs* right hemisphere was found [$F(1,526) = 1.515, p = .219$], mean correlation being 0.37 (95% CI=0.31–0.43) in the latter and 0.32 (95% CI: 0.27–0.37) in the former. Since higher correlation—although not significant—was observed in the right hemisphere, further analysis was performed in order to compare the two association measures in the two hemispheres (after converting them into a common scale, i.e. z-scores). This analysis indicated a significant *Association_measure*×*Hemisphere* interaction [$F(1,656) = 7.014; p < .008$], supporting the hypothesis of a different kind of association between muscles in the two hemispheres.

Comparison between supine and prone hand (left hand-right hemisphere, 3 subjects)

Hand posture influenced both measures of cortical association. In fact, the effect of *Position* was significant [$F(1,392) = 5.419,$

Table 1. Percentage of overlapping between each of the 66 pairs of muscles.

Pairs	Homogeneous subsets (Tukey's procedure)																		
	1	2	3	4	5	6	7												
EIP-TRI	12																		
DEL-EIP	12																		
DEL-FDI	15																		
DEL-OPP	15																		
DEL-EUC	15																		
BIC-EIP	15																		
ADM-DEL	16																		
ERC-TRI	16																		
DEL-ERC	16																		
DEL-EDC	16																		
DEL-FLC	17																		
FDI-TRI	17																		
ABD-DEL	17																		
OPP-TRI	17																		
EUC-TRI	17																		
ADM-TRI	18																		
FLC-TRI	18																		
DEL-TRI	18																		
EDC-TRI	19																		
ABD-TRI	19																		
BIC-FDI	20																		
BIC-OPP	21																		
BIC-ERC	21																		
BIC-EUC	21																		
ADM-BIC	22																		
BIC-EDC	22																		
ABD-BIC	23																		
BIC-DEL	23																		
BIC-FLC	24	24																	
BIC-TRI	29	29	29																
EIP-ERC		56	56	56															
EDC-EIP			58	58	58														
EIP-FLC				62	62	62													
EIP-OPP					64	64	64												
EIP-EUC						64	64												
ADM-EIP							67												
ABD-EIP								67											
ERC-FDI									69										
EDC-ERC										72									
EIP-FDI											72								
ERC-FLC												73							
EDC-FDI													74						
ERC-OPP														74					
ERC-EUC															75				
ADM-ERC																76			
FDI-FLC																	77		
ABD-ERC																		78	
EDC-FLC																			80

Table 1. cont.

Pairs	Homogeneous subsets (Tukey's procedure)						
	1	2	3	4	5	6	7
ABD-FDI				82	82	82	82
EDC-EUC				82	82	82	82
EDC-OPP				83	83	83	83
ADM-EDC				84	84	84	84
FLC-OPP				84	84	84	84
EUC-FLC				85	85	85	85
EUC-FDI				85	85	85	85
ABD-EDC				86	86	86	86
ADM-FLC				86	86	86	86
FDI-OPP				86	86	86	86
ADM-FDI				86	86	86	86
ABD-FLC				87	87	87	87
ABD-EUC					90	90	90
ABD-OPP					90	90	90
ABD-ADM						91	91
ADM-EUC						93	93
ADM-OPP						94	94
EUC-OPP							96
<i>p-value</i>	<i>0.999</i>	<i>0.906</i>	<i>0.978</i>	<i>0.951</i>	<i>0.922</i>	<i>0.931</i>	<i>0.901</i>

doi:10.1371/journal.pone.0003069.t001

$p = .021$] on co-activation with a 59.5% overlap (95% CI = 55.8–63.3) with prone hand and a 53.2% overlap (95% CI = 49.4–57.0) with supine hand, respectively. However, the larger overlapping with prone position was specifically evident for hand and forearm muscle pairs, while no changes occurred in the overlapping of proximal muscles with the more distal ones (Fig. 6).

The correlation between pairs of muscles resulted higher with prone than with supine hand [$F(1,160) = 7.025, p = .009$]. The presence of a significant *Couple_district* × *Position* interaction [$F(5,280) = 2.739, p = .020$] could be ascribed to the lower correlation between proximal and distal districts in case of supine hand.




Figure 3. TMS-MRI integration: number of co-activated muscles for each scalp position in a paradigmatic subject.

doi:10.1371/journal.pone.0003069.g003




Figure 4. Covariation measure of each pair between the 12 examined muscles. This measure was obtained from the right hemisphere of 10 healthy subjects lying with prone hands.
doi:10.1371/journal.pone.0003069.g004




Figure 5. Inter-hemispheric comparison of co-activation and covariation measures (obtained with hand in prone position) between pairs of muscles belonging to three different districts (hand, forearm, arm).
doi:10.1371/journal.pone.0003069.g005

Discussion

Modern concepts on the functional organization of cortical primary motor areas maintain a somatotopic view only for major body districts (such as face, arm, leg), each being sustained by a network serving broadly distributed functions which involve large populations of neurons in spatially separate clusters. Within this theoretical frame, the main goal of the present study was to investigate whether the use of a non-invasive technique like TMS could roughly disentangle the representational overlapping between different upper limb muscles and the eventual functional correlation in humans. [33–37].

The existence of multiple yet discrete efferent micro- and macrozones from primary motor cortex, is now accepted as the essential organizational principle of this area. Animal studies have in fact demonstrated that a particular movement can be elicited through stimulation of different M1 regions, often several millimeters apart and separated by non-responsive districts [24–27,39–43]. In addition, bi-directional projections interconnect motor cortex areas for different muscle districts [24,44] for a review. Any movement therefore, differently from the traditional view of the *labelled-line* hypothesis, is probably not controlled by single aggregates, but by a distributed network within the M1 cortex. In this model, the motor output from overlapping cortical territories converge onto individual muscles. Similarly, the output from any given cortical site diverges onto multiple muscles with different ‘gain’ according to the final movement to be performed, presumably regulated by horizontal intracortical projections interconnecting functionally related neuronal clusters within M1 [43,45,46].

Multiple representations of the cortical output to individual muscles, overlap each other. Multiple cortical representation of muscles controlling hand/fingers, represents the functional frame for the extraordinary repertoire of movement strategies (i.e. concerted actions orchestrated by many –sometimes the same-muscles acting on different joints at different times [24,37,47]). Such an organization also provides flexibility in motor planning and execution, and progressively substitutes a related dysfunctional area more easily than highly specialized and unique groups of cells, being thus fundamental either for restorative or for maladaptive plasticity.

Precise methods such as those in experimental models, allowing microneurographic stimulation/recording procedures from individual neurons/fibers in the motor cortex, pyramidal tract and spinal motoneurons cannot be used in humans to address physiological mechanisms for movement control like convergence, divergence and multiple muscle representation. Several techniques for functional brain imaging –namely PET and fMRI– allow detailed images on the relationship between function and anatomy showing the distributed network subtending a given motor act. However they cannot discriminate the temporal dynamics of a phenomenon taking place in few tens of milliseconds or differentiate neuronal firing decrease from increase (exciting *vs* inhibiting net effects); moreover, in the case of movement paradigms, the activation directly linked to motor programming and execution cannot be distinguished from the sensory input feed-back from the moving parts, and the chronological hierarchy –if any– governing activated areas cannot be studied appropriately [48,49]. Despite their potential ability to properly follow-up such a




Figure 6. Inter-position comparison of co-activation and covariation measures (obtained from the right hemisphere) between pairs of muscles belonging to three different districts (hand, forearm, arm).

doi:10.1371/journal.pone.0003069.g006

brain mechanism, non-invasive brain stimulation procedures have never been tested along this vein in human beings, despite their progressively growing use in research for movement physiology and pathophysiology see [7] for a review. In fact, transcranial magnetic stimulation of the primary motor cortex has largely been employed in investigating neurophysiological mechanisms of motor control in the healthy subjects [9,11,18,19,50–65] and in patients with neurological motor deficits [13,15,66–71]. Mapping procedures of motor cortical output have been progressively implemented and integrated with precise anatomical brain reconstructions [23,32,33,72–78]. However, no previous studies investigated in detail the functional linkage amongst upper limb muscles simultaneously recorded during TMS mapping of output from M1 cortex.

We document significant overlapping between proximal and distal muscles, in line with many animal studies involving the superposition of topographical maps of the motor cortex obtained by microstimulation and morphological connectivity maps obtained by tracer injections. The concordance between such studies and our TMS findings (related to trans-synaptic activation) could also be referred to the existence of intrinsic horizontal collaterals (intracortical connections) [79].

Findings from the present study demonstrate significant difference between the 66 pairs of muscles in terms of cortical overlapping: this index reached high scores for hand-forearm muscles and low for arm *vs* hand/forearm muscles. Moreover, a subset of low-overlapping muscles characterised by a low percentage of overlap was clearly identified by all pairs with at least one of the three arm muscles. In a separate cluster, there is a

group of pairs with high-overlapping muscles which include almost all hand-hand, hand-forearm and forearm-forearm possible pairs.

When covariation was taken into consideration interesting observations emerged. In particular, our data were obtained at rest and there was a lack of covariation between forearm/hand and more proximal arm muscles. Our covariation measure gives an evidence of the motor cortex representation of “muscle synergies” (in terms of the directionality of a coactivation) that should not be considered hard-wired entities. In fact, in the modern view of an integrated nature of motor cortical function, movement-related muscle synergies “may be dynamically created by the operations of motor cortical circuitry” [79–80].

Moreover, high covariations for wrist extensors and hand muscles were seen up to the most correlated pair EUC-OPP. One might argue that this functional organization of M1 output is strongly devoted to the control of wrist joint.

When right and left hemispheres were compared, a significant *Hemisphere* effect was found indicating that the index of overlapping in the left was significantly higher than in the right hemisphere. Even if this observation is intriguing, whether it has some relationship with manual dexterity and hemispheric dominance for hand motor preference, needs further evaluation in a population with right *vs* left-hand dexterity. It is worth noting that a larger overlapping in the left hemisphere, without a corresponding inter-hemispheric difference of the single muscle cortical representation was found. This larger overlapping in the left hemisphere could be related to its lifetime higher training of all twelve studied muscles with respect to the right hemisphere, resulting in a larger connectivity within M1 cortex.

Interhemispheric difference was not observed when the covariation measure was analysed. Covariation could be interpreted as an index of the intensity and direction of the co-activation; therefore, the larger overlapping observed in the left hemisphere did not involve a higher co-variation.

Of course, covariation/overlapping measures are significantly correlated ($r = 0.49$), indicating that about 25% of the variance of one could be accounted for by the other. In fact, to calculate the covariation the two muscles have to be simultaneously represented in the motor cortex (that is to say, there must be coactivation/overlapping of the two muscles). So we can affirm that overlapping is the basic phenomenon on which the covariation takes place, but also that covariation could be considered as an index of the intensity and direction of the co-activation. In other words, co-activation and covariation could be summarised only partially by one of them. The contamination between these two measures, although present, is not sufficient to conclude that they indicated the same phenomenon. In addition, the similarity of patterns represented in the two panels of Fig. 5 is only apparent, since co-activation was higher in right than in left hemisphere (left panel), while covariation was not different in the two hemispheres. For these reasons (the first neurophysiological, the second statistical) the two measures can reflect two different neurophysiological aspects of functional connectivity.

Network structure can adjust dynamically to meet the immediate needs of the motor system, as seen in response to postural changes [36]; indeed, in the present study hand position significantly influenced both overlapping and covariation indexes. The larger overlapping with prone hand seemed to be specific to pairs of muscles belonging to hand and forearm, while no changes occurred for proximal muscles. Therefore, this effect seemed specific to the muscles and districts where the main change in proprioceptive input occurred, and can be linked to task-devoted organization of synaptic contacts at the periphery of cortical maps of the muscles involved. Altogether, present findings with prone hand might be ascribed to cortical mechanisms facilitating coupling of muscle pairs for object' grasping and lifting -which needs involvement of more proximal muscles for joint stabilization and increased upper limb stiffness- compared to supine hand, facilitating a motor action most apt for object catching on a vertical axis, with less proximal muscle involvement [81].

The potential effect of different peripheral nerve territories involvement on association measures was also examined, but neither overlapping nor covariation indexes were statistically linked to muscle pairs innervated by branches belonging to the same nerve trunk respect to the pairs controlled by different nerves. This indicated that the observed patterns of muscle associations, take place within CNS, even if the present approach cannot discriminate between cortical and subcortical (including spinal) contributions.

In conclusion, simultaneous upper limb multiple-muscle recordings during mapping of M1 output with focal TMS has provided a bulk of information strongly related to the functionality of the arm/hand system in healthy humans. This approach will hopefully permit better understanding of upper limb motor control in healthy subjects and in patients with motor disorders, by providing prognostic indexes for motor recovery and in following-up 'plastic' reorganization of the motor system after acute or progressive lesions.

Materials and Methods

Mapping studies with TMS were performed on 10 healthy volunteers (6 females and 4 males), right handed, aged 28–40. The experimental protocol had the approval of the ethics committee of

Fatebenefratelli Hospital-Isola Tiberina (Rome) and all subjects gave written informed consent; international safety standards – including an EEG before subject were recruited for TMS- were strictly followed [20,82].

Simultaneous EMG records were obtained from the following 12 muscles in different experimental conditions: *Abductor Digiti Minimi* (ADM), *First Interosseus Dorsalis* (FDI), *Opponens Pollicis* (OPP), *Abductor Pollicis Brevis* (APB), *Extensor Digitorum Communis* (EDC), *Extensor Indicis Proprius* (EIP), *Extensor Carpi Ulnaris* (EUC), *Extensor Carpi Radialis* (ERC), *Flexor digitorum superficialis (communis)* (FLC), *Biceps brachii* (BIC), *Triceps* (TRI), *Deltoid* (DEL).

During the experimental session the subject, wearing a transparent and well-fitting plastic cap, lay supine on a bed in order to facilitate complete relaxation and was kept alert by the investigator.

Six subjects attended for two testing procedures, one for each hemisphere, with arm, forearm and hand relaxed in prone position. In three of them, another recording session was carried out only for right hemisphere with arm, forearm and hand relaxed in supine position. This further recording was performed in order to assess eventual modifications induced on cortical somatotopy by arm/hand posture. The remaining subjects only had one hemisphere and one hand position studied.

Focal single-pulse TMS was delivered to the scalp using a flat figure-of-eight coil with an inner diameter of 70 mm for each wing connected to a Magstim 200 magnetic stimulator (Magstim Company, Whitland, UK).

The coil was placed tangentially to the skull with the handle pointing backward and rotated away from the midline by 45°.

Stimulus intensity was adjusted at resting motor threshold defined on the hot spot following international guidelines as the stimulator's output able to elicit reproducible MEPs (at least 50 μ V in amplitude) in about 50% of 10–20 consecutive stimuli in a completely relaxed *Opponens Pollicis* muscle [20,82]. Once the excitability threshold was determined, TMS intensity was increased by 10% of the stimulator output to enhance response probability. This intensity was an obvious compromise between the different excitability thresholds of the upper limb muscles and is usually able to elicit responses with a decreasing probability from hand to shoulder muscles (the latter having a higher threshold and stronger intensities requirement). This was a forced choice in order to avoid an excessive "cortical spread" of the stimulus which would have occurred if we had utilized 110% intensity of the threshold of the less excitable (i.e. arm/shoulder) muscles.

The hot spot of the OPP, defined as the point from which stimuli triggered Motor Evoked Potentials (MEPs) of maximal amplitude and minimal latency, was found during a pilot stimulating/recording session with a frankly suprathreshold stimulus intensity. The hot spot position was then marked on a transparent, adherent elastic cap to facilitate an exact repositioning of the coil during the entire experiment. In one paradigmatic subject 3D head/brain Magnetic Resonance Imaging (MRI) was obtained, coregistered with the grid of recording scalp electrodes as represented by vitamin E capsules. Therefore, data presented in Fig. 3 are realistic and not merely a graphical representation.

Using surface electrodes, EMG activity was recorded simultaneously from each of the 12 target muscles. Skin was cleaned at the recording site followed by placement of Ag/AgCl cup electrodes filled with conductive gel in a belly/tendon montage. Recording electrodes were placed over the bellies (active) and tendons (reference) of muscles to be monitored and ground a electrode was placed over the wrist (Fig. 1). Interelectrode distance was maintained as short as possible; this -together with the TMS

intensity just above threshold- was deliberately chosen in order to reduce the amount of cross-talk from muscles different and adjacent from the one under examination. Skin/electrode resistances were kept <10 KOhms.

Signal recording was carried out with a modified version of the BASIS-Brain Surveyor Equipment (EB-Neuro; 12 channels; sampling rate 5 KHz/channel, filtering bandpass 3–1500 Hz).

Subjects were wearing an elastic, transparent cap containing a grid of 121 numbered squares marked at 1 cm intervals, in which square 1 corresponded to the designated hot-spot of the OPP hand muscle used to define the excitability threshold. The remaining squares were subsequently numbered in a spiral fashion. The scalp vertex (the point on the midsagittal line, midway between nasion and inion) and the scalp anatomical landmarks (nasion, inion) were taped in a stable position on the cap.

Four consecutive MEPs were gathered from each grid position starting from square 1 and following the progressive numbering with a stimulus repetition rate of 0.1–0.2 c/sec. The progression of stimulated sites of the scalp grid followed a spiral sequence, that could be considered a relatively safe method to avoid sequential stimulation of an ordered and topographically organized cortical representation of the muscles.

MEPs latency and amplitude were measured at peak onset and peak-to-peak of opposite polarity, respectively; they were calculated off-line for each stimulated site and for each muscle. Particular care was devoted to maintain full relaxation of the examined muscles and trials contaminated by involuntary movement or muscle activity were discarded from further analysis (Fig. 2).

Data analysis

A data matrix containing all raw latencies and amplitudes of MEPs was prepared for each of the 12 muscles elicited by stimulation on each of 121 scalp positions corresponding to the 11×11 grid, centred around the hot-spot. Since for each position, four successive stimuli were delivered, the matrix contained 424 rows for each subject. Ten subjects underwent TMS of the right hemisphere (with prone left hand), 6 subjects underwent the TMS of both hemispheres (with prone hands) and 3 subjects underwent the TMS of right hemisphere (with prone and supine left hand). Altogether 20 different TMS sessions entered data analysis. Therefore, the complete raw dataset consisted of 8056 rows and 27 columns (the first three to identify subjects, hemisphere and hand position, then 12 for latencies and 12 for amplitudes). It is worth noticing that a number of the matrix cells were empty, since no MEP was obtained from the corresponding scalp positions and muscles.

In order to reduce the large variability of MEP amplitude data, *log* transformation was applied [83]. In addition, the median value was considered for each scalp position: since four stimuli were delivered on each stimulation site, median value was the arithmetic mean in the case two MEPs were obtained, the unique central value in the case three MEPs were obtained, the arithmetic mean of the two central values after discarding the two extremes in case four MEPs were obtained. When just one MEP was observable, the amplitude was set at a missing value.

Measures for cortical associations between muscle representations

Many studies faced the issue of mapping the motor area and nowadays each mapping study should rely on a neuronavigation

tool allowing the integration between TMS and neuroimaging techniques. Thanks to such procedures, the cortical motor area of each muscle could be rigorously described and analysed by means of parameters characterizing such areas as excitability (number and extension of “active” points on the scalp), volume (sum of MEP amplitudes obtained stimulating the whole active area), center-of-gravity (the amplitude-weighted mean position) [10,14,76,84]. These parameters make it possible to address the issues of “convergence” (TMS on several different scalp positions produce a MEP in the same target muscle), “divergence” (TMS on a single scalp position produce a MEP in different target muscles) and somatotopy of motor area (based mainly on hot-spot and centre-of-gravity positions). It is however worth remembering that the aim of the present study is not to replicate previous mapping findings, but to describe and investigate the association between upper limb muscles in terms of their cortical representation. Since multivariate techniques, such as Multi-Dimensional Scaling (MDS) did not provide consistent findings, two simple measures of association between muscles were computed:

- 1) cortical overlapping of each pair of muscles, defined as the percentage of grid positions where TMS elicited a MEP response in both muscles respect to the total number of grid positions where TMS elicited a MEP response in at least one of the two muscles. For example, the overlapping between FDI and ADM was: $N_{FDI \cap ADM} / N_{FDI \cup ADM}$.
- 2) cortical covariation of each pair of muscles, measured by means of Pearson’s correlation based on MEP amplitude. To be noted that Pearson’s correlation was transformed according to Fisher’s transformation $0.5 \cdot \ln((1-r)/(1+r))$ in order to obtain a measure more suitable for statistical tests (that will be denoted hereafter *rPF*).

Neurophysiological interpretation of cortical overlapping mainly relies on co-activation of the target muscles and indicates how much two muscles are simultaneously represented in the motor cortex. On the other hand, covariation could be interpreted as an index of the intensity and direction of co-activation: for example, a negative correlation between two muscles would result when high MEP amplitude in one muscle corresponded to low MEP amplitude in the other and vice versa. In other words, this would be the case when, although simultaneously represented, a MEP in one muscle of a pair is elicited by recruitment of few 1st and 2nd motoneurons (small response with respect to the average) while the other is elicited by a larger pool of motoneurons (large response with respect to the average), provided the same scalp site is stimulated in both conditions.

Since 12 muscles were triggered by each stimulus, 66 pairs could be defined. During the statistical analysis and for graphic presentation, muscles were grouped in three districts (4 hand muscles-Group I, 5 forearm muscles-Group II, 3 arm muscles-Group III). The 66 muscle pairs were therefore distributed as follows: 6 hand-hand, 20 hand-forearm, 12 hand-arm, 10 forearm-forearm, 15 forearm-arm, 3 arm-arm.

Author Contributions

Conceived and designed the experiments: JMM PP PMR. Performed the experiments: JMM FP PMR. Analyzed the data: JMM PP. Wrote the paper: JMM PP FP PMR.

References

1. Merton PA, Morton HB (1980) Stimulation of the cerebral cortex in the intact human subjects. *Nature* 285: 227.
2. Merton PA, Morton HB (1980) Electrical stimulation of human motor and visual cortex through the scalp. *J Physiol* 305: 9–10.

3. Merton PA, Hill DK, Morton HB, Marsden CD (1982) Scope of a technique for electrical stimulation of human brain, spinal cord, and muscle. *Lancet* 2: 597–600.
4. Rossini PM, Marciani MG, Caramia M, Roma V, Zarola F (1985) Nervous propagation along “central” motor pathways in intact man: characteristics of motor responses to “bifocal” spine and scalp non-invasive stimulation. *Electroencephalogr Clin Neurophysiol* 61: 272–286.
5. Rossini PM, Di Stefano E, Stanzione P (1985) Nerve impulse propagation along central and peripheral fast conducting motor and sensory pathways in man. *Electroencephalogr Clin Neurophysiol* 60: 320–334.
6. Barker AT, Jalinous R, Freeston IL (1985) Non-invasive magnetic stimulation of the human motor cortex. *Lancet* 1: 1106–1107.
7. Rossini PM, Rossi S (2007) Transcranial magnetic stimulation: diagnostic, therapeutic, and research potential. *Neurology* 68: 484–8.
8. Cohen LG, Roth BJ, Nilsson J, Dang N, Panizza M, et al. (1990) Effects of coil design on delivery of focal magnetic stimulation. Technical considerations. *Electroencephalogr Clin Neurophysiol* 75: 350–357.
9. Rothwell JC, Day BL, Thompson PD, Dick JP, Marsden CD (1987) Some experiences of techniques for stimulation of the human cerebral motor cortex through the scalp. *Neurosurgery* 20: 156–163.
10. Wassermann EM, McShane LM, Hallett M, Cohen LG (1992) Noninvasive mapping of muscle representations in human motor cortex. *Electroencephalogr and Clinical Neurophysiol* 85: 1–8.
11. Pascual-Leone A, Cohen LG, Brasil-Neto JP, Hallett M (1994) Non-invasive differentiation of motor cortical representation of hand muscles by mapping of optimal current directions. *Electroencephalogr Clin Neurophysiol* 93: 42–48.
12. Rossini PM, Rossi S (1998) Clinical Application of Motor Evoked Potentials. *Electroencephalogr Clin Neurophysiol* 106: 180–194.
13. Traversa R, Cicinelli P, Bassi A, Rossini PM, Bernardi G (1997) Mapping of motor cortical reorganization after stroke. A brain stimulation study with focal magnetic pulses. *Stroke* 28: 110–117.
14. Cicinelli P, Traversa R, Bassi A, Scivoletto G, Rossini PM (1997) Interhemispheric differences of hand muscle representation in human motor cortex. *Muscle Nerve* 20: 535–542.
15. Cicinelli P, Traversa R, Rossini PM (1997) Post-stroke reorganization of brain motor output to the hand: a 2–4 month follow-up with focal magnetic transcranial stimulation. *Electroencephalogr Clin Neurophysiol* 105: 438–450.
16. Rossini PM, Rossi S, Tecchio F, Pasqualetti P, Finazzi-Agro A, et al. (1996) Focal brain stimulation in healthy humans: motor maps changes following partial hand sensory deprivation. *Neurosci Lett* 214: 191–195.
17. Rossini PM, Caltagirone C, Castriota-Scanderbeg A, Cicinelli P, Del Gratta C, et al. (1998) Hand motor cortical area reorganization in stroke: a study with fMRI, MEG and TCS maps. *Neuroreport* 9: 2141–2146.
18. Rossini PM, Pauri F, Cicinelli P, Pasqualetti P, Traversa R, et al. (1999) Neuromagnetic recordings and magnetic brain stimulation in the evaluation of sensorimotor hand area interhemispheric differences: normative, experimental and patients’ data. *Electroencephalogr Clin Neurophysiol Suppl* 50: 210–220.
19. Rossini PM, Rossi S, Pasqualetti P, Tecchio F (1999) Corticospinal excitability modulation to hand muscles during movement imagery. *Cereb Cortex* 9: 161–167.
20. Rossini PM, Berardelli A, Deuschl G, Hallett M, Maertens de Noordhout AM, et al. (1999) Applications of magnetic cortical stimulation. The International Federation of Clinical Neurophysiology. *Electroencephalogr Clin Neurophysiol Suppl* 52: 171–185.
21. Ferreri F, Pauri F, Pasqualetti P, Fini R, Dal Forno G, et al. (2003) Motor cortex excitability in Alzheimer’s disease: a transcranial magnetic stimulation study. *Ann Neurol* 53: 102–108.
22. Wilson SA, Thickbroom GW, Mastaglia FL (1993) Transcranial magnetic stimulation of the motor cortex in normal subjects. The representation of two intrinsic hand muscles. *Journal of the Neurological Science* 118: 134–144.
23. Krings T, Naujokat C, von Keyserlingk DG (1998) Representation of cortical motor function as revealed by stereotactic transcranial magnetic stimulation. *Electroencephalogr Clin Neurophysiol* 109: 85–93.
24. Kwan HC, MacKay WA, Murphy JT, Wong YC (1978) Spatial organization of precentral cortex in awake primates. II. Motor outputs. *J Neurophysiol* 41: 1120–1131.
25. Sessle BJ, Wiesendanger M (1982) Structural and functional definition of the motor cortex in the monkey (*Macaca fascicularis*). *J Physiol* 323: 245–265.
26. Gould HJ, Cusick CG, Pons TP, Kaas JH (1986) The relationship of corpus callosum connections to electrical stimulation maps of motor, supplementary motor, and the frontal eye fields in owl monkeys. *J Comp Neurol* 247: 297–325.
27. Donoghue JP, Leibovic S, Sanes JN (1992) Organization of the forelimb area in squirrel monkey motor cortex: representation of digit, wrist, and elbow muscles. *Exp Brain Res* 89: 1–19.
28. Nudo RJ, Jenkins WM, Merzenich MM, Prejean T, Grenda R (1992) Neurophysiological correlates of hand preference in primary motor cortex of adult squirrel monkeys. *J Neurosci* 12: 2918–2947.
29. Buonomano DV, Merzenich MM (1998) Cortical plasticity: from synapses to maps. *Annu Rev Neurosci* 21: 149–86.
30. Rossini PM, Calautti C, Pauri F, Baron JC (2003) Post-stroke plastic reorganization in the adult brain. *Lancet Neurol* 2: 493–502.
31. Rossini PM, Dal Forno G (2004) Neuronal post-stroke plasticity in the adult. *Restor Neurol Neurosci* 22: 193–206.
32. Rossini PM, Dal Forno G (2004) Integrated technology for evaluation of brain function and neural plasticity. *Phys Med Rehabil Clin N Am* 15: 263–306.
33. Schieber MH (2001) Constraints on somatotopic organization in the primary motor cortex. *J Neurophysiol* 86: 2125–2143.
34. Sanes JN, Schieber MH (2001) Orderly somatotopy in primary motor cortex: does it exist? *Neuroimage* 13: 968–974.
35. Sanes JN, Donoghue JP (1997) Static and dynamic organization of motor cortex. *Adv Neurol* 73: 277–296.
36. Sanes JN, Donoghue JP (2000) Plasticity and primary motor cortex. *Annu Rev Neurosci* 23: 393–415.
37. Sanes JN, Donoghue JP, Thangaraj V, Edelman RR, Warach S (1995) Shared neural substrates controlling hand movements in human motor cortex. *Science* 268: 1775–1777.
38. Schieber MH, Hibbard LS (1993) How somatotopic is the motor cortex hand area? *Science* 261: 489–492.
39. Penfield W, Boldrey E (1937) Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain* 37: 389–443.
40. Strick PL, Preston JB (1982) Two representations of the hand in area 4 of a primate. I. Motor output organization. *J Neurophysiol* 48: 139–149.
41. Humphrey DR (1986) Representation of movements and muscles within the primate precentral motor cortex: historical and current perspectives. *Federation Proc* 45: 2687–2699.
42. Sato KC, Tanji J (1989) Digit-muscle responses evoked from multiple intracortical foci in monkey precentral motor cortex. *J Neurophysiol* 62: 959–970.
43. Huntley GW, Jones EG (1991) Relationship of intrinsic connections to forelimb movement representations in monkey motor cortex: a correlative anatomic and physiological study. *J Neurophysiol* 66: 390–413.
44. Rossini PM, Pauri F (2000) Neuro magnetic integrated methods tracking human brain mechanism of sensorymotor areas “plastic” reorganisation. *Brain Res Rev* 33: 131–154.
45. Andersen P, Hagan PJ, Phillips CG, Powell TP (1975) Mapping by microstimulation of overlapping projections from area 4 to motor units of the baboon’s hand. *Proc R Soc Lond B Biol Sci* 188: 31–36.
46. Lemon RN, Muir RB, Mantel GW (1987) The effects upon the activity of hand and forearm muscles of intracortical stimulation in the vicinity of corticomotor neurones in the conscious monkey. *Exp Brain Res* 66: 621–637.
47. Kalaska JF, Crammond DJ (1992) Cerebral cortical mechanisms of reaching movements. *Science* 255: 1517–1523.
48. Korvenoja A, Huttunen J, Salli E, Pohjonen H, Martinkauppi S, et al. (1999) Activation of multiple cortical areas in response to somatosensory stimulation: combined magnetoencephalographic and functional magnetic resonance imaging. *Hum Brain Mapp* 8: 13–27.
49. Frostig RD, Lieke EE, Ts’o DY, Grinvald A (1990) Cortical functional architecture and local coupling between neuronal activity and the microcirculation revealed by in vivo high-resolution optical imaging of intrinsic signals. *Proc Natl Acad Sci U S A* 87: 6082–6086.
50. Berardelli A (1991) Electrical and magnetic spinal and cortical stimulation in man. *Curr Opin Neurol Neurosurg* 4: 770–776.
51. Berardelli A, Inghilleri M, Polidori L, Priori A, Mercuri B, et al. (1994) Effects of transcranial magnetic stimulation on single and sequential arm movements. *Exp Brain Res* 98: 501–506.
52. Gregori B, Curra A, Dinapoli L, Bologna M, Accornero N, et al. (2005) The timing and intensity of transcranial magnetic stimulation, and the scalp site stimulated, as variables influencing motor sequence performance in healthy subjects. *Exp Brain Res* 166: 43–55.
53. Rossini PM, Caramia MD, Iani C, Desiato MT, Sciarretta G, et al. (1995) Magnetic transcranial stimulation in healthy humans: influence on the behavior of upper limb motor units. *Brain Res* 676: 314–324.
54. Mills KR (1999) Magnetic brain stimulation: a review after 10 years experience. *Electroencephalogr Clin Neurophysiol Suppl* 49: 239–244.
55. Nithi KA, Mills KR (2000) Mapping motor cortex projections to single motor units in humans with transcranial magnetic stimulation. *Muscle Nerve* 23: 1542–1548.
56. Arunachalam R, Weerasinghe VS, Mills KR (2005) Motor control of rapid sequential finger tapping in humans. *J Neurophysiol* 94: 2162–2170.
57. Cohen LG, Bandinelli S, Topka HR, Fuhr P, Roth BJ, et al. (1991) Topographic maps of human motor cortex in normal and pathological conditions: mirror movements, amputations and spinal cord injuries. *Electroencephalogr Clin Neurophysiol Suppl* 43: 36–50.
58. Cohen LG, Hallett M (1988) Noninvasive mapping of human motor cortex. *Neurology* 38: 904–909.
59. Hallett M (2000) Transcranial magnetic stimulation and the human brain. *Nature* 406: 147–150.
60. Jabre JF, Hallett M, Lemon R, Kernell D (2000) Recent developments in our understanding of motor control, corticomotoneuronal connections and the activation of motoneurons and motoneuronal pools. *Suppl Clin Neurophysiol* 53: 171–177.
61. Ziemann U, Netz J, Szelenyi A, Homberg V (1993) Spinal and supraspinal mechanisms contribute to the silent period in the contracting soleus muscle after transcranial magnetic stimulation of human motor cortex. *Neurosci Lett* 156: 167–171.

62. Ziemann U, Hallett M (2001) Hemispheric asymmetry of ipsilateral motor cortex activation during unimanual motor tasks: further evidence for motor dominance. *Clin Neurophysiol* 112: 107–113.
63. Pascual-Leone A, Tarazona F, Catala MD (1999) Applications of transcranial magnetic stimulation in studies on motor learning. *Electroencephalogr Clin Neurophysiol Suppl* 51: 157–161.
64. Di Lazzaro V, Oliviero A, Pilato F, Saturno E, Di Leone M, et al. (2004) The physiological basis of transcranial motor cortex stimulation in conscious humans. *Clin Neurophysiol* 115: 255–266.
65. Di Lazzaro V, Oliviero A, Proficue P, Insola A, Mazzone P, et al. (1999) Direct demonstration of interhemispheric inhibition of the human motor cortex produced by transcranial magnetic stimulation. *Exp Brain Res* 124: 520–524.
66. Cicinelli P, Marconi B, Zaccagnini M, Pasqualetti P, Filippi MM, et al. (2005) Imagery-induced Cortical Excitability Changes in Stroke: A Transcranial Magnetic Stimulation Study. *Cereb Cortex*.
67. Cicinelli P, Pasqualetti P, Zaccagnini M, Traversa R, Olivetti M, et al. (2003) Interhemispheric asymmetries of motor cortex excitability in the postacute stroke stage: a paired-pulse transcranial magnetic stimulation study. *Stroke* 34: 2653–2658.
68. Cicinelli P, Mattia D, Spanedda F, Traversa R, Marciani MG, et al. (2000) Transcranial magnetic stimulation reveals an interhemispheric asymmetry of cortical inhibition in focal epilepsy. *Neuroreport* 11: 701–707.
69. Liepert J, Hamzei F, Weiller C (2004) Lesion-induced and training-induced brain reorganization. *Restor Neurol Neurosci* 22: 269–277.
70. Liepert J, Weiller C (1999) Mapping plastic brain changes after acute lesions. *Curr Opin Neurol* 12: 709–713.
71. Cantello R (2002) Applications of transcranial magnetic stimulation in movement disorders. *J Clin Neurophysiol* 19: 272–293.
72. Krings T, Chiappa KH, Foltys H, Reinges MH, Cosgrove GR, et al. (2001) Introducing navigated transcranial magnetic stimulation as a refined brain mapping methodology. *Neurosurg Rev* 24: 171–179.
73. Krings T, Reinges MH, Foltys H, Cosgrove GR, Thron A (2001) Multimodality neuroimaging: research and clinical applications. *Neurol Clin Neurophysiol* 1: 2–11.
74. Foltys H, Krings T, Meister IG, Sparing R, Boroojerdi B, et al. (2003) Motor representation in patients rapidly recovering after stroke: a functional magnetic resonance imaging and transcranial magnetic stimulation study. *Clin Neurophysiol* 114: 2404–2415.
75. Lotze M, Kaethner RJ, Erb M, Cohen LG, Grodd W, et al. (2003) Comparison of representational maps using functional magnetic resonance imaging and transcranial magnetic stimulation. *Clin Neurophysiol* 114: 306–312.
76. Neggers SF, Langerak TR, Schutter DJ, Mandl RC, Ramsey NF, et al. (2004) A stereotactic method for image-guided transcranial magnetic stimulation validated with fMRI and motor-evoked potentials. *Neuroimage* 21: 1805–1817.
77. Bohning DE, Denslow S, Bohning PA, Walker JA, George MS (2003) A TMS coil positioning/holding system for MR image-guided TMS interleaved with fMRI. *Clin Neurophysiol* 114: 2210–2219.
78. Gugino LD, Romero JR, Aglio L, Titone D, Ramirez M, et al. (2001) Transcranial magnetic stimulation coregistered with MRI: a comparison of a guided versus blind stimulation technique and its effect on evoked compound muscle action potentials. *Clin Neurophysiol* 112: 1781–1792.
79. Capaday C (2004) The integrated nature of motor cortical function. *Neuroscientist* 10: 207–220.
80. Devanne H, Cohen LG, Kouchtir-Devanne N, Capaday C (2002) Integrated motor cortical control of task-related muscles during pointing in humans. *J Neurophysiol* 87: 3006–3017.
81. Cheung VC, d'Avella A, Tresch MC, Bizzi E (2005) Central and sensory contributions to the activation and organization of muscle synergies during natural motor behaviors. *J Neurosci* 25: 6419–6434.
82. Rossini PM, Barker AT, Berardelli A, Caramia MD, Caruso G, et al. (1994) Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic principles and procedures for routine clinical application. Report of an IFCN committee. *Electroencephalogr Clin Neurophysiol* 91: 79–92.
83. Nielsen JF (1996) Logarithmic distribution of amplitudes of compound muscle action potentials evoked by transcranial magnetic stimulation. *J Clin Neurophysiol* 13: 423–434.
84. Wassermann EM, Wang B, Zeffiro TA, Sadato N, Pascual-Leone A, et al. (1996) Locating the motor cortex on the MRI with transcranial magnetic stimulation and PET. *Neuroimage* 3: 1–9.