

# Platelet-Derived Growth Factor Induces *Rad* Expression through Egr-1 in Vascular Smooth Muscle Cells

Yan Luo<sup>1</sup>, Meiling Zhang<sup>1</sup>, Ji Zhang<sup>1</sup>, Jifeng Zhang<sup>2</sup>, Chunlei Chen<sup>1</sup>, Y. Eugene Chen<sup>2</sup>, Jing-Wei Xiong<sup>1</sup>, Xiaojun Zhu<sup>1\*</sup>

**1** The Institute of Molecular Medicine, Peking University, Beijing, China, **2** The Cardiovascular Center, University of Michigan, Ann Arbor, Michigan, United States of America

## Abstract

**Background:** Ras associated with diabetes (*Rad*) inhibits vascular lesion formation by reducing the attachment and migration of vascular smooth muscle cells (VSMCs). However, the transcriptional regulation of *Rad* in VSMCs is unclear.

**Methodology and Principal Findings:** We found that Platelet-Derived Growth Factor (PDGF) induced *Rad* expression in a time- and dose-dependent manner in rat aortic smooth muscle cells (RASMCs) using quantitative real-time PCR. By serial deletion analysis of the *Rad* promoter, we identified that two GC-rich early growth response-1 (Egr-1) binding sites are essential for PDGF-induced *Rad* promoter activation. Overexpression of Egr-1 in RASMCs strongly stimulated *Rad* expression while the Egr-1 corepressor, NGFI-A binding protein 2 (NAB2), repressed PDGF-induced *Rad* up-regulation in a dose-dependent manner. Direct binding of Egr-1 to the *Rad* promoter region was further confirmed by chromatin immunoprecipitation assays.

**Conclusions:** Our results demonstrate that *Rad* is regulated by PDGF through the transcriptional factor Egr-1 in RASMCs.

**Citation:** Luo Y, Zhang M, Zhang J, Zhang J, Chen C, et al. (2011) Platelet-Derived Growth Factor Induces *Rad* Expression through Egr-1 in Vascular Smooth Muscle Cells. PLoS ONE 6(4): e19408. doi:10.1371/journal.pone.0019408

**Editor:** Carlo Gaetano, Istituto Dermopatico dell'Immacolata, Italy

**Received:** October 18, 2010; **Accepted:** April 5, 2011; **Published:** April 29, 2011

**Copyright:** © 2011 Luo et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by Natural Science Foundation of China [30400223 and 30671027 to X.Z.]; Peking University 985 Project; National Basic Research Program of China [2007CB512100 and 2010CB529503]; and American Heart Association National Career Development Grant [0835237N to J.Z.] The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: zhuxiaojun@pku.edu.cn

## Introduction

Ras associated with diabetes (*Rad*) is a member of the RGK family which is composed of *Rad*, *Gem/Kir*, *Rem* and *Rem2* [1]. It is expressed in the heart, skeletal muscle and lung [2]. *Rad* is highly expressed in the skeletal muscle of some type II diabetic patients [2], which suggests that *Rad* is related with glucose metabolism and insulin resistance. Our previous studies demonstrate that *Rad* is critical in maintaining normal cardiac functions. *Rad* expression decreases significantly in human failing hearts and *Rad* knockout (KO) mice are more susceptible to cardiac hypertrophy with increased CaMKII phosphorylation compared with their littermate controls [3]. Our findings as well as the others' indicate that *Rad* inhibits myocardium L-type calcium channel activity and attenuates the  $\beta$ -Adrenergic Receptor ( $\beta$ -AR) activity [4,5]. Dominant negative suppression of endogenous *Rad* in the heart up-regulate L-type  $Ca^{2+}$  channel expression on the plasma membrane, leading to  $I_{Ca,L}$  increase and action potential prolongation [6]. *Rad* is upregulated in vascular smooth muscle cells (VSMCs) during the formation of vascular lesions and overexpression of *Rad* attenuated neointimal formation by strongly inhibiting smooth muscle migration [7]. However, the molecular mechanism for the induction of *Rad* during vascular lesion formation is unknown.

Platelet-derived growth factor (PDGF) plays an important role in normal tissue growth and the patho-physiological processes of

vascular diseases like atherosclerosis and restenosis [8]. During the initiation and progression of atherosclerosis, VSMCs are activated by growth factors like PDGF or cytokines, then proliferate and migrate from the media into the intimal surface of the vessel, thus facilitating neointimal formation [8]. Egr-1 is a zinc-finger transcription factor that regulates cell proliferation and differentiation [9]. It is an immediate-early response protein that is rapidly and transiently stimulated by various growth factors including PDGF [10]. Egr-1 regulates gene transcription by the specific binding of its DNA binding domain, which consists of three zinc fingers, to the consensus GC-rich regions in the promoter of its target genes [11]. Structure analysis of Egr-1 identified a 34 amino acids inhibitory domain (R1) at the 5' zinc finger binding region [12]. Two corepressors, NGFI-A-binding proteins 1 and 2 (NAB1 and NAB2) can markedly decrease Egr-1 transcriptional activity by binding to this domain [13,14].

In the present study, we set out to explore how *Rad* is transcriptionally regulated in VSMCs. We found PDGF induced *Rad* expression in a dose- and time-dependent manner, which Egr-1 and its partners mediated this induction.

## Results

### PDGF induces *Rad* expression in RASMCs

To determine the effects of the growth factor PDGF on *Rad* expression in RASMCs, we treated cultured RASMCs with PDGF

(20 ng/ml) for 0, 0.5, 1, 2 and 6 hours. Quantitative real-time PCR revealed that expression of Rad increased 0.5 hours after PDGF stimulation, peaked at 1 hour, and returned to the baseline levels 6 hours later (Figure 1A). Rad was induced by PDGF in a dose-dependent manner (Figure 1B). One hour of 20 ng/ml PDGF treatment resulted in a 3.5-fold increase in Rad mRNA, compared to untreated cells (Figure 1B).

To identify the mechanisms by which PDGF activates Rad expression, we isolated Rad promoter and made reporter constructs containing different length Rad promoters. RASMCs were transfected with these constructs and then treated with PDGF. PDGF stimulated Rad promoter activity in all Rad promoter constructs except pRad-57, which also lacked basal promoter activity (Figure 1C).


### Egr-1 binding sites in the Rad promoter are required for PDGF-induced Rad promoter activation

Computer analysis revealed two GC-rich regions that may serve as Egr-1 binding sites within the -155 to -57 bp region of the Rad promoter (Figure 2A). To investigate whether these are functional Egr-1 binding sites, we over-expressed a constitutively active Egr-1 (Egr-1\*) and tested whether it could transactivate Rad promoter. Egr-1\* over-expression resulted in a ~25-fold increases of the pRad-155bp promoter activity compared with pcDNA3.1 vector transfection (Figure 2B). Then we mutated one or both of the predicted Egr-1 binding sites in the pRad-155 construct and co-transfected wild-type or mutant reporter constructs with a constitutive active Egr-1\* expression construct into 293A cells. Disruption of the first Egr-1 binding site in position -74 (pRad-155m1) caused a slight decrease in the Rad promoter activity but the induction upon Egr-1\* overexpression remain unchanged (Figure 2B). However, disruption of the second Egr-1 binding site in position -62 (pRad-155m2) resulted in a weaker induction of the Rad promoter activity by Egr-1\* (~7 folds) compared with the 25-fold activation in the pRad-155 construct, whereas the basal level remained unchanged. However, mutations on both Egr-1 binding sites (pRad-155m3) impaired both basal and Egr-1\*-induced transactivation of the Rad promoter (Figure 2B).

To further test whether Egr-1 binding site mutation affects PDGF-induced Rad promoter activation, we transfected RASMCs with wild-type (pRad-155) or Egr-1 binding site mutant constructs and then treated the cells with PDGF. PDGF triggered a nearly 3 fold promoter activity induction on pRad-155, and the induction dropped to ~1.8 fold in pRad-155m1, and no PDGF induction could be observed in pRad-155m2 construct meanwhile the basal activity of these constructs remained consistent. When both Egr-1 binding sites were disrupted as in pRad-155m3, the basal level of the Rad promoter activity dropped dramatically and no induction by PDGF was found (Figure 2C). These results are consistent with the above experiments performed in 293A cells overexpressing a constitutive active Egr-1\*. Furthermore, mutation of both Egr-1 binding sites in the pRad-3050 reporter construct caused ~ 54% decrease in the basal level of pRad-3050 promoter activity and PDGF-induced promoter activation was also completely abolished in pRad-3050m (Figure 2D). Taken together, the Egr-1-responsive regions located at -62 and -74 bp of Rad promoter are essential for PDGF-induced Rad promoter activation.


### Egr-1 mediates PDGF-induced Rad expression in RASMCs

Egr-1 was activated by various growth factors including PDGF, and we also found a remarkable induction of Egr-1 by PDGF in RASMCs (data not shown). Real-time quantitative PCR and Western blot demonstrated that Egr-1 markedly induced Rad expression at both mRNA and protein levels (Figure 3), which is



**Figure 1. PDGF induces Rad expression in RASMCs.** (A) Time course of Rad expression level in RASMCs after 20 ng/ml PDGF-BB treatment. (B) Dose-response of Rad expression after 1 hour treatment with the indicated concentrations of PDGF-BB. Rad expression was detected by real-time PCR and normalized by 18S (n=3, \*  $p < 0.05$ , \*\*  $p < 0.01$ ). (C) Responses of different Rad promoter reporters to PDGF-BB stimulation in RASMCs (n=4, \*\*  $p < 0.01$ ). doi:10.1371/journal.pone.0019408.g001

consistent with Egr-1 transactivation of the Rad promoter. These results suggest that Egr-1 is a potent transcriptional activator for Rad. We hypothesized that PDGF induced Rad expression via the activation of Egr-1. NAB2 functions as a corepressor of Egr-1 [14],




**Figure 2. Egr-1 binding sites are essential for Rad promoter activation.** (A) Nucleotide sequence between base pairs -76 and -51bp of Rad promoter. Egr-1 binding sites are boxed. The mutated bases are indicated in boldface. (B) Wild-type and mutant forms of pRad-155 were cotransfected with pcDNA3.1-Egr-1\* or pcDNA3.1 into 293A cells and luciferase activity were determined. The fold induction of Luciferase activity by Egr-1 was shown in the lower panel; (C) RASMCs were transfected with pRad-155 promoter reporter or its mutant forms then treated with PDGF and luciferase activity were determined (n=4, \*\*p<0.01). (D) pRad-3050 and pRad-3050m were transfected into RASMCs then treated with PDGF and luciferase activity were determined (n=4, \*\*p<0.01). doi:10.1371/journal.pone.0019408.g002

therefore we asked whether over-expression of NAB2 inhibits PDGF-induced Rad expression. RASMCs were infected with a recombinant adenovirus expressing NAB2 (Ad-NAB2) followed by PDGF treatment for 1 hour. Ad-GFP was used as a control in this experiment. We found that Rad expression increased in response to PDGF in Ad-GFP-infected RASMCs, and overexpression of NAB2 resulted in a dose-dependent abrogation of PDGF-induced Rad expression (Figure 4A). The induction of Rad mRNA was completely abolished when Ad-NAB2 reached a concentration of 250 MOI (multiplicity of infection). Furthermore, NAB2 inhibited Rad promoter activity elevated by PDGF-BB (Figure 4B). These results support that Egr-1 is a key mediator involved in PDGF-induced Rad expression in RASMCs.

### PDGF enhances Egr-1 binding to Rad promoter

ChIP assay was performed to further confirm the physiological relevance and functionality of Egr-1 through its putative binding sites in the Rad promoter. RASMCs untreated or treated with PDGF for 1 hour were incubated with formaldehyde to cross-link protein and binding sites in DNA. The -213 to -2 region of the Rad promoter region was amplified by PCR. After PDGF treatment, Egr-1 was found to bind to the Rad promoter. As for DNA from untreated cells or DNA precipitated by control IgG, we did not find any PCR amplifications (Figure 5). Our data strongly support that Egr-1 bound to the proximal Rad promoter following PDGF treatment in RASMCs.




**Figure 3. Egr-1 induces Rad expression.** (A) RASMCs were infected with Ad-Egr-1 or Ad-GFP for 24 hours. Rad mRNA levels were detected by real-time PCR (n=3, \*\*p<0.01 vs. control); (B) Rad protein levels were determined by Western blotting analysis. Expression of  $\beta$ -actin was detected as the control for equal loading. doi:10.1371/journal.pone.0019408.g003

**Discussion**


In the present study, we provide the first evidence that PDGF induces Rad expression in a time- and dose-dependent manner in rat aortic smooth muscle cells, and that Egr-1 is a key transcriptional factor to mediate PDGF-induced Rad expression.

Rad is a RSK-family small GTPase initially identified by subtractive cloning and found to be over-expressed in skeletal muscle of a group of patients with type II diabetes [2]. Rad possesses a structurally unique Ras-related core and COOH- and NH<sub>2</sub>-terminal extensions but lacks the CAAX-like prenylation motif at the COOH terminus possessed by Ras [15,16]. In atherosclerotic lesions, PDGF released from inflammatory and immune cells promotes VSMCs proliferation and attracts VSMCs to migrate from media to intima [17]. In different animal models of acute arterial injury, VSMCs accumulation in lesions is inhibited by the administration of various PDGF pathway inhibitors, including neutralizing PDGF antibodies [18], PDGFR kinase inhibitors [19], and PDGFR-neutralizing antibodies [20]. Here we demonstrated that PDGF stimulated Rad expression in RASMCs in a time- and dose-dependent manner. Our previous study indicated that Rad is a critical mediator that reduces vascular lesion formation by suppressing the attachment and migration of VSMCs via inhibition of Rho/ROK activity [7]. Rad induced by PDGF may serve as a suppressor for PDGF induced smooth muscle cell migration.

We have identified that normal Rad level is critical in maintaining cardiac and blood vessel functions. Rad expression decreases in human failing hearts and Rad knockout (KO) mice



**Figure 4. Egr-1 is an essential mediator for PDGF induced Rad expression.** (A) RASMCs were infected with Ad-NAB2 at MOI of 0, 50,100,150 and 250 before treated with PDGF-BB (20 ng/ml) for 1 hour. Ad-GFP was added to keep the amount of adenovirus consistent. Rad mRNA was quantified using real-time PCR (n = 3, \*p<0.05, \*\*p<0.01); (B) RASMCs were transfected with pRad-155 and infected with Ad-NAB2 or Ad-GFP for 24 hours and then stimulated with PDGF. Luciferase activity was measured 1 hour after PDGF stimulation (n = 4, \*p<0.05, \*\*p<0.01). doi:10.1371/journal.pone.0019408.g004



**Figure 5. Egr-1 binds to Rad promoter after PDGF treatment.** ChIP assay was performed in RASMC stimulated with or without PDGF (20 ng/ml) for 1 hour. After formaldehyde cross-linking, the protein-DNA complexes were recovered using anti-Egr-1 antibody or non-specific IgG. PCR was performed to detect the proximal Rad promoter. PCR products were detected by 2% agarose gel electrophoresis. doi:10.1371/journal.pone.0019408.g005

are more susceptible to cardiac hypertrophy [3]. Rad inhibits myocardium L-type calcium channel activity and attenuates the  $\beta$ -Adrenergic Receptor ( $\beta$ -AR) activity [4,5]. In blood vessel, Rad expression increases significantly after balloon injury. Unlike other small GTPase, the RGK family proteins are regulated at the transcriptional level. However, the understanding for Rad transcriptional regulation is very limited. Potential E box sequences (CANNTG), which could serve as binding sites for the HLH family transcription factors like myf5, MyoD, myogenin and MRF4, are predicted in the Rad proximal promoter region [21]. Rad expression increases significantly during skeletal muscle regeneration. Myogenic transcriptional factors like MEF2, MyoD and Myf-5 can increase the transcriptional activity of Rad promoter [22].

Until now, the molecular mechanism by which Rad is activated during vascular lesions formation is not clear. Rad promoter activity increased after PDGF treatment, suggesting that PDGF induces Rad expression at transcriptional level. Deletion and mutation analysis of Rad promoter revealed that potential Egr-1 binding sites would be critical for PDGF-induced Rad expression. It has been well documented by us and other groups that PDGF induces dramatic expression of Egr-1 [23]. Our data showed that Egr-1 activated the Rad promoter in RASMCs, and overexpression of Egr-1 resulted in a marked increase at Rad mRNA and protein levels in RASMCs. Egr-1 activity is negatively regulated by NGFI-A-binding proteins 1 and 2 (NAB 1 and NAB 2). Binding of NAB2 to Egr-1 through interaction between the NCD1 (NAB conserved domain 1, NCD1) and the R1 domain of Egr-1 represses the transcriptional activity mediated by Egr-1 [14]. We found that up-regulation of Rad induced by PDGF-BB was inhibited by the co-repressor NAB2 in a dose-dependent manner. Together, these data support that Egr-1 and its interacting partner(s) are the key mediators responsible for PDGF-induced Rad expression.

Egr-1 expression is strikingly elevated in the VSMCs of atherosclerotic lesions [24] and plays critical roles in regulating VSMCs growth and intimal thickening after vascular injury [25]. Egr-1 binds preferentially to GC-rich regions of the promoters of its target genes [26]. CHIP assay indicated that after PDGF stimulation, Egr-1 binded to the proximal Rad promoter region. Serial deletions of the Rad promoter also defined a -155 to -57 bp region responsible for both basal and Egr-1-inducible promoter activity. Two GC-rich motifs were predicted in this region as potential binding sites for Egr-1. Mutation of both Egr-1 binding sites in pRad-3050 and pRad-155 resulted in dramatic reduction in Rad basal promoter activity and complete abolishment of PDGF-induced Rad promoter activation. Egr1-activated Rad promoter activity was attenuated when the putative Egr-1 binding site at position -62 or both binding sites were mutated. Consistent with this, PDGF-stimulated Rad promoter activation was abolished when the putative Egr-1 binding site in position -62 or both binding sites were mutated. These results suggest that the Egr-1 response element located at -62 bp plays the crucial role in PDGF-induced Rad promoter activity.

Here we identified Rad as another target gene of transcriptional factor Egr-1. After vascular cell injury, Egr-1 is expressed primarily in the nucleus and is capable of altering the transcription of several genes implicated in the pathogenesis of vascular disease, including PDGF, FGF-2, TNF $\alpha$ , tissue factor, ICAM and p53. Genes activated by Egr-1 play important roles in VSMC proliferation, neointima formation and contribute to the development of vascular diseases. Many of those genes like PDGF are activated by Egr-1 and further induce Egr-1 expression. Osteopontin and Egr-1 also positively regulate each other in

VSMCs, which may play an important role in controlling inappropriate remodeling of vessel walls [27]. These positive feedback loops amplify gene transcription activated by Egr-1. On the other hand, the negative feedback loop between Egr-1 and other genes like NAB2 prevents the permanent activation of Egr-1 target genes. In response to extracellular stimuli Egr-1 induces the expression of NAB2, which in turn represses the activity of Egr-1 through binding to the R1 domain of Egr-1 [28]. Gene expression profiling by microarray analyses revealed enhanced expression of several Egr family members in the hearts of Rad Knockout mice compared with their wild-type littermates. These raised the possibility that there is a negative feedback loop between Rad and Egr-1, i.e. Egr-1 activation induced Rad expression and Rad inhibited the further activation of Egr-1. Furthermore, the effect of Rad on other transcription factors involved in neointima formation and VSMC migration remains to be further investigated.

## Materials and Methods

### Cell Culture

Rat aortic smooth muscle cells (RASMCs) were isolated from male Sprague-Dawley rats as described previously [23] and cultured in Dulbecco's modified Eagle's medium supplemented with 10% FBS in a 5% CO<sub>2</sub> humidified atmosphere at 37°C. Early passages (5 to 9) of cells grown to 80-90% confluence were used for all experiments. Cells were placed in serum-free medium for 24 hours before treatment with PDGF-BB (Sigma). This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Experiments were approved by the Peking University Animal Care and Use Committee (Permit Number: IMM-XiongJW-2).

### Plasmids and adenoviral recombinants

The pcDNA3.1-Egr-1\* plasmid containing constitutively active Egr-1 as well as the adenoviruses Ad-Egr-1 and Ad-NAB2 were prepared as described previously [29]. Fragments of the Rad promoter region (3050 bp, 1040 bp, 155 bp and 57 bp) were amplified by PCR and subcloned into the pGL3-basic vector to generate the pRad-3050, pRad-1040, pRad-155 and pRad-57 reporter constructs. These constructs were confirmed by DNA sequencing.

### RNA Isolation and Quantitative Real-time PCR

Total RNA was extracted from cultured RASMCs with Trizol (Invitrogen) and 1  $\mu$ g was used for cDNA synthesis using a first strand cDNA synthesis kit according to the manufacturer's instructions (Invitrogen). Rad mRNA level was assessed by real-time quantitative RT-PCR as previously described [7] and 18S level was measured for normalization of variations in RNA input and cDNA synthesis.

### Western Blot Analyses

Cells were lysed with lysis buffer (Cell Signaling Technology). The lysates were resolved on 12% SDS-polyacrylamide gels, and proteins were then transferred to PVDF membranes (Bio-Rad). The membranes were blocked for 1 hour at room temperature and incubated overnight at 4°C with anti-Rad antibody (1:2000). Rabbit anti-Rad polyclonal antibody was a kind gift from Dr. Ronald Kahn [30]. Blots were then incubated with anti-rabbit secondary antibody (1:2000) for 1 hour at room temperature and the Rad/antibody complexes were visualized by chemiluminescence (Bio-Rad) according to the manufacturer's recommendations.

## Transient Transfection and Luciferase Assays

Transfection of RASMCs or 293A cells were performed using Lipofectamine<sup>TM</sup>2000 (Invitrogen) according to the manufacturer's instructions. The pRL-CMV plasmid was cotransfected as the control for transfection efficiency. Luciferase activity was measured by the Dual Luciferase Reporter System using a TD20/20 Luminometer (Turner Biosystems).

## Site-Directed Mutagenesis

Mutations were introduced in the pRad-155 or pRad-3050 plasmids using QuikChange II XL Site-Directed Mutagenesis Kit (Stratagene) according to the manufacturer's recommendations. The following oligonucleotides were used to create the desired mutations: Egr1-mut1 (for pRad-155), 5'-CCGCAGCGTG-GACCG**TTTT**CGGGAGCGGGGGCGG-3'; Egr1-mut2 (for pRad-155), 5'-GACCGGGGGCGGGAGCG**TTTT**CGGGTG-GAGGCTTAAAT-3'; and Egr1-mut3 (for pRad-155 and pRad-3050), 5'-GGACCG**TTTT**CGGGAGCG**TTTT**CGGGTGGA-GGCTTAAATA-3'. The mutated nucleotides are indicated in bold. Mutations were confirmed by DNA sequencing. The resulting pRad-155 mutants with one or both Egr1 binding sites mutated were designated pRad-155m1, pRad-155m2 and pRad-155m3. pRad-3050 with both Egr1 binding sites mutated was designated pRad-3050m.

## Chromatin Immunoprecipitation (ChIP)

ChIP assay was performed using a chromatin immunoprecipitation assay kit (Upstates). Cultured RASMCs were treated with PDGF or vehicle for 1 hour. Proteins were cross-linked with DNA by 1% formaldehyde treatment for 10 minutes at 37°C. Cells were then lysed and DNA was sheared by sonication (Microson<sup>TM</sup>,

strength 40%, pulse 10 seconds, 4 times). The sonicated DNA was diluted in ChIP dilution buffer. The diluted cell supernatant was pre-cleared with protein A agarose before immunoprecipitation with anti-Egr-1 antibody (Santa Cruz) or non-specific IgG. The immune complexes were collected by adsorption to protein A agarose, precipitated by gentle centrifugation at 3,000 rpm of 4°C for 1 min, and the supernatant containing unbound nonspecific DNA was discarded. The precipitate was washed and the immune complexes were eluted, adjusted to 200 mM NaCl and incubated at 65°C for 5 hours to reverse the crosslinks. After successive treatments with RNase A and proteinase K, the DNA was extracted with phenol-chloroform, precipitated with ethanol and resuspended in H<sub>2</sub>O. The immunoprecipitated DNA were analyzed by PCR with primers spanning -213 to -2 of the Rad promoter. The primer sequences were as follows: forward: 5'-TCGCTCTCTCTCTCCCTTCTCACAC-3' and reverse: 5'-AC-CCTCTTCCCTCGGACCTTACATC-3'. An aliquot of the sonicated DNA was used as the input. PCR products were detected by 2% agarose gel electrophoresis.

## Statistical Analysis

Each experiment was repeated for a minimum of three times. Statistics were analyzed using either ANOVA (for multiple comparisons) or Student's 2-tailed t test for comparing two means. Data are presented as Means ± SD. In all cases p < 0.05 was considered statistically significant.

## Author Contributions

Conceived and designed the experiments: XZ. Performed the experiments: YL MZ Ji Zhang CC. Analyzed the data: YEC J-WX. Contributed reagents/materials/analysis tools: Jifeng Zhang. Wrote the paper: YL XZ.

## References

- Kelly K (2005) The RGK family: a regulatory tail of small GTP-binding proteins. *Trends in Cell Biology* 15: 640–643.
- Reynet C, Kahn CR (1993) Rad: a member of the Ras family overexpressed in muscle of type II diabetic humans. *Science* 262: 1441–1444.
- Chang L, Zhang J, Tseng Y-H, Xie C-Q, Ilany J, et al. (2007) Rad GTPase Deficiency Leads to Cardiac Hypertrophy. *Circulation* 116: 2976–2983.
- Finlin BS, Crump SM, Satin J, Andres DA (2003) Regulation of voltage-gated calcium channel activity by the Rem and Rad GTPases. *Proc Natl Acad Sci U S A* 100: 14469–14474.
- Wang G, Zhu X, Xie W, Han P, Li K, et al. (2010) Rad As a Novel Regulator of Excitation-Contraction Coupling and {beta}-Adrenergic Signaling in Heart. *Circ Res* 106: 317–327.
- Yada H, Murata M, Shimoda K, Yuasa S, Kawaguchi H, et al. (2007) Dominant negative suppression of Rad leads to QT prolongation and causes ventricular arrhythmias via modulation of L-type Ca<sup>2+</sup> channels in the heart. *Circ Res* 101: 69–77.
- Fu M, Zhang J, Tseng YH, Cui T, Zhu X, et al. (2005) Rad GTPase attenuates vascular lesion formation by inhibition of vascular smooth muscle cell migration. *Circulation* 111: 1071–1077.
- Heldin CH, Westermark B (1999) Mechanism of action and in vivo role of platelet-derived growth factor. *Physiol Rev* 79: 1283–1316.
- Silverman ES, Collins T (1999) Pathways of Egr-1-mediated gene transcription in vascular biology. *Am J Pathol* 154: 665–670.
- Fu M, Zhang J, Lin Y, Zhu X, Ehrenguber MU, et al. (2002) Early Growth Response Factor-1 Is a Critical Transcriptional Mediator of Peroxisome Proliferator-activated Receptor-γ1 Gene Expression in Human Aortic Smooth Muscle Cells. *Journal of Biological Chemistry* 277: 26808–26814.
- Gashler A, Sukhatme VP (1995) Early growth response protein 1 (Egr-1): prototype of a zinc-finger family of transcription factors. *Prog Nucleic Acid Res Mol Biol* 50: 191–224.
- Gashler AL, Swaminathan S, Sukhatme VP (1993) A novel repression module, an extensive activation domain, and a bipartite nuclear localization signal defined in the immediate-early transcription factor Egr-1. *Mol Cell Biol* 13: 4556–4571.
- Russo MW, Severson BR, Milbrandt J (1995) Identification of NAB1, a repressor of NGFI-A- and Krox20-mediated transcription. *Proceedings of the National Academy of Sciences of the United States of America* 92: 6873–6877.
- Svaren J, Severson BR, Apel ED, Zimonjic DB, Popescu NC, et al. (1996) NAB2, a corepressor of NGFI-A (Egr-1) and Krox20, is induced by proliferative and differentiative stimuli. *Mol Cell Biol* 16: 3545–3553.
- Kelly K (2005) The RGK family: a regulatory tail of small GTP-binding proteins. *Trends Cell Biol* 15: 640–643.
- Bilan PJ, Moyers JS, Kahn CR (1998) The ras-related protein rad associates with the cytoskeleton in a non-lipid-dependent manner. *Exp Cell Res* 242: 391–400.
- Andrae J, Gallini R, Betsholtz C (2008) Role of platelet-derived growth factors in physiology and medicine. *Genes Dev* 22: 1276–1312.
- Lewis CD, Olson NE, Raines EW, Reidy MA, Jackson CL (2001) Modulation of smooth muscle proliferation in rat carotid artery by platelet-derived mediators and fibroblast growth factor-2. *Platelets* 12: 352–358.
- Yamasaki Y, Miyoshi K, Oda N, Watanabe M, Miyake H, et al. (2001) Weekly dosing with the platelet-derived growth factor receptor tyrosine kinase inhibitor SU9518 significantly inhibits arterial stenosis. *Circ Res* 88: 630–636.
- Hart CE, Kraiss LW, Vergel S, Gilbertson D, Kenagy R, et al. (1999) PDGFbeta receptor blockade inhibits intimal hyperplasia in the baboon. *Circulation* 99: 564–569.
- Caldwell JS, Moyers JS, Doria A, Reynet C, Kahn RC (1996) Molecular cloning of the human rad gene: gene structure and complete nucleotide sequence. *Biochim Biophys Acta* 1316: 145–148.
- Hawke TJ, Kanatous SB, Martin CM, Goetsch SC, Garry DJ (2006) Rad is temporally regulated within myogenic progenitor cells during skeletal muscle regeneration. *Am J Physiol Cell Physiol* 290: C379–387.
- Kamimura M, Bea F, Akizawa T, Katus HA, Kreuzer J, et al. (2004) Platelet-derived growth factor induces tissue factor expression in vascular smooth muscle cells via activation of Egr-1. *Hypertension* 44: 944–951.
- McCaffrey TA, Fu C, Du B, Eksinar S, Kent KC, et al. (2000) High-level expression of Egr-1 and Egr-1-inducible genes in mouse and human atherosclerosis. *J Clin Invest* 105: 653–662.
- Harja E, Bucciarelli LG, Lu Y, Stern DM, Zou YS, et al. (2004) Early growth response-1 promotes atherosclerosis: mice deficient in early growth response-1 and apolipoprotein E display decreased atherosclerosis and vascular inflammation. *Circ Res* 94: 333–339.
- Paillard G, Deremble C, Lavery R (2004) Looking into DNA recognition: zinc finger binding specificity. *Nucleic Acids Res* 32: 6673–6682.

27. Liu QF, Yu HW, Liu GN (2009) Egr-1 upregulates OPN through direct binding to its promoter and OPN upregulates Egr-1 via the ERK pathway. *Mol Cell Biochem*.
28. Kumbrink J, Gerlinger M, Johnson JP (2005) Egr-1 induces the expression of its corepressor nab2 by activation of the nab2 promoter thereby establishing a negative feedback loop. *J Biol Chem* 280: 42785–42793.
29. Zhu X, Lin Y, Bacanamwo M, Chang L, Chai R, et al. (2007) Interleukin-1 beta-induced Id2 gene expression is mediated by Egr-1 in vascular smooth muscle cells. *Cardiovasc Res* 76: 141–148.
30. Zhu J, Bilan PJ, Moyers JS, Antonetti DA, Kahn CR (1996) Rad, a novel Ras-related GTPase, interacts with skeletal muscle beta-tropomyosin. *J Biol Chem* 271: 768–773.