

# Tau Phosphorylation and $\mu$ -Calpain Activation Mediate the Dexamethasone-Induced Inhibition on the Insulin-Stimulated Akt Phosphorylation

Yudong Liu<sup>1</sup>, Ying Su<sup>2</sup>, Shenggang Sun<sup>2</sup>, Tao Wang<sup>2</sup>, Xian Qiao<sup>2</sup>, Xiaoqin Run<sup>3</sup>, Zhihou Liang<sup>2\*</sup>

**1** Department of Orthopedics, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China, **2** Department of Neurology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China, **3** Department of General Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

## Abstract

Evidence has suggested that insulin resistance (IR) or high levels of glucocorticoids (GCs) may be linked with the pathogenesis and/or progression of Alzheimer's disease (AD). Although studies have shown that a high level of GCs results in IR, little is known about the molecular details that link GCs and IR in the context of AD. Abnormal phosphorylation of tau and activation of  $\mu$ -calpain are two key events in the pathology of AD. Importantly, these two events are also related with GCs and IR. We therefore speculate that tau phosphorylation and  $\mu$ -calpain activation may mediate the GCs-induced IR. Akt phosphorylation at Ser-473 (pAkt) is commonly used as a marker for assessing IR. We employed two cell lines, wild-type HEK293 cells and HEK293 cells stably expressing the longest human tau isoform (tau-441; HEK293/tau441 cells). We examined whether DEX, a synthetic GCs, induces tau phosphorylation and  $\mu$ -calpain activation. If so, we examined whether the DEX-induced tau phosphorylation and  $\mu$ -calpain activation mediate the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation. The results showed that DEX increased tau phosphorylation and induced tau-mediated  $\mu$ -calpain activation. Furthermore, pre-treatment with LiCl prevented the effects of DEX on tau phosphorylation and  $\mu$ -calpain activation. Finally, both LiCl pre-treatment and calpain inhibition prevented the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation. In conclusion, our study suggests that the tau phosphorylation and  $\mu$ -calpain activation mediate the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation.

**Citation:** Liu Y, Su Y, Sun S, Wang T, Qiao X, et al. (2012) Tau Phosphorylation and  $\mu$ -Calpain Activation Mediate the Dexamethasone-Induced Inhibition on the Insulin-Stimulated Akt Phosphorylation. PLoS ONE 7(4): e35783. doi:10.1371/journal.pone.0035783

**Editor:** Victor Sanchez-Margalet, Virgen Macarena University Hospital, Spain

**Received:** November 24, 2011; **Accepted:** March 21, 2012; **Published:** April 20, 2012

**Copyright:** © 2012 Liu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by grants from the National Nature Science Foundation of China (30901386), the Wuhan Science and Technology Bureau, China (200960323132), and the Research Fund for the Doctoral Program of Higher Education of China (200804871026). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: zhihouliang@msn.com

## Introduction

Insulin resistance (IR) is defined as a state in which normal amounts of insulin produce a subnormal biological response [1]. Binding of insulin to the insulin receptor promotes Akt phosphorylation (and activation) at Ser-473 (pAkt) [2]. Inhibition on the insulin-stimulated Akt phosphorylation is commonly used as a marker of IR [2]. IR is a feature of type 2 diabetes mellitus (DM2) and is involved in a wide spectrum of other clinical diseases. It is increasingly recognized that IR may be linked with Alzheimer's disease (AD) [3,4]. For instance, clinical investigations suggest that IR may occur before the onset of mild cognitive impairment (MCI) and may be a marker of AD risk [5,6]. Moreover, rodent studies show that IR increases tau phosphorylation and  $\beta$ -amyloid production, two hallmarks of AD pathology [7,8]. Furthermore, IR exacerbates memory deficits in transgenic mouse models of AD [9,10]. Hence, these studies have led to the qualification of AD as 'an insulin-resistant brain state'. However, little is known about the pathogenic mechanisms that impair insulin action in the context of AD.

Glucocorticoids (GCs) are well known to induce IR in a variety of clinical and experimental settings [11]. In chronic IR and DM2,

the concentration of circulating GCs increases. AD patients may also have higher plasma GCs levels than normal controls [12]. A high level of GCs has been shown to result in impaired synaptic plasticity, decreased neurogenesis and hippocampal atrophy, thereby yielding memory deficits [13,14]. It is thus suggested that there may be a link between a high level of GCs and AD pathogenesis [12,13,14]. Although a high level of GCs is a confounding factor in both IR and AD, little is known about the molecular link between GCs and IR in the context of AD.

Tau hyperphosphorylation is one of the hallmark pathological alterations in AD [15]. Tau is a microtubule-stabilizing protein and supports the microtubule system responsible for intracellular transport, axonal morphology and cell physiology. Tau hyperphosphorylation occurs long before the onset of AD symptoms and may play a crucial role in neurodegeneration in AD [15,16]. Although it has been suggested that tau hyperphosphorylation may contribute independently to AD pathogenesis and/or progression [17,18], little is known about the molecular mechanisms underlying tau-mediated toxicities. Because both GCs and IR have been shown to lead to tau hyperphosphorylation [19,20], it is reasonable to postulate that GCs and IR may be linked by tau. We have previously shown that tau might mediate the GCs-

induced regulation of cAMP-dependent kinase and cAMP response element binding protein in HEK293 cells stably expressing the longest human tau isoform (tau-441; HEK293/tau441 cells) [21]. We were interested to examine whether tau mediates the GCs-induced IR.

Calpains, a family of calcium-activated cysteine proteases, have been reported to be involved in the pathogenesis of AD [22,23]. Our previous work and other studies have shown that  $\mu$ -calpain (also called calpain 1) of the calpains family may play a significant role in AD [24,25]. Dysregulation of calpains is also associated with the risk of IR and DM2 [26,27,28,29]. Although many studies have focused on calpain 10, more recent evidence also highlights the role of  $\mu$ -calpain in IR and DM2 [30,31]. Moreover, activation of  $\mu$ -calpain is also implicated in the GCs-induced toxicities in muscles and brains [32,33]. We thus assumed that GCs and IR may be also linked by  $\mu$ -calpain. It has been shown that activation of  $\mu$ -calpain mediates the tau-induced toxicities in rats cortical neurons over-expressing human tau [34]. We thus wanted to examine whether  $\mu$ -calpain also mediates the GCs-induced IR, and if so, whether  $\mu$ -calpain is linked with tau.

To address the issues mentioned above, we employed two cell lines, wild-type HEK293 cells HEK293/tau441 cells, a good cellular model to study tau pathology [35,36]. We examined whether DEX, a synthetic GCs, inhibits the insulin-stimulated Akt phosphorylation and if so, whether tau or  $\mu$ -calpain is involved in the inhibitory effect of DEX. In HEK293/tau441 cells, we made the following observations. 1) DEX increases tau phosphorylation and induces tau-mediated  $\mu$ -calpain activation. 2) The effects of DEX on tau phosphorylation and  $\mu$ -calpain activation are prevented by LiCl pre-treatment. 3) Either LiCl pre-treatment or calpain inhibition prevents the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation.

## Materials and Methods

### 1. Antibodies and chemicals

Mouse monoclonal antibody (mAb) against  $\beta$ -actin was purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Rabbit mAb against Akt and phosphorylated Akt at Ser473 (pAkt) were obtained from Cell Signaling Technology (Beverly, MA, USA). Mouse mAb against  $\mu$ -calpain was purchased from Sigma (St. Louis, MO, USA). Mouse mAb against Tau-1 was obtained from Millipore (Billerica, MA, USA). The antibody R134d that recognizes total tau was generously provided by Gong Cheng-Xin (NYS Institute for Basic Research, SI, NY, USA). Dexamethasone (DEX), insulin, mifepristone (RU), lithium chloride (LiCl) and E-64d (calpain inhibitor) were purchased from Sigma (St. Louis, MO, USA).

### 2. Cell lines, culture and treatment

HEK293 cells stably transfected with the longest isoform of recombinant human tau (tau441; HEK293/tau441) and HEK293 cells transfected with an empty vector (HEK293/vec) were a generous gift from Professor Qing Tian (Department of Pathophysiology, Tongji Medical College, Huazhong University of Science and Technology). The construction of HEK293/tau441 and HEK293/vec cells is as previously described [35,36]. For control purposes, wild-type HEK293 cells were used and compared with HEK293/vec cells; no differences between the two cell lines were found. Both wild-type HEK293 and HEK293/tau441 were grown in Dulbecco's modified Eagle's medium (DMEM; Invitrogen, Eggenstein, Germany), supplemented with 10% (v/v) fetal bovine serum (FBS; Gibco, Carlsbad, CA, USA) and maintained in 37°C and 5% CO<sub>2</sub>.

For the experiments, both HEK293/tau441 and wild-type HEK293 cells were first incubated in DMEM with 2% FBS overnight in order to reduce the serum effects. The cells were then pre-treated with RU (20  $\mu$ M, 30 minutes), LiCl (10 mM, 1 hour), or E-64d (30  $\mu$ g/ml, 1 hour) and then treated with DEX (1  $\mu$ M), or treated with DEX (1  $\mu$ M) alone. For treatment longer than three days, the medium was changed at 72 hours (h) after exposure to DEX. Then RU (20  $\mu$ M), LiCl (10 mM) or E-64d (30  $\mu$ g/ml) with DEX (1  $\mu$ M) or DEX (1  $\mu$ M) were added to the media. After incubation with the above agents, cells were exposed to 100 nM insulin for 15 minutes (min) and then harvested for Western blotting.

### 3. Western blotting

The total proteins from the cultures were extracted with the ice-cold buffer containing 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 0.5% sodium deoxycholate, 1% NP-40, 0.1% SDS, 100  $\mu$ M sodium orthovanadate, 1 mM phenylmethylsulfonyl fluoride and 1  $\mu$ g/ml protease inhibitor cocktail (Beyotime Biotechnology, Haimen, Jiangsu, China). The cell lysates were boiled, briefly sonicated and centrifuged at 12500 $\times$ g for 15 min. The protein concentration in each lysate was determined using a bicinchoninic acid (BCA) Protein Assay Kit (Beyotime Biotechnology, Haimen, China) with the bovine serum albumin as the standard. Equal amounts of proteins (60–80  $\mu$ g) were separated on 10% SDS-polyacrylamide gel (SDS-PAGE) and transferred to nitrocellulose membranes. The membranes were blocked in 5% nonfat milk in TTBS (10 mM Tris-HCl, 150 mM NaCl, 0.02% Tween-20, pH 7.5) and probed overnight at 4°C with antibodies against  $\beta$ -actin (1:400), pAkt, Akt, Tau-1 and  $\mu$ -calpain (1:1000) and total tau (R134d, 1:10 000). The blots were developed with horseradish peroxidase-conjugated secondary anti-mouse (1: 1000; Thermo scientific, Rockford, IL, USA) or anti-rabbit antibodies (1: 3000; Signalway antibody, Pearland, TX, USA) and visualized with an enhanced chemiluminescent substrate kit (Thermo scientific, Rockford, IL, USA) and exposure to CL-XPosure film (Kodak). The films were scanned and the density of each of band was quantified using Kodak Digital Science 1D software (Eastman Kodak Co., New Haven, CT).


### 4. Statistical analysis

Numerical data were expressed as group means  $\pm$  SEM and analyzed with SPSS statistical software (version 13.0). All data sets were subjected to one-way ANOVA before application of *post hoc* pair-wise comparisons.  $P < 0.05$  or lower was considered to represent statistically significant differences.

## Results

### 1. DEX induces inhibition on the insulin-stimulated Akt phosphorylation in wild-type HEK293 and HEK293/tau441 cells

The insulin-stimulated increase in pAkt is a key event in the proximal insulin signalling pathway and the pAkt/Akt ratio has been widely used as a marker for evaluating IR [2]. We therefore determined whether GCs induce IR by examining the insulin-stimulated Akt phosphorylation (pAkt/Akt ratio) in wild-type HEK293 and HEK293/tau441 cells. Dexamethasone (DEX), a synthetic GCs and specific glucocorticoid receptor (GR) agonist, was used in our study as GCs. The concentration of DEX was 1  $\mu$ M because it is in the range of the amount given for the therapeutic purpose in clinical practice (e.g., dexamethasone suppression test, prevention of respiratory distress in preterm infants) and also applied for induction of long-term stress [37,38].



**Figure 1. Effects of dexamethasone on the insulin-stimulated Akt phosphorylation.** A): Both wild-type HEK293 and HEK293/tau441 cells were treated with 1  $\mu$ M dexamethasone (DEX) for 1–6 days and then stimulated with 100 nM insulin (i) for 15 min. Representative immunoblots from the results on the third day (D3) and the sixth day (D6) after DEX treatment were shown. Bars representing means  $\pm$  SEM were shown below. Total amounts of Akt remained stable under the conditions. In HEK293/tau441 cells, DEX prevented the insulin-stimulated increases in pAkt on D3 and D6. In wild-type HEK293 cells, the inhibitory effect of DEX was evident on D6 but not on D3. Each

experiment was repeated three times unless stated otherwise. \* $P < 0.05$  versus control. B): Culture cells were pre-treated with mifepristone (RU; 20  $\mu$ M, 30 min) and then treated with DEX for 3 days or 6 days. Representative immunoblots from wild-type HEK293 cells on D6 and those from HEK293/tau441 cells on D3 were shown, with bars representing means  $\pm$  SEM below. Pre-treatment with RU prevented the inhibitory effects of DEX. \* $P < 0.05$  versus DEX group. doi:10.1371/journal.pone.0035783.g001


Long-term (1–6 days) treatment with DEX was applied to both cell lines (see *Materials and Methods*) to mimic the induction of long-term stress *in vivo* [37,39]. Representative immunoblots from the results obtained on the third day (D3) and on the sixth day (D6) after DEX treatment were shown in Fig. 1. As shown in Fig. 1, total amounts of Akt remained stable under the conditions and were used as internal standards for measurement of Akt phosphorylation. Under basal conditions, insulin induced similar increases in pAkt in both cell lines (lane 2 and 6, Fig. 1A). This suggests that in HEK293/tau441 cells, the exogenously expressed tau by itself does not affect the insulin-stimulated Akt phosphorylation. Treatment with DEX alone did not affect the basal pAkt levels in either of the two cell lines (lane 3 and 7, Fig. 1A). In wild-type HEK293 cells, the inhibitory effect of DEX was not evident until D6 (D6, lane 4, Fig. 1A). In HEK293/tau441 cells, the inhibitory effect of DEX occurred on D3 (D3, lane 8, Fig. 1A) and then sustained on day 4, day 5 (data not shown) and D6 (D6, lane 8, Fig. 1A). We thus chose D3 and D6 as two representative time points for the following experiments.

There two types of receptors that GCs can bind with, one is glucocorticoid receptor (GR) and the other is mineralocorticoid receptor (MR). Because HEK293 cells do not express MR [40], we only used mifepristone (RU), a GR antagonist. Representative immunoblots from wild-type HEK293 cells on D6 and those from HEK293/tau441 cells on D3 were shown in Fig. 1B. Without insulin stimulation, pre-treatment with RU (20  $\mu$ M, 30 min; see *Materials and Methods*) did not affect the pAkt levels upon DEX treatment (lane 3, Fig. 1B). However, pre-treatment with RU prevented the inhibitory effects of DEX in both cell lines (lane 4, Fig. 1B), indicating that the inhibitory effect of DEX is mediated by the GR.

Although DEX inhibited the insulin-stimulated Akt phosphorylation in wild-type HEK293 and HEK293/tau441 cells, the inhibitory effect of DEX occurred earlier in HEK293/tau441 cells (D3 versus D6). Because GCs are known to induce tau phosphorylation [14], we speculate that it is the DEX-induced increase in tau phosphorylation that causes the difference described above.

## 2. DEX increases tau phosphorylation in HEK293/tau441 cells, which is prevented by lithium chloride

We next assessed whether DEX increases tau phosphorylation in HEK293/tau441 cells, and if so, whether lithium chloride (LiCl), an inhibitor of glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ) that has been shown to inhibit tau hyperphosphorylation in different contexts [41,42,43], can prevent the action of DEX. To this end, Western blots were performed using Tau-1 and R134d antibodies. Tau-1 is an antibody that recognizes tau when it is dephosphorylated within the epitope 189–207 [44]. R134d is an antibody that recognizes total tau. HEK293/tau441 cells were pre-treated with LiCl (10 mM, 1 h) and then treated with DEX (see *Materials and Methods*) for 3 days (D3) or 6 days (D6). Total amounts of tau (detected by R134d) remained stable under the conditions and were used as internal standards for measurement of the relative level of Tau-1 (Tau-1/R134d ratio, Fig. 2). As expected, DEX



**Figure 2. Effects of DEX and lithium chloride on tau phosphorylation.** HEK293/tau441 cells were pre-treated with lithium chloride (LiCl; 10 mM, 1 h) and then treated with DEX for 3 days (D3) or 6 days (D6), followed by Western blotting analysis of tau phosphorylation with Tau-1 (against dephosphorylated tau within the epitope 189–207) and R134d (against total tau) antibodies. Total amounts of tau remained largely stable and the level of tau phosphorylation was determined by the Tau-1/R134d ratio. Bars representing means  $\pm$  SEM. On D3, DEX induced an increase in tau phosphorylation, as evidenced by the decrease in the Tau-1/R134d ratio. Pre-treatment with LiCl prevented the effect of DEX on tau phosphorylation on D3. \* $P < 0.05$  versus control, # $P < 0.05$  between indicated groups. doi:10.1371/journal.pone.0035783.g002

induced an increase in tau phosphorylation on D3, as evidenced by the decrease in the level of Tau-1 (lane 2, Fig. 2). However, DEX did not affect the Tau-1 level on D6 (lane 6, Fig. 2). LiCl treatment alone did not affect basal Tau-1 levels (lane 3 and 7, Fig. 2). However, pre-treatment with LiCl prevented the DEX-induced decrease in Tau-1 on D3 (lane 4, Fig. 2), suggesting that the DEX-induced increase in tau phosphorylation is prevented by LiCl. Interestingly, although LiCl is widely accepted as an inhibitor to GSK-3 $\beta$  [41], GSK-3 $\beta$  activity, measured by the ratio of the phosphorylated GSK-3 $\beta$  at Ser-9 (p GSK-3 $\beta$ ) and total GSK-3 $\beta$ , was not obviously affected by DEX in HEK293/tau441 cells on D3 (data not shown). DEX actually decreased GSK-3 $\beta$  activity (an increase in pGSK-3 $\beta$ /GSK-3 $\beta$  ratio) in wild-type HEK293 cells on D3 (data not shown). The mechanisms responsible for the effects of DEX and LiCl on tau phosphorylation in our system warrants further research. Nevertheless, results described above show that LiCl pre-treatment prevents the DEX-induced increase in tau phosphorylation in HEK293/tau441 cells.


### 3. Insulin does affect tau phosphorylation in HEK293/tau441 cells

We have shown that LiCl prevented the DEX-induced increase in tau phosphorylation in HEK293/tau441 cells. To test our hypothesis that the DEX-induced increase in tau phosphorylation

mediates the inhibition on the insulin-stimulated Akt phosphorylation, we should examine whether LiCl corrects the inhibitory effects of DEX. However, before that we also wanted to see how insulin affects tau phosphorylation because previous studies demonstrated that insulin could induce either tau phosphorylation or dephosphorylation [45,46,47]. Moreover, insulin is well known to activate the Akt/GSK-3 $\beta$  signaling pathway and thus inhibits the activity of GSK-3 $\beta$  [45,46]. Because we have shown GSK-3 $\beta$  inhibition by LiCl prevented the DEX-induced tau phosphorylation (Fig. 2), we examined the effect of insulin on tau phosphorylation. HEK293/tau441 cells were pre-treated with LiCl (10 mM, 1 h) and then treated with DEX (see *Materials and Methods*) for 3 days (D3), followed by insulin stimulation. Total amounts of tau remained stable under the conditions (Fig. 3). Insulin stimulation under basal condition did affect the Tau-1 level (lane 2, Fig. 3). Upon treatment with LiCl or co-treatment with DEX and LiCl, insulin did not affect Tau-1 levels either (lane 6 and 8, Fig. 3). Importantly, insulin did not prevent the DEX-induced decrease in Tau-1 (lane 4, Fig. 3). These results suggest that insulin stimulation, at least in our system, does not affect tau phosphorylation.

### 4. Lithium chloride prevents the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation in HEK293/tau441

We have shown that LiCl prevented the DEX-induced increase in tau phosphorylation and insulin did not affect tau phosphorylation. We next examined whether LiCl corrects the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation in HEK293/tau441 cells. However, it has also been reported that




**Figure 3. Effects of insulin on tau phosphorylation.** HEK293/tau441 cells were pre-treated with LiCl (10 mM, 1 h) and then treated with DEX for 3 days, followed by stimulation of insulin for 15 min and Western analysis of Tau-1 and R134d. Bars representing means  $\pm$  SEM. Insulin did not prevent the DEX-induced decrease in Tau-1 or affected Tau-1 levels under other conditions, suggesting that insulin does not affect tau phosphorylation. \* $P < 0.05$  versus control. doi:10.1371/journal.pone.0035783.g003

LiCl affected insulin signaling through tau-independent mechanisms [48]. We therefore also examined how LiCl affects the insulin-stimulated Akt phosphorylation in wild-type HEK293 cells. Both cell lines were treated with LiCl, DEX, or in combination of them for 3 days (D3) or 6 days (D6). In both cell lines, LiCl did not have obvious effects on pAkt levels under basal conditions, under insulin stimulation alone or upon DEX treatment alone (lane 5, 6 and 7, Fig. 4). LiCl prevented the inhibitory effect of DEX in HEK293/tau441 cells on D3 (HEK293/tau441, lane 8, Fig. 4A). However, on D6, when the inhibitory effects of DEX were evident in both cell lines (lane 4, Fig. 4B), pre-treatment with LiCl did not seem to have obvious effects on the DEX-induced inhibition in either of the cell lines on D6 (lane 8, Fig. 4B). Notice that the DEX-induced increase in tau phosphorylation was evident on D3 but not on D6 in HEK293/tau441 cells (Fig. 2). Taken together, the above results suggest that the DEX-induced increase in tau phosphorylation may mediate the inhibitory effect of DEX in HEK293/tau441 cells.

### 5. DEX induces tau-mediated $\mu$ -calpain activation and inhibition of calpain activity prevents the negative effects of DEX in HEK293/tau441 cells

We have shown that the DEX-induced increase in tau phosphorylation mediated the inhibitory effect of DEX in HEK293/tau441 cells. According to our hypothesis, we further determined if  $\mu$ -calpain activation also mediates the inhibitory effects of DEX, and if so, whether the DEX-induced  $\mu$ -calpain activation is mediated by tau phosphorylation. Activation of  $\mu$ -calpain was determined by the ratio of the active/truncated calpain (78-kDa band) and the inactive/full-length calpain (80-kDa band). DEX did not affect  $\mu$ -calpain activation in wild-type HEK293 cells on D3 or on D6 (lane 2, Fig. 5A). DEX did not affect  $\mu$ -calpain activation in HEK293/tau441 cells on D6 either (D6, lane 6, Fig. 5A). In contrast, DEX induced an increase in  $\mu$ -calpain activation in HEK293/tau441 cells on D3 (D3, lane 2, Fig. 5A). Because the DEX-induced increase in tau phosphorylation was evident on D3 (lane 2, Fig. 2), the above results suggest that the DEX-induced tau phosphorylation may correlate with  $\mu$ -



**Figure 4. Effects of LiCl on the DEX-induced inhibitory effect.** Both wild-type HEK293 and HEK293/tau441 cells were pre-treated with LiCl (10 mM, 1 h) and then treated with DEX for 3 days (D3; A) or 6 days (D6; B), followed by stimulation of insulin and Western analysis of pAkt and Akt. Bars representing means  $\pm$  SEM. Pre-treatment with LiCl prevented the inhibitory effect of DEX in HEK293/tau441 cells on D3. doi:10.1371/journal.pone.0035783.g004

calpain activation in HEK293/tau441 on D3. However, it has been shown that  $\mu$ -calpain could result in tau phosphorylation [49]. To examine whether  $\mu$ -calpain activation mediates the DEX-induced tau phosphorylation, we pre-treated HEK293/tau441 cells with E-64d (30  $\mu$ g/ml, 1 h), a cell-membrane-permeable calpain inhibitor, and then determined the levels of Tau-1 and total tau by Western blotting. As Fig. 5B shows, pretreatment with E-64d did not prevent the DEX-induced decrease in Tau-1 in HEK293/tau441 cells. This suggests that  $\mu$ -calpain may not mediate the DEX-induced tau phosphorylation. In contrast, pretreatment with LiCl (10 mM, 1 h) prevented the DEX-induced  $\mu$ -calpain activation in HEK293/tau441 cells on D3 (D3, lane 4, Fig. 5A). Notice that LiCl did not affect  $\mu$ -calpain activation in HEK293/tau441 cells on D6 or in wild-type HEK293 cells (Fig. 5A). Because  $\mu$ -calpain activation was observed only in the tau-expressing HEK293/tau441 cells and prevention of tau phosphorylation by LiCl inhibited the DEX-induced  $\mu$ -calpain activation, these results suggest that the DEX-induced tau phosphorylation mediates  $\mu$ -calpain activation in HEK293/tau441 cells.

Next we assessed whether the DEX-induced and tau-mediated  $\mu$ -calpain activation mediates the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation on D3. However, previous studies have shown that calpain may inhibit insulin signaling [50–53]. We also examined how calpain inhibition would affect the inhibitory effect of DEX on D6, when no activation of  $\mu$ -calpain was observed upon DEX treatment (D6, lane 2, Fig. 5A). Both cell lines were pre-treated with E-64d (30  $\mu$ g/ml, 1 h) and then treated with DEX for 3 days (D3) or 6 days (D6) (See *Materials and Methods*). In wild-type HEK293 cells, E-64d did not affect the insulin-stimulated increase in pAkt upon DEX treatment on D3 (wild-type HEK293, D3, lane 6, Fig. 5C) or prevented the inhibitory effect of DEX on D6 (wild-type HEK293, D6, lane 8, Fig. 5C). In HEK293/tau441 cells, E-64d did not prevent the inhibitory effect of DEX on D6 (HEK293/tau441, D6, lane 6, Fig. 5C). Notice the DEX-induced tau phosphorylation was not evident on D6 (Fig. 2). On D3, when the DEX-induced tau phosphorylation was evident (Fig. 2), E-64d prevented the inhibitory effect of DEX (HEK293/tau441, D3, lane 6, Fig. 5C). These results suggest that the involvement of  $\mu$ -calpain in the inhibitory effect of DEX may depend on tau phosphorylation in our system. Taken together, the above results suggest that the DEX-induced  $\mu$ -calpain activation may also mediate the inhibitory effect of DEX. Moreover, the involvement of  $\mu$ -calpain seems to depend on the DEX-induced tau phosphorylation.

## Discussion


In the current study, several observations have been made using a cellular model system for studying tau pathology, i.e., a HEK293 cell line stably expressing the longest isoform of human tau (HEK293/tau441 cells). First, DEX increased tau phosphorylation and induced tau-mediated  $\mu$ -calpain activation. Second, the effects of DEX on tau phosphorylation and  $\mu$ -calpain activation were prevented by LiCl pre-treatment. Third, both LiCl pre-treatment and calpain inhibition by E-64d prevented the DEX-induced inhibition on the insulin-stimulated Akt phosphorylation.

Numerous potential mechanisms have been proposed to explain the pathogenic mechanisms of IR [54]. However, these studies mainly focused in three main targets, i.e., liver, skeletal muscle and adipose tissue [55,56]. The origin of IR in the context of AD is difficult to elucidate due to the diverse set of risk factors linked to AD. Our results suggest that the GCs-induced tau phosphorylation and consequent  $\mu$ -calpain activation may mediate IR. This

suggests that GCs and IR, two potent driving forces for AD [4,13], may be linked by tau phosphorylation in the context of AD. However, neither LiCl nor E-64d prevented the inhibitory effect of DEX on D6, when the inhibitory effects of DEX were evident in both cell lines. This suggests that the involvement of the GCs-induced tau phosphorylation and consequent  $\mu$ -calpain activation in IR may be only within a particular time window.

Previous rodent studies suggest that IR or lack of insulin signaling can result in tau hyperphosphorylation [20,57,58]. Indeed, it is generally accepted that insulin activates Akt signaling, inhibits GSK-3 $\beta$  activity and thus may reduce tau phosphorylation [45]. It is reasonable to speculate that IR may lead to tau phosphorylation by activating GSK-3 $\beta$ . However, our results show that insulin does not affect tau phosphorylation under basal condition or upon treatment with DEX (Fig. 3). These results suggest that i) insulin signaling may not affect tau phosphorylation, and ii) the DEX-induced tau phosphorylation is not mediated by inhibition of the insulin signaling in our system. In fact, there is a debate about the relationship with the insulin-activated Akt/GSK-3 $\beta$  signaling and tau phosphorylation. In brain/neuron-specific insulin receptor knockout mice, lack of insulin signaling in the brain activates GSK-3 $\beta$  and leads to tau hyperphosphorylation [57]. Intriguingly, in high fat diet C57BL6 mice, IR actually reduces tau phosphorylation at Ser396, Ser202/Thr205 and Thr231 [59]. Moreover, in C57BL mice, insulin administration does not affect tau phosphorylation at GSK-3 $\beta$ -relevant sites but promotes tau phosphorylation at sites that are dependent on extracellular regulated kinase [60]. Interestingly, insulin transiently promotes tau phosphorylation by activating GSK-3 $\beta$  in SH-SY5Y neuroblastoma cells and primary cortical neurons [46,61]. Furthermore, in C57BL mice intraperitoneally injected with streptozotocin, which leads to insulin depletion, tau phosphorylation is mediated by the decrease in protein phosphatase 2A activity but not by the increase in GSK-3 $\beta$  activity [58]. How lack of insulin signaling and IR affect tau phosphorylation *in vivo* and in AD patients warrants further research. Nevertheless, our study suggests that i) the GCs-induced tau phosphorylation may not be mediated by inhibition of insulin signaling, and ii) the GCs-induced tau phosphorylation could function upstream as a mediator of IR.

Clinical observations suggest that DM2 is among the risk factors for AD [5,6,62–64]. Because tau hyperphosphorylation is one of the key pathological features of AD, our results that the DEX-induced tau phosphorylation could occur upstream to IR do not seem to be supported by these clinical data. However, it should be noticed that tau hyperphosphorylation and the detectable clinical symptoms of AD are two different notions. Tau hyperphosphorylation may actually occur decades before individuals manifest clinical symptoms of AD [65]. Tau phosphorylation may also increase during normal aging [66]. Thus it is not surprising that tau hyperphosphorylation also occurs in asymptomatic patients with Braak-stage I neuropathology where clinical symptoms of AD are absent [65]. Because tau hyperphosphorylation can occur long before the onset of symptoms and may play a key role in AD pathogenesis and/or progression, it is possible that tau hyperphosphorylation could contribute to the brain IR in AD. Because both tau phosphorylation and the prevalence of IR increases with aging [67], it might be difficult to identify which of these two events first occurs in brain. It should be also noticed that brain IR and peripheral IR may be two different pathological alterations [68]. The extant literature does not resolve whether the brain IR in AD represents a local disease process, or complication/extension of peripheral IR, i.e. chronic hyperglycemia, hyperinsulinemia, and DM2.



**Figure 5. Involvement of  $\mu$ -calpain in the inhibitory effect of DEX.** A): Both wild-type HEK293 and HEK293/tau441 cells were pre-treated with LiCl (10 mM, 1 h) and then treated with DEX for 3 days (D3) or 6 days (D6). Activation of  $\mu$ -calpain was determined by the ratios of the active/truncated calpain (78-kDa bands) and the inactive/full-length calpain (80-kDa bands). Bars representing means  $\pm$  SEM. On D3 in HEK293/tau441 cells, DEX induced  $\mu$ -calpain activation and pre-treatment with LiCl prevented the activation of  $\mu$ -calpain. \* $P$ <0.05 versus control. B): HEK293/tau441 cells were pre-treated with E-64d (30  $\mu$ g/ml, 1 h) or LiCl (10 mM, 1 h), and then treated with DEX for 3 days. The level of tau phosphorylation was determined by the ratio of Tau-1/R134d. Bars representing means  $\pm$  SEM. E-64d did not have obvious effect on the DEX-induced increase in tau phosphorylation. \* $P$ <0.05 versus control, # $P$ <0.05 between indicated groups. C) Both wild-type HEK293 and HEK293/tau441 cells were pre-treated with E-64d (30  $\mu$ g/ml, 1 h) and then treated with DEX for 3 days (D3) or 6 days (D6), followed by stimulation of insulin and Western analysis of pAkt and Akt. Bars representing means  $\pm$  SEM. Pre-treatment with E-64d prevented the inhibitory effect of DEX in HEK293/tau441 cells on D3. \* $P$ <0.05 versus control.

doi:10.1371/journal.pone.0035783.g005

Although the inhibitory effects of DEX were evident in both cell lines on D6, calpain inhibition did not prevent the inhibitory effects in either of the cell lines (Fig. 5C). In contrast, calpain inhibition prevented the inhibitory effect of DEX only on D3, when the DEX-induced tau phosphorylation was evident (Fig. 2 and Fig. 5C). These results suggest that the involvement of  $\mu$ -calpain in the GCs-induced IR may not be a universal phenomenon and the action of  $\mu$ -calpain seems to depend on the GCs-induced tau phosphorylation. The roles of calpains in the pathogenesis of IR and DM2 have been extensively studied [28,29,30,31]. Although Calpain 10 is the first identified diabetes gene through a genome scan [69], little is known about its protease activity and physiological functions [70]. Furthermore, whether calpain 10 really links with IR and DM2 is still in controversy [70–72]. Some recent studies have suggested that  $\mu$ -calpain activation may contribute to the inhibition of insulin signaling. For instance,  $\mu$ -calpain cleaves phosphoinositide 3-kinases (PI3Ks), an important upstream activator of Akt, and reduces PI3Ks enzymatic activities in NIH 3T3 cells [53]. Moreover,  $\mu$ -calpain degrades insulin receptor substrates (IRS), major substrates of insulin signaling, in human neuroblastoma cells and prostate epithelial cells [50,52]. Furthermore, calpain activation by calcium incubation reduced total amount of Akt and heat shock protein 90, a key Akt regulator protein, in rats skeletal muscle cells [51]. Although these studies suggest that  $\mu$ -calpain activation may mediate IR, other studies also show contradictory results. For instance, calpain inhibition by overexpression of calpastatin (the endogenous calpain inhibitor) induced a decrease in Akt level and subsequent IR in mice [73]. Moreover, in soleus muscle strips and adipocytes and adipocytes isolated from Wistar rats, calpain inhibition by E-64d reduced insulin-mediated signaling [74]. Furthermore, in 3T3-L1 adipocytes and HepG2 hepatoma cells, inhibition of calpain did not

affect the insulin-mediated PI3K/Akt pathway but it did prevent the insulin-stimulated glucose uptake [75,76]. These studies suggest that the roles of  $\mu$ -calpain in IR may be tissue-specific. Our results suggest that the involvement of  $\mu$ -calpain in IR, at least in tau-expressing tissues, may depend on the GCs-induced tau phosphorylation.

However, how the GCs-induced tau phosphorylation mediates  $\mu$ -calpain activation is not known. The activation and regulation of calpain activity in vivo is not well understood, although it is a ubiquitously expressed protein with limited proteolytic activity and therefore it is assumed to be tightly regulated [77]. We found an enhanced co-localization of phosphorylated tau and  $\mu$ -calpain in HEK293/tau441 and SH-SY5Y cells upon DEX treatment (unpublished data). We speculate that the GCs-induced tau phosphorylation may enhance the structural link between tau and  $\mu$ -calpain and thus mediate the activation of  $\mu$ -calpain. Interestingly, in a postmortem study of AD patients' brains, it was found that the hyperphosphorylated tau can bind with a calcium-binding protein—EF hand domain protein 2, which may be involved in modification of calpain activity [78].

In conclusion, we have demonstrated that the GCs-induced tau phosphorylation and consequent  $\mu$ -calpain activation may contribute to IR. Our study suggests that tau phosphorylation and  $\mu$ -calpain activation may be two key events that link GCs, IR and AD.

## Author Contributions

Conceived and designed the experiments: YL YS ZL TW XR SS. Performed the experiments: YL YS. Analyzed the data: YL YS XQ XR ZL. Contributed reagents/materials/analysis tools: SS TW XR ZL. Wrote the paper: YL ZL.

## References

- Reaven GM (2011) Insulin resistance: the link between obesity and cardiovascular disease. *Med Clin North Am* 95: 875–892.
- Tsoyi K, Jang HJ, Nizamutdinova IT, Park K, Kim YM, et al. (2010) PTEN differentially regulates expression of ICAM-1 and VCAM-1 through PI3K/Akt/GSK-3 $\beta$ /GATA-6 signaling pathways in TNF- $\alpha$ -activated human endothelial cells. *Atherosclerosis* 213: 115–121.
- Lu FP, Lin KP, Kuo HK (2009) Diabetes and the risk of multi-system aging phenotypes: a systematic review and meta-analysis. *PLoS One* 4: e4144.
- Park SA (2011) A common pathogenic mechanism linking type-2 diabetes and Alzheimer's disease: evidence from animal models. *J Clin Neurol* 7: 10–18.
- Baker LD, Cross DJ, Minoshima S, Belongia D, Watson GS, et al. (2011) Insulin resistance and Alzheimer-like reductions in regional cerebral glucose metabolism for cognitively normal adults with prediabetes or early type 2 diabetes. *Arch Neurol* 68: 51–57.
- Craft S, Baker LD, Montine TJ, Minoshima S, Watson GS, et al. (2012) Intranasal insulin therapy for Alzheimer disease and amnesic mild cognitive impairment: a pilot clinical trial. *Arch Neurol* 69: 29–38.
- Luo D, Hou X, Hou L, Wang M, Xu S, et al. (2011) Effect of pioglitazone on altered expression of A $\beta$  metabolism-associated molecules in the brain of fructose-drinking rats, a rodent model of insulin resistance. *Eur J Pharmacol* 664: 14–19.
- Hiltunen M, Khandelwal VK, Yaluri N, Tiilikainen T, Tusa M, et al. (2011) Contribution of genetic and dietary insulin resistance to Alzheimer phenotype in APP/PS1 transgenic mice. *J Cell Mol Med* in press.
- Cao D, Lu H, Lewis T, Li L (2007) Intake of sucrose-sweetened water induces insulin resistance and exacerbates memory deficits and amyloidosis in a transgenic mouse model of Alzheimer Disease. *J Bio Chem* 282: 36275–36282.
- Ke YD, Delerue F, Gladbach A, Gotz J, Itner LM (2009) Experimental diabetes mellitus exacerbates tau pathology in a transgenic mouse model of Alzheimer's disease. *PLoS One* 4: e7917.
- Houstis N, Rosen ED, Lander ES (2006) Reactive oxygen species have a causal role in multiple forms of insulin resistance. *Nature* 440: 944–948.
- Harmann A, Veldhuis JD, Deuschle M, Standhardt H, Heuser I (1998) Twenty-four hour cortisol release profiles in patients with Alzheimer's and Parkinson's disease compared to normal controls: ultradian secretory pulsatility and diurnal variation. *Neurobiol Aging* 18: 285–289.
- Lupien SJ, Leon MD, Santi SD, Antonio C, Tarshish C, et al. (1998) Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nat Neurosci* 1: 69–73.
- Rissman RA (2009) Stress-induced tau phosphorylation: functional neuroplasticity or neuronal vulnerability? *J Alzheim Dis* 18: 453–457.
- Congdon EE, Duff KE (2008) Is tau aggregation toxic or protective. *J Alzheimers Dis* 14: 453–457.
- Gong CX, Grundke-Iqbal I, Iqbal K (2010) Targeting tau protein in Alzheimer's disease. *Drugs Aging* 27: 351–365.
- Dimakopoulos AC (2005) Protein aggregation in Alzheimer's disease and other neuropathological disorders. *Curr Alzheimer Res* 2: 19–28.
- Cowan CM, Bossing T, Page A, Shepherd D, Muhler A (2010) Soluble hyperphosphorylated tau causes microtubule breakdown and functionally compromises normal tau in vivo. *Acta Neuropathol* 120: 593–604.
- Green KN, Billings LM, Roozendaal B, McLaugh JL, LaFerla FM (2006) Glucocorticoids increase amyloid-beta and tau pathology in a mouse model of Alzheimer's disease. *J Neurosci* 26: 9047–9056.
- Bosco D, Fava A, Plastino M, Montalcini T, Pujia A (2011) Possible implications of insulin resistance and glucose metabolism in Alzheimer's disease pathogenesis. *J Cell Mol Med* 15: 1807–1821.
- Liu Y, Su Y, Sun S, Wang T, Qiao X, et al. (2012) Human tau may modify glucocorticoids-mediated regulation of cAMP-dependent kinase and phosphorylated cAMP response element binding protein. *Neurochem Res* Feb 1. [Epub ahead of print].
- Nixon RA (2003) The calpains in aging and aging related disease. *Ageing Res Rev* 2: 407–418.
- Trinchese F, Fa M, Liu S, Zhang H, Hidalgo A, et al. (2008) Inhibition of calpains improves memory and synaptic transmission in a mouse model of Alzheimer disease. *J Clin Invest* 118: 2796–2807.
- Liu F, Grundke-Iqbal I, Iqbal K, Oda Y, Tomizawa K, et al. (2005) Truncation and activation of calcineurin A by calpain I in Alzheimer disease brain. *J Biol Chem* 280: 37755–37762.
- Liang Z, Liu F, Grundke-Iqbal I, Iqbal K, Gong CX (2007) Down-regulation of cAMP-dependent protein kinase by over-activated calpain in Alzheimer disease brain. *J Neurochem* 103: 2462–2470.
- Permut MA, Bernal-Mizrachi E, Inoue H (2000) Calpain 10: the first positional cloning of a gene for type 2 diabetes? *J Clin Invest* 106: 819–821.

27. Derbel S, Doumaquet C, Hubert D, Monsnier-Pudar H, Grabar S, et al. (2006) Calpain 10 and development of diabetes mellitus in cystic fibrosis. *J Cyst Fibros* 5: 47–51.
28. Kang ES, Kim HJ, Nam M, Nam CM, Ahn CW, et al. (2006) A novel 111/121 diplotype in the calpain-10 gene is associated with type 2 diabetes. *J Hum Genet* 51: 629–633.
29. Randriamboavonjy V, Pistrosch F, Bolck B, Schwinger RHG, Dixit M, et al. (2008) Platelet sacroplasmic endoplasmic reticulum  $Ca^{2+}$ -ATPase and  $\mu$ -calpain activity are altered in type 2 diabetes mellitus and restored by rosiglitazone. *Circulation* 117: 52–60.
30. Hsieh YY, Chang CC, Hsu KH, Tsai FJ, Chen CP, et al. (2008) Effect of exercise training on calpain systems in lean and obese Zucker rats. *Int J Biol Sci* 4: 300–308.
31. Liu Y, Liu F, Grundke-Iqbal I, Iqbal K, Gong CX (2011) Deficient brain insulin signaling pathway in Alzheimer's disease and diabetes. *J Pathol* 225: 54–82.
32. Elliott EM, Mattson MP, Vanderklis P, Lynch G, Chang I, et al. (1993) Corticosterone exacerbates kainate-induced alterations in hippocampal tau immunoreactivity and spectrin proteolysis in vivo. *J Neurochem* 61: 57–67.
33. Smith IJ, Lecker SH, Hasselgren PO (2008) Calpain activity and muscle wasting in sepsis. *Endocrinology and metabolism* 295: E762–E771.
34. Amadoro G, Ciotti MT, Costanzi M, Cestari V, Calissano P, et al. (2006) NMDA receptor mediates tau-induced neurotoxicity by calpain and ERK/ MAPK activation. *Prog Natl Acad Sci USA* 103: 2892–2897.
35. Ren QG, Liao XM, Chen XQ, Liu GP, Wang JZ (2007) Effects of tau phosphorylation on proteasome activity. *FEBS Letters* 581: 1521–1528.
36. Yu G, Li Y, Tian Q, Liu R, Wang Q, et al. (2011) Berberine attenuates calyculin A-induced cytotoxicity and tau hyperphosphorylation in HEK293 cells. *J Alzheimer Dis* 24: 525–535.
37. Jameson RR, Seidler FJ, Qiao D, Slotkin TA (2006) Adverse neurodevelopmental effects of dexamethasone modeled in PC12 cells: identifying the critical stages and concentration thresholds for the targeting of cell acquisition, differentiation and viability. *Neuropsychopharmacology* 31: 1647–1658.
38. Yaniv SP, Ben-Shachar D, Klein E (2008) Norepinephrine-glucocorticoids interaction does not annul the opposite effects of the individual treatments on cellular plasticity in neuroblastoma cells. *Eur J Pharmacol* 596: 14–24.
39. Holsboer F (2000) The corticosteroid receptor hypothesis of depression. *Neuropsychopharmacology* 23: 477–501.
40. Pfau A, Grossmann C, Freudinger R, Mildnerberger S, Benesic A, et al. (2007) Ca but not H<sub>2</sub>O<sub>2</sub> modulates GRE-element activation by the human mineralocorticoid receptor in HEK cells. *Mol Cell Endocrinol* 264: 35–43.
41. Sato N, Meijer L, Skaltsounis L, Greengard P, Brivanlou AH (2004) Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3-specific inhibitor. *Nat Med* 10: 55–63.
42. Caccamo A, Oddo S, Tran LX, LaFerla FM (2007) Lithium reduces tau phosphorylation but not A $\beta$  or working memory deficits in a transgenic model of both plaques and tangles. *Am J Pathol* 170: 1669–1678.
43. Leroy K, Ando K, Heraud C, Yilmaz Z, Authalet M, et al. (2010) Lithium treatment arrests the development of neurofibrillary tangles in mutant tau transgenic mice with advanced neurofibrillary pathology. *J Alzheimer Dis* 19: 705–719.
44. Szendrei GI, Lee VM, Otvos Jr. (1993) Recognition of the minimal epitope of monoclonal antibody Tau-1 depends upon the presence of a phosphate group but not its location. *J Neurosci Res* 34: 243–249.
45. Hong M, Lee VMY (1997) Insulin and insulin-like growth factor-1 regulate tau phosphorylation in cultured human neurons. *J Biol Chem* 272: 19547–19553.
46. Lesort M, Jope RS, Johnson GVW (1999) Insulin transiently increases tau phosphorylation. Involvement of glycogen synthase kinase-3 $\beta$  and Fyn Tyrosine kinase. *J Neurochem* 72: 576–584.
47. Planel E, Tatebayashi Y, Miyasaka T, Liu L, Wang L, et al. (2007) Insulin dysfunction induces in vivo tau hyperphosphorylation through distinct mechanisms. *J Neurosci* 27: 13635–13648.
48. Cohen P, Goedert M (2004) GSK3 inhibitors: development and therapeutic potential. *Nat Rev* 3: 479–487.
49. Town T, Zolton J, Shaffner R, Schnell B, Crescentini R, et al. (2002) p35/Cdk5 pathway mediates soluble amyloid-beta peptide-induced tau phosphorylation in vitro. *J Neurosci Res* 69: 362–372.
50. Zhang H, Hoff H, Sell C (2003) Downregulation of IRS-1 protein in thapsigargin-treated human prostate epithelial cells. *Exp Cell Res* 289: 352–358.
51. Smith IJ, Dodd SL (2007) Calpain activation causes a proteasome-dependent increase in protein degradation and inhibits the Akt signalling pathway in rat diaphragm muscle. *Exp Physiol* 92: 561–573.
52. Kim B, Feldman EL (2009) Insulin receptor substrate (IRS)-2, but not IRS-1, protects human neuroblastoma cells against apoptosis. *Apoptosis* 14: 665–673.
53. Beltran L, Chaussade C, Vanhaesebroeck B, Cutillas PR (2011) Calpain interacts with class IA phosphoinositide 3-kinases regulating their stability and signaling activity. *Proc Natl Acad Sci USA* 108: 16217–16222.
54. Nolan CJ, Damm P, Prentki M (2011) Type 2 diabetes across generations: from pathophysiology to prevention and management. *Lancet* 378: 169–181.
55. Gallagher EJ, Leroith D, Karnieli E (2011) The metabolic syndrome—from insulin resistance to obesity and diabetes. *Med Clin North Am* 95: 855–873.
56. Kahn BB, Flier JS (2000) Obesity and insulin resistance. *J Clin Invest* 106: 473–481.
57. Schuber M, Gautam D, Surjo D, Ueki K, Baudler S, et al. (2004) Role for neuronal insulin resistance in neurodegenerative diseases. *Proc Natl Acad Sci U S A* 101: 3100–3105.
58. Clodfelder-Miller BJ, Zmijewska AA, Johnson GVW, Jope RS (2006) Tau is hyperphosphorylated at multiple sites in mouse brain in vivo after streptozotocin-induced insulin deficiency. *Diabetes* 55: 3320–3325.
59. To AW, Ribe EM, Chuang TT, Schroeder JE, Lovestone S (2011) The  $\epsilon$ 3 and  $\epsilon$ 4 alleles of human APOE differentially affect tau phosphorylation in hyperinsulinemic and pioglitazone treated mice. *PLoS One* 6: e16991.
60. Freude S, Plum L, Schnitker J, Leeser U, Udelhoven M, et al. (2005) Peripheral hyperinsulinemia promotes tau phosphorylation in vivo. *Diabetes* 54: 3343–3348.
61. Lesort M, Johnson GV (2000) Insulin-like growth factor-1 and insulin mediate transient site-selective increases in tau phosphorylation in primary cortical neurons. *Neuroscience* 99: 305–316.
62. Craft S (2007) Insulin resistance and Alzheimer's disease pathogenesis: potential mechanisms and implications for treatment. *Curr Alzheimer Res* 4: 147–152.
63. Frisardi V, Solfrizzi V, Capurso C, Imbimbo BP, Vendemiale G, et al. (2010) Is insulin resistant brain state a central feature of the metabolic-cognitive syndrome? *J Alzheimers Dis* 21: 57–63.
64. Riedere RP, Bartl J, Laux G, Grunblatt E (2011) Diabetes type II: a risk factor for depression-Parkinson-Alzheimer? *Neurotox Res* 19: 253–265.
65. Maccioni RB, Farias G, Morales I, Navarrete L (2010) The revitalized tau hypothesis on Alzheimer's disease. *Arch Med Res* 41: 226–231.
66. Mukactova-Ladinska EB, Harrington CR, Roth M, Wischik CM (1996) Alterations in tau protein metabolism during normal aging. *Dementia* 7: 95–103.
67. Grundy SM, Brewer HB, Jr., Cleeman JI, Smith SC, Jr., Lenfant C, et al. (2004) Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation* 109: 433–438.
68. de la Monte SM, Wands JR (2005) Review of insulin and insulin-like growth factor expression, signaling, and malfunctions in the central nervous system: relevance to Alzheimer's disease. *J Alzheimers Dis* 7: 45–61.
69. Turner MD, Cassell PG, Hitman GA (2005) Calpain-10: from genome search to function. *Diabetes Metab Res Rev* 21: 505–514.
70. Suzuki K, Hata S, Kawabata Y, Sorimachi H (2004) Structure, activation, and biology of calpain. *Diabetes Suppl.* 1: S12–S18.
71. Baier IJ, Permana PA, Yang X, Pratley RE, Hanson RL, et al. (2000) A calpain-10 gene polymorphism is associated with reduced muscle mRNA levels and insulin resistance. *J Clin Invest* 106: R69–R73.
72. Horikawa Y, Oda N, Yu L, Imamura S, Fujiwara K, et al. (2003) Genetic variations in calpain-10 gene are not a major factor in the occurrence of type 2 diabetes in Japanese. *J Clin Endocrinol Metab* 88: 244–247.
73. Otani K, Polonsky KS, Holloszy JO, Han DH (2006) Inhibition of calpain results in impaired contraction-stimulated GLUT4 translocation in skeletal muscle. *Am J Physiol Endocrinol Metab* 291: E544–E548.
74. Sreenan SK, Zhou YP, Otani K, Hansen P, Currie KP, et al. (2001) Calpains play a role in insulin secretion and action. *Diabetes* 50: 2013–2020.
75. Paul DS, Harmon AW, Winston CP, Patel YM (2003) Calpain facilitates GLUT4 vesicle translocation during insulin-stimulated glucose uptake in adipocytes. *Biochem J* 376: 625–632.
76. Meier M, Klein HH, Kramer J, Drenckhan M, Schutt M (2007) Calpain inhibition impairs glycogen syntheses in HepG2 hepatoma cells without altering insulin signaling. *J Endocrinol* 193: 45–51.
77. Perrin BJ, Huttenlocher A (2002) Calpain. *Int J Biochem Cell Biol* 34: 722–725.
78. Vega IE, Traverso EE, Ferrer-Acosta Y, Matos E, Colon M, et al. (2008) A novel calcium-binding protein is associated with tau proteins in tauopathy. *J Neurochem* 106: 96–106.