

Analysis of *RAD51D* in Ovarian Cancer Patients and Families with a History of Ovarian or Breast Cancer

Ella R. Thompson^{1*}, Simone M. Rowley¹, Sarah Sawyer², kConFab³, Diana M. Eccles⁴, Alison H. Trainer^{1,2,5}, Gillian Mitchell^{2,5}, Paul A. James^{2,5}, Ian G. Campbell^{1,5,6}

1 VBCRC Cancer Genetics Laboratory, Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia, **2** Familial Cancer Centre, Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia, **3** Kathleen Cuninghame Foundation for Research into Familial Breast Cancer (kConFab), Peter MacCallum Cancer Centre, East Melbourne, Victoria, Australia, **4** Cancer Sciences Division, Faculty of Medicine, University of Southampton, Princess Anne Hospital, Southampton, United Kingdom, **5** Sir Peter MacCallum Department of Oncology, University of Melbourne, Parkville, Victoria, Australia, **6** Department of Pathology, University of Melbourne, Melbourne, Victoria, Australia

Abstract

Mutations in *RAD51D* have been associated with an increased risk of hereditary ovarian cancer and although they have been observed in the context of breast and ovarian cancer families, the association with breast cancer is unclear. The aim of this current study was to validate the reported association of *RAD51D* with ovarian cancer and assess for an association with breast cancer. We screened for *RAD51D* mutations in *BRCA1/2* mutation-negative index cases from 1,060 familial breast and/or ovarian cancer families (including 741 affected by breast cancer only) and in 245 unselected ovarian cancer cases. Exons containing novel non-synonymous variants were screened in 466 controls. Two overtly deleterious *RAD51D* mutations were identified among the unselected ovarian cancer cases (0.82%) but none were detected among the 1,060 families. Our data provide additional evidence that *RAD51D* mutations are enriched among ovarian cancer patients, but are extremely rare among familial breast cancer patients.

Citation: Thompson ER, Rowley SM, Sawyer S, kConFab, Eccles DM, et al. (2013) Analysis of *RAD51D* in Ovarian Cancer Patients and Families with a History of Ovarian or Breast Cancer. PLoS ONE 8(1): e54772. doi:10.1371/journal.pone.0054772

Editor: Nathan A. Ellis, University of Illinois at Chicago, United States of America

Received: October 2, 2012; **Accepted:** December 14, 2012; **Published:** January 25, 2013

Copyright: © 2013 Thompson et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: kConFab is supported by grants from the National Breast Cancer Foundation and the National Health and Medical Research Council (NHMRC) and by the Queensland Cancer Fund; the Cancer Councils of New South Wales, Victoria, Tasmania, and South Australia; and the Cancer Foundation of Western Australia. The kConFab Clinical Follow Up study was funded between 2001–2009 by NHMRC and currently by the National Breast Cancer Foundation and Cancer Australia (NBCF) (ID 628333). This study was supported by the Victorian Cancer Agency, Cancer Australia, and the NBCF (Priority-driven Collaborative Cancer Research Scheme grant, ID 628610), and the Wessex Cancer Trust. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ella.thompson@petermac.org

Introduction

RAD51 homolog D (*S. cerevisiae*) (*RAD51D/RAD51L3*; MIM#602954) is a component of the homologous recombination DNA repair pathway. The *RAD51D* protein forms a protein complex with *RAD51B*, *RAD51C* and *XRCC2* that binds to single stranded DNA (including single stranded gaps in double stranded DNA) and is required for the formation of *RAD51* foci in response to DNA damage [1,2]. Loveday et al [3] recently reported the identification of eight truncating mutations in *RAD51D* among 911 families with histories of breast and ovarian cancer, compared to one mutation among 1,060 population controls. They reported a significantly elevated risk of ovarian cancer (6.30, 95% CI 2.86–13.85) but did not detect a significantly elevated risk of breast cancer (1.32, 95% CI 0.59–2.96). They also reported that mutations are more prevalent in multiple case ovarian cancer families. *RAD51D* has subsequently been investigated in an additional series of 175 breast and ovarian cancer families, with an additional mutation being identified among the 51 families with at least two ovarian cancers (and among the 75 probands affected by ovarian cancer) [4]. Similarly, Pelttari et al [5] identified a splice site mutation (c.576+1G) in two breast cancer affected probands from 95 Finnish breast and/or ovarian

cancer families. Pelttari et al then screened for the c.576+1G variant in an additional 2,200 breast and 553 ovarian cancer patients and overall identified 5/707 patients with a personal or family history of ovarian cancer compared to 2/2,105 breast cancer only patients/families.

Until recently, *BRCA1* and *BRCA2* were the only genes known to confer a considerable risk of ovarian cancer (in conjunction with breast cancer) with two recent studies reporting that 13.3–14.1% of unselected high grade ovarian cancers are accounted for by mutations in one of these two genes [6,7]. A further small proportion of unselected cases carry mutations in *RAD51C* [8,9]. Loveday et al [3] estimated that 0.6% of unselected ovarian cancer cases will carry *RAD51D* mutations. To validate the association of *RAD51D* mutations to ovarian cancer and assess if there is any risk for breast cancer risk, we screened all coding exons in germline DNA from an unselected cohort of 245 unselected ovarian cancer patients and *BRCA1/2*-unrelated index cases from 1,060 breast and/or ovarian cancer families. Exons containing novel, non-synonymous variants among these cases were screened in a panel of 466 cancer-naïve control samples.

Table 1. Primers used for mutation analysis of *RAD51D*.

Primer pair	Forward primer (5'-3')	Reverse primer (5'-3')	Product size (bp)	Annealing temperature (°C)
<i>RAD51D</i> _ex1	CCGCGAATGCCACGTGA	AGGTATGCCAGGGCAGTG	228	62
<i>RAD51D</i> _ex2	GGGTAGAATTGACACCCATT	CTCCCAAAGTCTGGGATTA	264	62
<i>RAD51D</i> _ex3	GGTGAATGACACCCTGGGA	AGCATCAAAGCAGAGCTGAG	241	62
<i>RAD51D</i> _ex4	CAGAACCAGTGCTTAAAGAAA	CCCTGGGCTATGCATCTACC	248	62
<i>RAD51D</i> _ex5	GAATCTGGGCAAGTTTGGT	GGGGTTTTCTGTGCAGAA	266	62
<i>RAD51D</i> _ex6	TCTTCTTCTCAGCCTTACC	ATTGCACATCTGCATTCCA	277	62
<i>RAD51D</i> _ex7	TGTGCTAGAGGCTGACAGG	GCCAGAGACCAGACTCCAGA	215	62
<i>RAD51D</i> _ex8	CCAGCTCTGGAGTCTGGTCT	TTTGGGGTTCAGAAGCTGAC	239	62
<i>RAD51D</i> _ex9	CGATGCTCTATACTAGCA	CCTCCAGGGCCCAAGATT	261	62
<i>RAD51D</i> _ex10	GAGGCTGAAACCTTGAACCT	AGTGCCAGGTGGCAGTAAAC	253	62

doi:10.1371/journal.pone.0054772.t001

Materials and Methods

The unselected ovarian cancer cohort included 245 individuals with various histological subtypes of ovarian cancer (130 serous, 73 endometrioid, 35 mucinous, two clear cell, two granulosa cell tumours, two adenocarcinomas and one mixed mullerian tumour). These samples were obtained from patients presenting to hospitals

in the south of England, UK [10]. Undocumented, verbal consent was obtained from patients as approved by the governing ethics committee at the time.

The familial cohort included 540 individuals with verified personal and family histories of breast and/or ovarian cancer who were previously assessed at the Peter MacCallum Cancer Centre Familial Cancer Centre (Australia), as well as index cases from 520 multiple case breast cancer families (with or without ovarian cancer) obtained from the Kathleen Cunningham Foundation Consortium for Research into Familial Breast Cancer (kConFab) [11]. kConFab families are recruited through Familial Cancer Centres throughout Australia and New Zealand. All families were recruited based on multiple affected, multigenerational family and personal history of breast and/or ovarian cancer. The families fulfilled diagnostic criteria for BRCA testing, with no underlying *BRCA1* or *BRCA2* mutation having been identified. The ethnicity of the index cases was self-reported as Caucasian in the vast majority of cases. All individuals provided written, informed consent for genetic testing of the genetic causes of hereditary breast and ovarian cancer and subsequently tested negative for mutations in *BRCA1* and *BRCA2*. This study was approved by the Peter MacCallum Cancer Centre Human Research Ethics Committee. In total, index cases from 1,060 families were examined in this study, including 16 with a family history of ovarian cancer only, and 303 with a family history of both breast and ovarian cancer. Of these index cases, 98 had a personal history of ovarian cancer. The remaining 741 families had a personal and family history of breast cancer only.

Cancer-naive control DNA samples were obtained from kConFab (231 age- and ethnicity-matched best friend controls) and from the Princess Anne Hospital, UK (235 Caucasian female volunteers, as described previously) [12]. kConFab control individuals provided written, informed consent. Controls from the Princess Anne Hospital provided undocumented, verbal consent as approved by the governing ethics committee at the time.

DNA for mutation screening underwent whole genome amplification (WGA) using the Repli-G amplification system (Qiagen). Ten primer pairs were designed to amplify the ten coding exons of *RAD51D* with amplicons ranging in size from 215–277 bp for high resolution melt (HRM) analysis (Table 1). HRM analysis and DNA resequencing were performed as described previously [13]. Variant positions were determined with reference to GenBank reference sequence NM_002878.3. Nucle-




Figure 1. Loss of heterozygosity analysis of the **c.556C>T (p.(Arg186*))** variant. Sequencing (forward and reverse) of the heterozygous c.556C>T variant in the germline sample, and tumour DNA showing loss of the wildtype allele (with some contamination from normal DNA).
doi:10.1371/journal.pone.0054772.g001

Table 2. Summary of germline RAD51D variants.

Site ^a	Nucleotide change ^{b,c}	Protein ^b Change ^b	DBSNP (V135)	Previously described in RAD51D mutation studies?	1000 G MAF (%) ^d	EVS MAF (%) ^e	BC families (n = 741) ^f	BC/OC families (n = 303) ^f	OC families (n = 16) ^f	Unselected ovarian cases (n = 245)	Controls (n = 466) ^g	Grantham score ^h	Mutation Taster	MutPred	PMUT	POLYPHEN-2	SNPs&GO	
Truncating	Ex4 c.270_271dup	p.(Lys911lefs*13)	no	Yes [3]	-	-	-	-	-	-	1	-	-	-	-	-	-	-
	Ex6 c.556C>T	p.(Arg186*)	no	Yes [3,4]	-	-	-	-	1	-	-	-	-	-	-	-	-	-
	Ex9 c.803G>A	p.(Trp268*)	no	Yes [3]	-	-	-	-	1	-	-	-	-	-	-	-	-	-
Coding non-synonymous	Ex1 c.47T>C	p.(Met161Thr)	no	-	-	-	1	-	-	-	-	81	Polymorphism	0.587	Neutral	Benign	Neutral	
	Ex4 c.286G>T	p.(Gly96Cys)	no	-	-	-	1	-	-	-	-	159	Disease causing	0.938	Pathological	Probably damaging	Disease	
	Ex5 c.394G>A	p.(Val132Ile)	rs150498754	-	-	0.05	1	-	-	-	N/A	29	Disease causing	0.487	Neutral	Probably damaging	Neutral	
	Ex9 c.793G>A	p.(Gly265Arg)	rs140285068	Yes (control) [3,4]	-	0.02	1	-	-	-	-	125	Disease causing	0.924	Neutral	Probably damaging	Disease	
	Ex9 c.796C>T	p.(Arg266Cys)	no	-	-	-	1	-	-	-	-	180	Disease causing	0.621	Pathological	Probably damaging	Neutral	
	Ex2 c.117A>T	p.(=)	no	-	-	-	1	-	-	-	N/A	-	-	-	-	-	-	-
Coding synonymous	Ex9 c.864C>T	p.(=)	rs138557828	-	-	0.01	-	-	-	-	1	-	-	-	-	-	-	-
	Ex9 c.879G>A	p.(=)	no	-	-	0.01	1	-	-	-	-	-	-	-	-	-	-	-
	Ex9 c.900A>G	p.(=)	no	Yes [3]	-	-	-	-	-	-	1	-	-	-	-	-	-	-
	5'UTR c.-39C>T	p.(=)	no	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-
Non-coding	IVS1 c.82+60C>T	p.(=)	no	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-
	IVS1 c.83-20T>C	p.(=)	rs182793287	-	0.05	-	-	-	1	-	N/A	-	-	-	-	-	-	-
	IVS3 c.264-6C>T	p.(=)	no	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-

^aExon (Ex), intervening sequence (IVS).
^bVariant positions are reported in reference to NCBI RefSeq NM_002878.3 (mRNA) and NP_002869 (protein).
^cIn addition to the variants listed, common variants rs4796033, rs28563284 (non-synonymous) and rs9901455 (synonymous) were detected at high frequency in both cases and controls.
^dAll variants were queried against 1000 Genomes (1000 G) data using the 1000 Genomes Browser which integrates SNP and indel calls from 1,092 individuals (data release 20110521 v3). The minor allele frequency (MAF) is provided here.
^eAll variants were queried against Exome Variant Server (EVS), NHLBI Exome Sequencing Project (ESP). EVS contains SNP information from 5,379 individuals (data release ESP400). The minor allele frequency (MAF) is provided here.
^fBreast cancer only family (BC), ovarian cancer only family (OC), breast and ovarian cancer family (BC/OC).
^gvariants annotated with N/A were not assessed in control samples.
^h[22].
 doi:10.1371/journal.pone.0054772.t002

otide numbering reflects cDNA numbering with +1 corresponding to the A of the ATG translation initiation codon in the reference sequence, according to HGVS guidelines (www.hgvs.org/mutnomen). All novel variants were verified by Sanger resequencing of non-WGA DNA. Tumour cells were needle dissected from 10 µm sections to obtain tumour DNA, which was subsequently whole genome amplified.

The following *in silico* prediction tools were used to assess the likely functional effect of the missense variants identified in this study: PolyPhen-2 [14], SNPs&Go [15], MutPred [16], PMut [17] and MutationTaster [18]. Human Splicing Finder (HSF) was used to assess the effect of all non-truncating variants on splice sites [19].

Results and Discussion

Two previously reported truncating mutations, p.(Arg186*) and p.(Trp268*) were identified among a series of 245 unselected ovarian cancer patients (0.82%). The p.(Arg186*) variant was detected in a patient diagnosed with a grade 2 papillary serous cystadenocarcinoma at 66 years of age. DNA sequence analysis of tumour tissue obtained from this tumour showed reduction of the wildtype allele consistent with loss of heterozygosity (LOH) (Figure 1). The p.(Trp268*) variant was detected in a patient diagnosed with an endometrioid carcinoma (no grade information) at 70 years of age. No tumour tissue was available for LOH analysis. No family history information is available from either case. The histology of the two ovarian cases with truncating *RAD51D* mutations (*i.e.* high grade serous and endometrioid) is consistent with the majority of mutations reported in other *RAD51D* studies [3,4,5,20], and with other ovarian cancers associated with mutations in double strand break DNA repair genes (*e.g.* *BRCA1* and *BRCA2*), but the number of mutations in *RAD51D* identified to date is too few to determine the significance of this observation. A third truncating mutation, p.(Lys91Ilefs*13), was identified in one of 466 control samples (0.21%). All three of these mutations have previously been reported [3,4]. Table 2 provides a summary of all detected variants.

Analysis of 1,060 index cases from breast and/or ovarian cancer families did not identify any further truncating mutations. Interestingly, five rare (*i.e.* allele frequency <1%) nonsynonymous variants were detected, once each among 741 breast only cancer families. Three of these variants were novel: p.(Met16Thr), p.(Gly96Cys) and p.(Arg266Cys); the remaining two variants, rs150498754 and rs140285068, are reported in the Exome

Variant Server (EVS) database at frequencies of <0.02%. *In silico* analysis tools predicted that variants p.(Gly96Cys) and p.(Arg266Cys) would likely affect protein function (Table 2). Four of eight synonymous or intronic variants detected were novel; these were observed once each in either cases (c.117A>T, c.-39C>T, c.264-6C>T) or controls (c.82+60C>T). None of the synonymous or intronic variants were predicted to alter splicing.

The frequency of truncating germline *RAD51D* mutations detected in all patients with a personal history of ovarian cancer in this study (2/343 = 0.58%) is in keeping with that (0.6%) estimated by Loveday et al., and higher than observed in controls (0.21%). However, it is possible that the variant frequency reported here could be an underestimate due to the reduced sensitivity of HRM analysis compared to direct resequencing (used by Loveday et al.). Five of the variants detected in this study have previously been reported in *RAD51D* mutation studies by Loveday et al. or Osher et al. [3,4], and may represent founder mutations. However, there is no overlap with variants reported in more recent studies by Peltari et al. or Wickramanyake et al. [5,20]. To date, no truncating mutations have been detected among 1,092 individuals in the 1000 genomes cohort (data release 20110521 v3) [21] or 5,379 individuals in the Exome Variant Server (release ESP5400; NHLBI Exome Sequencing Project (ESP), Seattle, WA (<http://evs.gs.washington.edu/EVS/>) [June 2012]).

The absence of truncating mutations in 741 breast cancer only families (or 962 breast cancer-affected probands) provides further evidence that *RAD51D* mutations do not contribute significantly to breast cancer risk.

Acknowledgments

We wish to thank Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, Rebecca Driessen from the Victorian Familial Cancer Clinical Trials Group, staff of the Family Cancer Clinics, staff of the Peter MacCallum Cancer Centre Pathology Department and the Clinical Follow Up Study for their contributions to this resource, and the many families who contribute to kConFab.

Author Contributions

Conceived and designed the experiments: ET IC. Performed the experiments: SR. Analyzed the data: SR SS ET. Contributed reagents/materials/analysis tools: PJ GM AT DE kConFab. Wrote the paper: ET AT IC.

References

1. Masson J-Y, Tarsounas MC, Stasiak AZ, Stasiak A, Shah R, et al. (2001) Identification and purification of two distinct complexes containing the five RAD51 paralogs. *Genes & Development* 15: 3296–3307.
2. Smiraldi PG, Gruver AM, Osborn JC, Pittman DL (2005) Extensive Chromosomal Instability in Rad51d-Deficient Mouse Cells. *Cancer Research* 65: 2089–2096.
3. Loveday C, Turnbull C, Ramsay E, Hughes D, Ruark E, et al. (2011) Germline mutations in RAD51D confer susceptibility to ovarian cancer. *Nat Genet* 43: 879–882.
4. Osher DJ, De Leener K, Michils G, Hamel N, Tomiak E, et al. (2012) Mutation analysis of RAD51D in non-BRCA1/2 ovarian and breast cancer families. *Br J Cancer* 106: 1460–1463.
5. Peltari LM, Kiiski J, Nurminen R, Kallioniemi A, Schleutker J, et al. (2012) A Finnish founder mutation in RAD51D: analysis in breast, ovarian, prostate, and colorectal cancer. *J Med Genet*.
6. Zhang S, Royer R, Li S, McLaughlin JR, Rosen B, et al. (2011) Frequencies of BRCA1 and BRCA2 mutations among 1,342 unselected patients with invasive ovarian cancer. *Gynecol Oncol* 121: 353–357.
7. Alsop K, Fereday S, Meldrum C, deFazio A, Emmanuel C, et al. (2012) BRCA Mutation Frequency and Patterns of Treatment Response in BRCA Mutation-Positive Women With Ovarian Cancer: A Report From the Australian Ovarian Cancer Study Group. *Journal of Clinical Oncology*.
8. Peltari LM, Heikinen T, Thompson D, Kallioniemi A, Schleutker J, et al. (2011) RAD51C is a susceptibility gene for ovarian cancer. *Hum Mol Genet* 20: 3278–3288.
9. Thompson ER, Boyle SE, Johnson J, Ryland GL, Sawyer S, et al. (2011) Analysis of RAD51C Germline Mutations in High-Risk Breast and Ovarian Cancer Families and Ovarian Cancer Patients. *Hum Mutat* 33: 95.
10. Bryan EJ, Watson RH, Davis M, Hitchcock A, Foulkes WD, et al. (1996) Localization of an ovarian cancer tumor suppressor gene to a 0.5-cM region between D22S284 and CYP2D, on chromosome 22q. *Cancer Res* 56: 719–721.
11. Mann GJ, Thorne H, Balleine RL, Butow PN, Clarke CL, et al. (2006) Analysis of cancer risk and BRCA1 and BRCA2 mutation prevalence in the kConFab familial breast cancer resource. *Breast Cancer Res* 8: R12.
12. Baxter SW, Choong DY, Eccles DM, Campbell IG (2002) Transforming growth factor beta receptor 1 polyalanine polymorphism and exon 5 mutation analysis in breast and ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 11: 211–214.
13. Gorringer KL, Choong DY, Williams LH, Ramakrishna M, Sridhar A, et al. (2008) Mutation and methylation analysis of the chromodomain-helicase-DNA binding 5 gene in ovarian cancer. *Neoplasia* 10: 1253–1258.
14. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, et al. (2010) A method and server for predicting damaging missense mutations. *Nat Methods* 7: 248–249.

15. Calabrese R, Capriotti E, Fariselli P, Martelli PL, Casadio R (2009) Functional annotations improve the predictive score of human disease-related mutations in proteins. *Hum Mutat* 30: 1237–1244.
16. Li B, Krishnan VG, Mort ME, Xin F, Kamati KK, et al. (2009) Automated inference of molecular mechanisms of disease from amino acid substitutions. *Bioinformatics* 25: 2744–2750.
17. Ferrer-Costa C, Gelpi JL, Zamakola L, Parraga I, de la Cruz X, et al. (2005) PMUT: a web-based tool for the annotation of pathological mutations on proteins. *Bioinformatics* 21: 3176–3178.
18. Schwarz JM, Rodelsperger C, Schuelke M, Seelow D (2010) MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods* 7: 575–576.
19. Desmet FO, Hamroun D, Lalande M, Collod-Beroud G, Claustres M, et al. (2009) Human Splicing Finder: an online bioinformatics tool to predict splicing signals. *Nucleic Acids Res* 37: e67.
20. Wickramanyake A, Bernier G, Pennil C, Casadei S, Agnew KJ, et al. (2012) Loss of function germline mutations in RAD51D in women with ovarian carcinoma. *Gynecol Oncol* 127: 552–555.
21. The 1000 Genomes Project Consortium (2010) A map of human genome variation from population-scale sequencing. *Nature* 467: 1061–1073.
22. Grantham R (1974) Amino acid difference formula to help explain protein evolution. *Science* 185: 862–864.