

Elemental Concentrations in the Seed of Mutants and Natural Variants of *Arabidopsis thaliana* Grown under Varying Soil Conditions

Stephen C. McDowell¹, Garo Akmakjian², Chris Sladek¹, David Mendoza-Cozatl², Joe B. Morrissey³, Nick Saini¹, Ron Mittler⁶, Ivan Baxter⁷, David E. Salt⁴, John M. Ward⁵, Julian I. Schroeder², Mary Lou Guerinot³, Jeffrey F. Harper^{1*}

1 Department of Biochemistry and Molecular Biology, University of Nevada, Reno, Nevada, United States of America, **2** Division of Biological Sciences, University of California, San Diego, California, United States of America, **3** Department of Biological Sciences, Dartmouth College, Hanover, New Hampshire, United States of America, **4** Institute of Biological and Environmental Science, University of Aberdeen, Aberdeen, Scotland, United Kingdom, **5** Department of Plant Biology, University of Minnesota, St. Paul, Minnesota, United States of America, **6** Department of Biological Sciences, University of North Texas, Denton, Texas, United States of America, **7** United States Department of Agriculture—Agricultural Research Service, Plant Genetics Research Unit, Donald Danforth Plant Science Center, St. Louis, Missouri, United States of America

Abstract

The concentrations of mineral nutrients in seeds are critical to both the life cycle of plants as well as human nutrition. These concentrations are strongly influenced by soil conditions, as shown here by quantifying the concentration of 14 elements in seeds from *Arabidopsis thaliana* plants grown under four different soil conditions: standard, or modified with NaCl, heavy metals, or alkali. Each of the modified soils resulted in a unique change to the seed ionome (the mineral nutrient content of the seeds). To help identify the genetic networks regulating the seed ionome, changes in elemental concentrations were evaluated using mutants corresponding to 760 genes as well as 10 naturally occurring accessions. The frequency of ionic phenotypes supports an estimate that as much as 11% of the *A. thaliana* genome encodes proteins of functional relevance to ion homeostasis in seeds. A subset of mutants were analyzed with two independent alleles, providing five examples of genes important for regulation of the seed ionome: *SOS2*, *ABH1*, *CCC*, *At3g14280* and *CNGC2*. In a comparison of nine different accessions to a Col-0 reference, eight accessions were observed to have reproducible differences in elemental concentrations, seven of which were dependent on specific soil conditions. These results indicate that the *A. thaliana* seed ionome is distinct from the vegetative ionome, and that elemental analysis is a sensitive approach to identify genes controlling ion homeostasis, including those that regulate gene expression, phospho-regulation, and ion transport.

Citation: McDowell SC, Akmakjian G, Sladek C, Mendoza-Cozatl D, Morrissey JB, et al. (2013) Elemental Concentrations in the Seed of Mutants and Natural Variants of *Arabidopsis thaliana* Grown under Varying Soil Conditions. PLoS ONE 8(5): e63014. doi:10.1371/journal.pone.0063014

Editor: Els J M van Damme, Ghent University, Belgium

Received: November 14, 2012; **Accepted:** March 27, 2013; **Published:** May 6, 2013

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: The seed ionome mutant screen was supported by a grant from the National Science Foundation (<http://www.nsf.gov>, DBI-0077378) to MLG/JFH/JIS/JW. Grants to JFH from the National Institutes of Health (<http://www.nih.gov>, 1R01 GM070813-01) and the Department of Energy (<http://www.energy.gov>, DE-FG03-94ER20152) supported genetic studies on CNGCs/ACAs, and ALAs, respectively. Grants to JIS from the Department of Energy (<http://www.energy.gov>, DE-FG02-03ER15449) and the National Institutes of Health (<http://www.nih.gov>, P42 E5010337) in part supported sos mutant analysis and metal accession analysis. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: It should be noted that Ivan Baxter is a co-author of this manuscript and is also an editor for PLOS ONE. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: jfharper@unr.edu.

Introduction

Ion homeostasis is a complex process that is essential to the health and survival of all organisms. While at least 14 mineral nutrients are essential for higher plants [1], they can also be toxic if present in excess. Most land plants obtain their mineral nutrients from the soil, with homeostasis involving the regulation of uptake, binding, transport and sequestration; each of which is controlled by a network of genes. The resulting elemental composition of a specific tissue or cell has been referred to as the ionome [2–4].

Large scale screens to identify ionic mutants have been undertaken in yeast, *A. thaliana* and Lotus [5–8]. These studies have provided estimates that approximately 5% of the genome might be involved in ion homeostasis. Genetic variation controlling the leaf ionome has also been observed between naturally occurring populations of *A. thaliana* [9–17] and used in several

cases to identify genes of phenotypic consequence [18–23]. Ionic techniques have also been applied to crop plants including maize, rice, tomato and wheat [24–27].

Land plants have evolved the ability to adapt to highly variable soil conditions. Many studies have correlated different soil conditions with changes in elemental concentrations [14,17,28–30]. Soils that are deficient in specific nutrients, such as Fe and P, can sometimes be diagnosed by assaying the leaf ionome for a specific pattern of changes [31]. However, relatively little is known about the estimated 5% of the genome expected to regulate ion homeostasis, and even less about how genetic networks coordinate plant-environment interactions.

The purpose of this study was to explore different strategies to identify genes regulating ion homeostasis in seeds. While previous screens have uncovered mutations affecting the leaf ionome, this study was focused on the seed for two reasons. First, the elemental

concentrations of seed are expected to be strongly influenced by the physiological functions of the root and shoot; thereby providing a test that can potentially detect ionome defects throughout the plant [3]. Second, seed represent a primary source of mineral nutrition in the human diet, as well as a gateway for the entry of toxic elements into the food chain, and therefore an important target for biofortification or toxic element exclusion.

Here we report on a pilot study in which we quantified 14 elements in the seed of plants grown under four different soil conditions: standard, or modified with NaCl, heavy metals, or alkali. This survey included mutants corresponding to 760 genes, including membrane proteins, transcription factors, stress regulated genes, and unknowns. In addition, we included a comparison of 10 natural variants of *A. thaliana*. Our survey provides evidence that as much as 11% of the *A. thaliana* genome encodes proteins of functional relevance to the seed ionome. Here we identify seed ionome phenotypes for mutations in 15 different genes, five of which were corroborated with two independent mutant alleles. Phenotypes were also observed in eight of the nine accessions, in comparison to Col-0. The use of four different soil conditions revealed several conditional phenotypes, providing evidence that the seed ionome is highly sensitive to changes in the soil environment.

Results

Soil Modifications Affect Ion Homeostasis in Col-0 Seed

To evaluate the impact of growth conditions on the *A. thaliana* seed ionome, the concentration of 14 elements (Ba, Ca, Cd, Cu, Fe, K, Li, Mg, Mn, Mo, Na, P, S and Zn) was quantified in the seed of plants grown under four different soil conditions: standard, or modified with NaCl (75 mM NaCl), heavy metals (Al, As, Cd, Co, Ba, Cs, Cr, Cu, Ca, Fe, Mn, Mg, Li, Rb), or alkali (pH ~ 7.9 with CaO). Under standard conditions, the average elemental concentrations of Col-0 seed were comparable to previous reports [14]. Under heavy-metal growth conditions, most of the elements added to the soil accumulated in the seeds at concentrations below the lower limit of detection due to sensitivity limitations in the ICP-AES instrument used in this study. Each soil modification was chosen to challenge plants with a sub-optimal soil environment, while still allowing plants to grow and provide sufficient seed for ICP analyses. For example, Col-0 plants showed no consistent differences in plant health, bolt length, or seed set between the standard and 75 mM NaCl soil conditions. Furthermore, a quantitative analysis of Col-0 seed size found no statistically significant differences between standard and 75 mM NaCl soil conditions (Figure S1). Nevertheless, each of the three challenge conditions resulted in a unique perturbation of the Col-0 seed ionome (Figure 1). NaCl conditions increased relative concentrations of K, Li and Na; heavy metal conditions increased Ca, Cd, Cu, and S, but decreased Fe and Mn; and alkali conditions increased Ca, Cu, K, and S, but decreased Ba and Fe. This indicated that the chosen conditions were sufficient to alter ion homeostasis in the seed, supporting an expectation that these conditions could be used in the context of a mutant screen to increase the probability of identifying mutations that alter ion accumulation in the seed.

Plants were grown in trays of 32, with eight Col-0 control plants present in each tray. An experimental plant was considered to have a statistically significant change in elemental concentrations (i.e., an ionome phenotype) if one or more elements was different from the Col-0 control mean by three or more standard deviations (3 s.d.). Therefore, the sensitivity of phenotype detection was directly related to the variability of each specific element within

the Col-0 control seeds. Figure 2 shows the variability of each element in Col-0 control seeds under each soil condition, expressed in terms of the standard deviation relative to the mean (%RSD, percent relative standard deviation). While each tray was considered individually, the average variability of eight elements across the entire dataset (Ca, Cu, K, Mg, Mn, P, S, and Zn) was between 5–15%RSD for every soil condition. Higher variability (15–30%RSD) was observed for Ba, Cd, Li, Fe, Mo, and Na, resulting in lower sensitivity of phenotype detection. The highest variation was seen for Na (50%RSD) and Li (70%RSD), specifically in soils modified with NaCl and alkali, respectively.

Mutant Screen: T-DNA Insertion and Fast Neutron Deletion Lines

To identify genes that contribute to the regulation of the seed ionome, the elemental concentrations of mutant and Col-0 (i.e., wild-type) control seeds were compared under different soil conditions (File S1). Mutant lines representing 760 genes were chosen for analysis and included a nearly proportional representation from all gene ontology (GO) functional categories (Figure S2). The exception was transport-related genes, which were over-represented in our screen by approximately three-fold. The best candidates for disruption of gene functions were chosen when selecting mutant alleles for analysis. For example, all mutants used had T-DNA insertions located between the start and stop codons of the target gene. However, many of the lines had insertions located in introns, and no analyses were done to confirm the absence of corresponding full-length mRNA expression. In addition, most of the T-DNA insertion lines obtained from the stock center as homozygous mutants were used without reconfirming their zygosity. Thus, it is possible that some mutant phenotypes were missed in this survey due to leaky expression or mis-represented zygosity.

Three different labs screened subsets of mutants for the 760 genes, each using a different soil modification. The results were compiled for this report (File S1). 731 genes were screened in NaCl soil, 203 genes in heavy metal soil, 88 genes in alkali soil and 38 genes in untreated soil. Mutant lines were initially subjected to a first pass screen in which at least two plants were analyzed. The first pass screen identified 299 genes as candidates for having ionic phenotypes; revealing potential increases and decreases in every element detectable with our instrumentation. A follow up analysis of genes randomly selected from the 299 candidates yielded a 22% confirmation rate (File S1), suggesting that ~9% of the 760 total genes analyzed in this study are likely to have ionic phenotypes. An extension of this allows us to speculate that 7%–11% (at 95% confidence) of the *A. thaliana* genome encodes proteins of functional relevance to the seed ionome.

Ionic phenotypes were corroborated in mutant lines corresponding to 15 genes (Figure 3). The corroboration included meeting stringency criteria such as: reproducibility across multiple flats, soil conditions, or multiple alleles. Of these 15 genes, five were shown to have mutant phenotypes supported by two independent alleles (*ABH1*, *SOS2*, *At3g14280*, *CCC*, and *CNGC2*). Corroboration with two alleles provides strong evidence that the ionic phenotype is associated with a specific gene, as opposed to a possible second site mutation in some other gene. However, ten of the 15 genes with ionic phenotypes were only analyzed with a single allele (Figure 3), and still need to be verified by second alleles or transgene rescue experiments.

Accession Screen

To explore the extent of elemental variation between naturally occurring populations of *A. thaliana*, 10 accessions (Col-0, Cvi-1,




Figure 1. The elemental concentrations of Col-0 seed change under modified soil conditions. The average (\pm SE) concentration of each element is given in PPM and plotted on the y-axis on a log scale. In total, 139 flats were processed for NaCl soil, 92 for heavy metal soil, 35 for alkali soil and 37 for untreated soil. However, some elements were not detected in every flat. The numbers above the columns represent the number of flats from which data was collected (i.e., n-values). Circumstances for which no data was collected are represented by an “X”. * Elements for which a statistically significant difference between standard and modified soil conditions was observed (t-test, $p < 0.05$, Bonferroni multiple test correction). doi:10.1371/journal.pone.0063014.g001

Kas-1, Ler-0, Mrk-0, Nd-1, Sha, Ts-1, Tsu-1 and WS-0) were grown under standard, NaCl, heavy metal and alkali soil conditions. Compared to Col-0, the only accession that did not show an ionic difference under one or more soil conditions was Cvi-1. In contrast, the other eight accessions (Kas-1, Ler-0, Mrk-0, Nd-1, Sha, Ts-1, Tsu-1 and WS) showed a total of 13 ionic changes (Figure 4). For example, Kas-1 showed an increase in Fe, while Mrk-0 showed an increase in Zn. In the example of a Zn increase in Mrk-0, the increase was soil-dependent, as it was seen here in seed from plants grown under a NaCl challenge, but not with unmodified soil.

Discussion

To identify genes regulating ion homeostasis in seeds, this survey utilized four different soil conditions as a way to challenge plants to adjust their ion homeostasis set points, and thereby help reveal conditional phenotypes. Fourteen elements were quantified in an analysis of mutants corresponding to 760 genes, as well as the natural variation present in nine different accessions compared to Col-0.




Figure 2. Similar variability was found for 9 of the 14 elements under all soil conditions. Columns represent the average variability (\pm SE) of an element in Col-0 control seeds under a specific soil condition, expressed in terms of the standard deviation relative to the mean (%RSD, percent relative standard deviation). In total, 139 flats were processed for NaCl soil, 92 for heavy metal soil, 35 for alkali soil and 37 for untreated soil. However, some elements were not detected in every flat. The exact n-value for each circumstance can be found in Figure 1. Circumstances for which no data was collected are represented by an "X". * Elements for which a statistically significant difference between standard and modified soil conditions was observed (t-test, $p < 0.05$, Bonferroni multiple test correction). doi:10.1371/journal.pone.0063014.g002

Soil Modifications Alter Seed Ion Homeostasis

Two lines of evidence support a conclusion that the challenge conditions used in our assays altered ion homeostasis in *A. thaliana* seed. First, each challenge uniquely altered the concentration of elements in the seed of Col-0 control plants (Figure 1). The fact that changes were seen in elements other than those used to modify the soil supports an expectation that homeostasis is maintained by interconnected regulatory networks, and that each of the modified soil conditions used here was sufficient to perturb some of these connections. Second, in both the mutant and accession screens (Figures 3 and 4) many of the phenotypes were observed to be soil-specific. For example, while five elements showed significant changes in *abh1* seed obtained from plants grown under standard soil conditions, the only change that was consistent under all four soil conditions was a decrease in sulfur. Similarly, in the accession survey, soil-specific phenotypes were observed in seven of the eight accessions.

While each modified soil condition triggered distinct changes in the Col-0 seed ionome compared to standard soil, the steady-state concentration of most elements still appeared to be under relatively tight homeostasis controls. This was indicated by the observation of similar levels of variability for nine elements (Ba, Ca, K, Mg, Mn, Mo, P, S, and Zn) under standard and modified soil conditions (Figure 2). There were however, two notable exceptions: Na in NaCl soil and Li in alkali soil. As the sensitivity of detecting a phenotype in our screen is directly related to the elemental variability in control plants, our ability to detect Na phenotypes under NaCl challenge and Li phenotypes under alkali challenge was greatly diminished. These examples illustrate that, in some cases, the use of modified soil conditions can increase the variability of specific elements and thereby cause phenotypes to be missed that would have otherwise been detected as statistically significant under standard soil conditions.

Mutant Screen

An extrapolation from our screen of 760 genes supports an estimate that up to 11% (at 95% confidence) of the *A. thaliana* genome might be involved in the regulation of the seed ionome. This estimate is higher than those for *A. thaliana* leaf tissue (2–4%) [7] or for yeast (4.8%) [6]. It is possible that the 11% extrapolated from our results is an over-estimate. For example, 760 genes is still

only a small fraction of the genome, and some subgroup(s) of gene functions may be over-represented. However, the seed ionome is expected to be strongly influenced by root and shoot physiological functions as well as sink loading controls within the seed. It is therefore also possible that the seed ionome is dependent on more genes than other plant structures, or a single-cell system like yeast.

While the 760 genes include representatives of all GO annotations, transport-related genes were over-represented by approximately three-fold (Figure S2). Interestingly, this over-representation was actually in contrast to an under-representation of transport-related genes with ionic phenotypes (Figure S3). This under-representation is similar to that observed in a screen of 4,385 yeast mutants [6]. One possible explanation is that, because of a need to precisely regulate ion concentrations in many cell types, ion homeostasis systems might have evolved with functional redundancies provided by alternative transport pathways [6].

From a subset of genes chosen for further analyses, 15 were found to have reproducible mutant phenotypes (Figure 3). Of these, five were corroborated with two independent alleles and are discussed below. In two cases (ABH1 and CCC), the two independent sets of alleles showed overlapping but non-identical phenotypes, potentially due to different degrees of phenotypic penetrance. Regardless, these five genes provide examples of how ion homeostasis can be influenced by diverse sets of genes, including those involved in regulating gene expression, phosphoregulation, and ion transport. Additional mutations that warrant further consideration are listed in File S1, along with their corresponding ionome profiles and preliminary confidence rankings.

SOS2: Protein Kinase

An increase in the relative concentration of Na was observed in the seed of *sos2* mutants under all soil conditions tested, including unchallenged (Figure 3). In contrast, evidence from the Purdue Ionomics Information Management System (PiiMS, <http://www.ionomichub.org> [32]) (File S2) and a previous study [33] indicate that loss of SOS2 does not alter the leaf ionome under unchallenged conditions. *SOS2* encodes a protein kinase that is thought to be important for a plant's response to NaCl stress by activating transporters that can regulate the concentration of Na^+ in the cytosol [34–37]. Among the proteins activated by SOS2 are:

Gene Accession	Allele Accession	Soil Treatment	Significant Median % Change Compared to WT													n Plants	n Flats	Gene Name
			Ba	Ca	Cu	Fe	K	Li	Mg	Mn	Na	P	S	Zn				
At2g13540 ^a	Salk_016753	Untreated	-44	33			-15	15	31					-38	12	2	ABH1	
		NaCl		35										-45	14	3		
		Metal		26			-19							-23	16	4		
	Salk_024285	Untreated	-59	26			-16	17						-40	9	2		
		NaCl												-47	13	3		
		Metal												-13	5	2		
At5g35410 ^a	Salk_016683	Untreated											73	6	1	SOS2		
		NaCl											144	8	2			
		Metal											604	10	2			
	CS3863	Untreated											102	6	1			
		NaCl											880	3	1			
		Metal																
At3g14280 ^a	SALK_008085	Untreated												-22	12	3	Unknown	
		NaCl												-23	5	2		
	SALK_063934	Untreated												-27	12	3		
		NaCl												-28	6	2		
At1g30450 ^a	SALK_145300	Untreated	53	16			-40						-45	19	9	2	CCC	
	CS16311	Untreated		15	60	-47							-66	29	8	2		
At5g15410 ^a	SALK_019922	NaCl		48									-41		5	1	CNGC2	
	<i>cngc2-2^s</i>	NaCl		59									-18		2	1		
At1g17500, At1g72700 ^b	Salk_129561, Salk_049232	Untreated					-17								14	3	ALA4/5	
NaCl														-18	11	3		
At1g09700 ^b	SALK_064863	Untreated		30			-36	79					-45		8	2	HYL1	
		NaCl		68			-43	ND							2	1		
At2g29040 ^b	SALK_108349	Untreated			16										8	2	Unknown	
		NaCl			ND										2	1		
		Metal			20										7	2		
At3g50910 ^b	SALK_132810	Untreated		60			-62	-26							6	1	Unknown	
		NaCl		42			-48	-18				30			8	2		
		Alkali		40			-22	-18							6	1		
At3g22590 ^b	SALK_150644	Untreated		19			37								12	3	PHP	
		NaCl		33											6	2		
		Metal		65											4	1		
		Alkali		15											6	1		
At1g75820 ^b	WiscDsLox 489_492B1	Untreated					38							8	2	CLAVATA		
At1g04300 ^b	SAIL_701_C06	Untreated					-21								12	2	TRAF-Like	
		NaCl					-25								12	2		
At5g40780 ^b	SALK_036871	Untreated							23	686	-22				8	1	LHT1	
		NaCl							29	203	-25				15	3		
		Metal													12	3		
		Alkali							22			-15			6	1		
Not Identified ^b	FN 116.49	NaCl		34											6	1	Not Identified	
		Alkali		47											5	1		
At2g28670 ^b	FN145.01	Untreated		-14											1	1	ESB1	
		NaCl		-25											6	1		
		Metal		-45											6	1		
		Alkali		-11											5	1		

Figure 3. Percent changes in mutant elemental concentrations identify 15 genes that regulate the seed ionome. Data represent the median percent changes in the concentration of elements with reproducibly significant differences between the mutants and the Col-0 wild-type controls. An ionic change was considered significant if more than three standard deviations from the control mean and reproducible if consistent

across multiple flats, soil conditions, or alleles. Circumstances for which no data was collected are represented by "ND". ^a Genes with ionic phenotypes corroborated by two independent alleles ^b Candidate genes with putative ionic phenotypes evaluated with only one allele ^c T-DNA insertional mutant of CNGC2 [66]: ig# 581 and Harper Lab ss57.
doi:10.1371/journal.pone.0063014.g003

SOS1, a Na⁺/H⁺ antiporter that effluxes Na⁺ across the PM [38–40]; and NHX1, a Na⁺/H⁺ antiporter that sequesters Na⁺ in the vacuole [41]. The activity of the SOS1 Na⁺/H⁺ antiporter has been linked to long-distance Na⁺ transport in plants [42]. The observation that seeds from *sos2* plants over-accumulate Na⁺ relative to Col-0 wild-type is consistent with the involvement of the SOS2 kinase in either reducing the levels of Na⁺ translocated to

the seed, or as part of an efflux pathway to exclude Na from accumulating in developing seeds.

ABH1: Nuclear mRNA Cap Binding Protein

Multiple soil-dependent phenotypes were observed in the seed of *abh1* mutants, with a decrease in S being the only change that was consistent under all growth conditions tested (Figure 3). These changes might be specific to the seed, as no changes have yet been

Accession	Soil Treatment	Significant Median % Change Compared to WT										n Plants	n Flats
		Ba	Ca	Fe	K	Mg	Mn	Na	P	S	Zn		
KAS-1	Untreated	-44	-34	39								9	1
	Alkali	-32	-41	43			24					8	1
LER-0	Untreated											43	6
	NaCl							626				40	5
	Metal							108				2	1
	Alkali											6	1
MRK-0	Untreated											17	3
	NaCl										38	11	2
	Metal											22	4
	Alkali										12	6	1
ND-1	Untreated						73					18	3
	NaCl						48					10	2
	Metal											13	3
	Alkali											6	1
SHA	Untreated											18	3
	NaCl		-32				70					11	2
	Metal											16	3
	Alkali											6	1
TS-1	Untreated							79				19	4
	NaCl							219				4	2
	Metal							560				13	3
	Alkali							262				6	1
TSU-1	Untreated											22	4
	NaCl							295				13	3
	Metal							57				15	3
	Alkali											6	1
WS-0	Untreated						41					17	3
	NaCl						37					12	2
	Metal						24					19	4
	Alkali											6	1

Figure 4. Percent changes in elemental concentrations relative to Col-0 reveal soil-dependent differences in natural variants. Data represents the median percent change in the concentration of elements with reproducibly significant differences between the experimental accessions and the Col-0 control accession. An ionic change was considered significant if more than three standard deviations from the control mean and reproducible if consistent across multiple flats, soil conditions, or alleles.
doi:10.1371/journal.pone.0063014.g004

detected in assays of the leaf ionome (File S2). An *abh1* mutant plant was initially characterized as being hypersensitive to ABA [43,44]. ABH1 encodes the large subunit of the heterodimeric mRNA cap-binding complex (CBC) [43,44]. The eukaryotic CBC complex binds the 5' end of RNA polymerase II transcripts and has multiple roles in mRNA biogenesis, including nuclear export [45,46] and protecting transcripts from nuclease degradation [47]. Loss of ABH1 results in as many as 144 changes in gene expression of two-fold or greater in the leaf, including kinases, transporters and transcription factors (File S3) [48]. ABH1 has also been shown to be involved in alternate splicing [49], miRNA maturation [50–53], and possibly protein translation [54]. It is tempting to speculate that the changes in ion homeostasis observed in *abh1* seed are the direct result of altered expression and/or function of one or more genes involved seed in ion homeostasis. However, the possibility cannot be excluded that ionic changes in the *abh1* seed ionome are the indirect result of changes in whole-plant physiology caused by the ABA hypersensitivity of *abh1* mutants.

CCC: Cation Chloride Co-transporter

When grown with unmodified soil, *ccc* mutants were observed to produce seed that over-accumulate Ca and S, and under-accumulate Na and K (Figure 3). *CCC* encodes the only known cation chloride co-transporter in *A. thaliana*. Promoter-GUS [55] and public expression profiling analyses [56] suggest that *CCC* is expressed throughout the plant, including roots, shoots, seeds and pollen. In animals, cation chloride co-transporters are divided into three groups based on ion co-transport specificity: K^+Cl^- (KCC), Na^+Cl^- (NCC) and $Na^+K^+Cl^-$ (NKCC) [57]. In plants, two lines of evidence suggest a significant role for CCCs in cellular ion homeostasis. First, heterologous expression of CCC from *A. thaliana* in *Xenopus laevis* oocytes provides evidence of Cl^- ion transport in the presence of both Na^+ and K^+ , consistent with transport characteristics observed for the animal NKCC group [55]. Second, gene knockout and RNAi experiments in *A. thaliana* and *Oryza sativa*, respectively, indicate that loss of CCC can disrupt the ability of the plant to regulate the concentrations of Na, K and Cl, especially when grown under salt stress conditions [55,58]. Both knockout and RNAi silenced plants showed multiple growth defects, consistent with a model in which CCCs play a critical role in ion homeostasis in multiple cell types and tissues. The observation here that a *ccc* mutant over-accumulates Ca and S in seeds raises the question of whether the *in-vivo* ion transport properties of CCC may also include Ca^{2+} or SO_4^{2-} , or whether an increase in seed Ca and S could be indirect consequences of pleiotropic changes that affects ion loading or fluxes within the transpiration stream or nutritional loading into seeds.

At3g14280: Protein with Obscure Features

A decrease in seed S was observed in mutant plants with T-DNA insertions in the gene At3g14280 under all growth conditions tested (Figure 3). The protein encoded by gene At3g14280 is currently annotated as a protein with obscure features (POF) [59,60], with putative orthologs in monocots, dicots and mosses; none of which are annotated with known motifs, domains, or biological functions. The protein encoded by At3g14280 is predicted to be soluble [61], and public expression profiling data suggests it is expressed throughout the plant, including roots, shoots and pollen [56]. This highlights the potential of using a seed ionome screen to provide experimental evidence for annotating unknown genes with potential physiological functions. From our first pass screen of 760 genes, 45 “unknowns” were scored with potential phenotypes worthy of follow up analyses (File S1). In the case of the POF/unknown

encoded by At3g14280, we suggest its designation as a sulfur homeostasis gene (SHO), as two independent T-DNA insertions in its coding sequence result in a lower concentration of S in the *A. thaliana* seed.

CNGC2: Cyclic Nucleotide Gated Channel

An increase in Ca and a decrease in Mg were observed in the seed of *cngc2* mutants under NaCl challenge conditions (Figure 3). Whether these changes might be specific to the seed is not yet clear, but current evidence from the PiiMS database suggests that the leaf ionome is unaltered under an unchallenged soil condition (File S2). Cyclic nucleotide gated channels (CNGCs) are nonselective cation-conducting channels that can be activated by cyclic nucleotides. 20 CNGC isoforms (CNGC 1–20) have been identified in *A. thaliana* [62]. Electrophysiological experiments in *Xenopus laevis* oocytes, human embryonic kidney (HEK) cells and *A. thaliana* guard cells suggest that CNGC2 is permeable to cations including K^+ and Ca^{2+} , [63–65]. Loss of CNGC2 results in reduced plant health with a specific hypersensitivity to high Ca^{2+} levels (10 mM) [66,67]. While the possibility cannot be excluded that the changes in ion homeostasis for *cngc2* are an indirect consequence of reduced plant health, other CNGC isoforms have been implicated in the maintenance of ion homeostasis. For example, *cngc3* mutant seedlings under-accumulate K when grown on high levels of NaCl or KCl [68], *cngc1* mutant seedlings under-accumulate Ca [69], and antisense plants under-expressing CNGC10 under-accumulate Ca, Mg and K in shoot tissue [70]. Evidence from studies on CNGC18 are consistent with the potential of CNGCs to change the accumulation of Ca [71]. These examples of *cngc* mutants that under-accumulate Ca are in contrast to our finding that Ca is over-accumulated in *cngc2* seed. The observation that *cngc2* mutants also under-accumulate Mg in seeds is consistent with previous reports of a strong correlation between Ca and Mg concentrations [2], and raises the question of how Mg^{2+} transport might be influenced by CNGC2.

Accession Screen

Changes in the seed ionome, relative to Col-0, were observed for eight of the nine accessions studied. This is similar to what was seen for a seed and leaf ionome comparison in a previous study with 12 accessions grown with soil amended with subtoxic levels of various elements, including Cd and As [14]. However, our results using plants grown on four different soils indicate that most of the differences are likely to be soil-dependent.

Only one of the nine ionic changes observed in accessions screened under all four soil conditions occurred consistently under all soil conditions. Nevertheless, the observed high frequency of natural variation is consistent with the hypothesis that the seed and leaf ionomes are both sensitive indicators for changes in genetic networks that accompany adaptive changes in the evolution of plant accessions [72]. In the context of the leaf ionome, natural variation has already been used to identify several genes of ionic importance, for example: *MOT1*, Mo concentration [19]; *FPN2*, Co concentration [22]; *APR2*, sulfate concentration [20]; and *HKT1*, Na concentration [18,21,25,73].

Importantly, two accessions analyzed here show increased seed concentrations for Zn (Mrk-0) or Fe (Kas-1). As these two elements are considered important nutrients for biofortification of seeds, understanding the genetic basis of this natural variation might be instructive towards genetic improvements in crop plants. In the case of increased Zn in Mrk-0 seed, this change was only seen in plants grown under NaCl and alkali challenges. This suggests that understanding a NaCl or alkali stress response in Mrk-0 might

provide insights into genetic networks that regulate Zn homeostasis.

In the case of increased Fe in Kas-1 seed, this change was seen for both untreated and alkali soil conditions. A seed analysis was not possible for the NaCl or heavy metal challenges, as these plants were highly sensitive to the NaCl and heavy metal conditions used here, and produced very few seed. However, a previous study with a different amended soil treatment also showed an increase in seed Fe [14]. Interestingly, the previous study did not see any Fe changes in the leaf ionome. This suggests that the increased levels of Fe in Kas-1 seeds are not soil-dependent, and that the mechanism responsible has a seed specific feature.

Seed and Leaf Ionomes are Distinct

Previous studies have shown that the relative elemental compositions of root, leaf and seed tissues are very different [14,16,17,74]. For nine of the 15 seed ionome mutants identified here, an initial comparison with changes in the leaf ionome was possible through information available either from a public database (PiiMS) or the literature. With this subset of mutants, there were no examples in which elemental concentrations were altered in the same way (File S2). Similar results have emerged from analyses of three RIL populations of *A. thaliana* [17]. These examples support an expectation that the regulatory controls over the seed ionome are distinct from those that regulate the leaf, despite a dependence of the seed on vegetative tissues for their source of nutrients. Thus, in working towards a goal of biofortifying seeds with increased levels of mineral nutrients, specific attention will be required to understand the ion homeostasis controls that are specific to the seed ionome.

Conclusions

Here, we identified seed ionome functions for 15 genes, five of which were corroborated with two independent mutant alleles. Significantly, all 15 mutants showed at least one ionic change that could be identified in plants grown under standard conditions. This suggests that the sole use of a standard soil condition might provide the most efficient strategy for a first pass screen of the entire genome to identify genes that can impact the seed ionome. Nevertheless, our results show that modified soil conditions have significant influence on the seed ionome (Figure 1), and can be used to uncover soil-dependent ionic phenotypes in mutants (Figure 3) and naturally occurring accessions (Figure 4). Thus, a comprehensive understanding of the genetic networks governing ion homeostasis will eventually require careful attention to factors impacting the soil environment.

Materials and Methods

Resources

Seeds for the accessions [75], the SALK [76], SAIL [77] and WiscDsLox [78] T-DNA disruption lines and the CSHL [79] transposon disruption lines were obtained from the Arabidopsis Biological Resource Facility at Ohio State University. Seeds for SM [80] transposon disruption lines were obtained from the NASC European Arabidopsis Stock Centre. The locations of all inserts were estimated using flanking sequences available through the Salk SIGnAL T-DNA Express portal (<http://signal.salk.edu/cgi-bin/tdnaexpress>). Plant lines in the Wisconsin Collection [81] were isolated by individual labs from pooled seed and are available from the Harper Lab. Seeds designated as being from the Harper Lab Collection and all multiple knockout plant lines are also available from the Harper Lab. Seed lines with the prefix “FN” carry fast neutron deletions [7] and are available from the Salt

Lab. The *sos2-1* allele [37] carries a fast neutron deletion and is available from the Arabidopsis Biological Resource Facility at Ohio State University.

Untreated Soil Condition

Seeds were sown on plates containing ½ strength Murashige and Skoog basal salts, 0.05% MES (pH 5.7) and 1% agar. The seeds were stratified (4°C, dark) for 48 h and then grown at room temperature under 24 h light for 10 d. Seedlings were transplanted to soil (Sunshine LB-2 or Sunshine SMB283) and grown in a growth chamber at 21°C under long-day light conditions (16 h light/8 h dark) until bolting. After bolting, the plants were moved to a greenhouse where they were allowed to grow to maturity. The plants were frequently subirrigated by soaking the flats for ~16 h in an ebb-and-flow tray with an excess of 1/10 strength Hoaglands #2 basal salts supplemented with 5 µM Sprint138 Iron Chelate (Becker-Underwood, Ames, IA, USA).

75 mM NaCl Soil Condition

Plants were grown as described for the untreated soil condition until bolting. Upon bolting, plants were moved to a greenhouse and subirrigated by soaking the flats for ~16 h in an ebb-and-flow tray with an excess of 1/10 strength Hoaglands #2 supplemented with 5 µM Sprint138 Iron Chelate (Becker-Underwood, Ames, IA, USA) and 75 mM NaCl. This watering strategy was used to maintain a constant 75 mM NaCl challenge to the plants over the course of the experiment. Unless the NaCl concentration in the soil was allowed to equilibrate with an excess of fertilizer +75 mM NaCl, NaCl would accumulate in the soil resulting in a NaCl challenge far greater than 75 mM.

Heavy Metal Soil Condition

Seeds were sown on plates containing ¼ strength Murashige and Skoog with Gamborg’s vitamins, 10 mM MES (pH 5.6) and 1% phytoagar, and two week old seedlings were transplanted to soil. Heavy metal-containing soil was made by mixing 1 kg of soil (Sunshine Basic Mix 2) with 1 L heavy metal solution [0.005% AlCl₃, 0.005% KH₂AsO₄, 0.005% CdCl₂, 0.01% CoCl₂ and 0.0025% each of BaCl₂, CsNO₃, K₂CrO₄, CuSO₄, Ca(NO₃)₂, Fe(SO₄)₃, MnCl₂, MgCl₂, LiCl, RbCl and ZnSO₄ (w/v)], and the soil was split evenly among three trays. Trays were subirrigated thrice weekly with 700 ml of ¼ strength Hoagland’s #2 basal salts supplemented with 25 µg/L Sprint 330 Iron Chelate (Becker-Underwood, Ames, IA, USA).

Alkali Soil Condition

Seedlings were grown on plates containing Gamborg’s B5 media until they reached the 3–4 true leaf stage and were transplanted to high pH soil. High pH soil was prepared by supplementing Sunshine LB-2 soil with CaO (7.8g CaO/kg dry soil). The plants were subirrigated frequently with ¼ strength Hoagland’s #2 basal salts supplemented with 5 µM Sprint138 Iron Chelate (Becker-Underwood, Ames, IA, USA).

Elemental Concentration Measurements

Elemental analysis was performed using a Varian Vista PRO ICP-AES after a HNO₃ digestion. Samples were placed in 15 ml Falcon polypropylene centrifuge tubes (Fisher Scientific, Waltham, MA, USA) to which 0.5–1 ml of trace metal grade HNO₃ was added. The samples tubes were capped and allowed to react overnight followed by heating in an oven to 80°C for 2 h. After cooling, tubes were brought up to a volume of 3.5 ml in the case of a 0.5 ml HNO₃ digestion or 5 ml in the case of a 1 ml HNO₃

digestion. Analysis was conducted using conventional ICP-AES methods with the addition of a 0.5 ml Y +10 g CsNO₃/L solution introduced into the sample uptake for internal standardization.

Calculations

Sample weights were calculated using a method derived from that of Lahner et al. [7]. Briefly, sample weights were calculated by comparing the concentration (signal) of specific elements in experimental samples to weighed control samples. These calculations are based on the assumption that the signal/weight ratio (albeit different for every element) is consistent for experimental and control samples. In contrast to Lahner et al., [7], the subset of elements used in the weight calculations was not fixed, but rather elements were included if the variation in signal/weight ratio was 25%RSD or less. The concentrations of elements within mutant plants relative to the controls were considered in terms of z-Score (number of standard deviations from control mean) using:

Equation 1:

$$z - \text{Score} = \frac{x_{\text{sample}} - \bar{x}_{\text{control}}}{\sigma_{\text{control}}}$$

and in terms of percent change using:

Equation 2:

$$\% \text{ Change} = 100\% \times \frac{x_{\text{sample}} - \bar{x}_{\text{control}}}{\bar{x}_{\text{control}}}$$

Where x_{sample} is an individual sample, \bar{x}_{control} is the mean of control samples and σ_{control} is the standard deviation of the control samples. Data analysis was done in the Harper Lab using the PERL and R programming languages.

Supporting Information

Figure S1 75 mM NaCl soil modification does not alter Col-0 seed size. Average results (±SE) of four independent experiments are presented for untreated (open bars, n = 12 plants) and 75 mM NaCl (crosshatched bars, n = 12 plants) soil conditions. Seed sizes were measured from scanned images as two-dimensional areas using the ImageJ software package. No statistically significant differences were observed between untreated and 75 mM NaCl soil conditions (p = 0.7605, Welch’s t-test). (PDF)

Figure S2 Gene ontology analyses comparing the genes from the mutant screen to the *A. thaliana* genome. Gene Ontology (GO) analyses were done using gene ontology annotations from TAIR (<http://www.arabidopsis.org>). Three aspects of each gene product are considered individually: biological process, cellular component and molecular function. Each aspect is divided up into non-overlapping functional categories. Columns represent the percentage of genes within a functional category. The most significant difference between the genes analyzed in this study and the *A. thaliana* genome is an approximately three-fold increase in the number of transport-related genes in the Biological Process and Molecular Function categories. Eight of the 760 genes screened were not included in the GO analysis because they were represented by fast-neutron mutant lines and their identity is unknown. (PDF)

Figure S3 Gene ontology analyses comparing genes with potential phenotypes to genes without potential phenotypes. Analyses were done as described in Figure S2.

Genes were considered to have a potential ionic phenotype if they were identified as candidates in the first pass screen. Eight of the 760 genes screened were not included in the GO analysis because they were represented by fast-neutron mutant lines and their identity is unknown. Transport-related genes from both the Biological Process and Molecular Function GO categories were slightly less likely to show a putative phenotype. This suggests that their approximately three-fold over-representation in the mutant screen (Figure S2) did not produce a disproportionate increase in the estimate that 7–11% of the *A. thaliana* genome is involved in regulating seed ion homeostasis.

(PDF)

File S1 Cumulative results of trace element screen. The concentrations of elements in experimental plants relative to the Col-0 controls are represented in terms of z-Scores (number of standard deviations from the control mean). Data represents the median z-Scores of an experimental line within an individual flat. Boxes shaded in gray represent circumstances for which no data was collected. A change in elemental concentration is considered statistically significant if different from the control mean by three or more standard deviations (z-Score ≥3 or z-Score ≤ -3). Statistically significant changes in elemental concentrations are represented by blue (lower concentration) and yellow (higher concentration) shaded boxes. The phenotype of a line is described for each growth condition in one of four possible ways: reproducible, low confidence, inconsistent and none. A phenotype is considered “reproducible” if consistent within multiple trays of the same soil condition, across different soil conditions, or for independent alleles of the same gene. A phenotype is considered “low confidence” if observed in the first pass screen and not chosen for follow up analysis. A phenotype is considered “inconsistent” if observed in the first pass screen and not corroborated by a follow up analysis. A phenotype description of “none” indicates that no statistically significant changes in elemental concentrations were observed. IG#’s represent internal seed stock designations. Except in the case of double knockouts, common names and TAIR descriptions were obtained from the TAIR website (<http://www.arabidopsis.org>) and were current on 6/1/12.

(XLS)

File S2 Elemental abundances in vegetative tissue. The elemental contents of mutant plants relative to the tray average is represented in terms of z-Scores (number of standard deviations from the tray mean). Data represents the median z-Scores of an experimental line within an individual flat. Boxes shaded in gray represent circumstances for which no data was collected. A change in elemental accumulation is considered statistically significant if different from the tray average by three or more standard deviations (z-Score ≥3 or z-Score ≤ -3). Statistically significant over- and under-accumulations are represented by yellow and blue shaded boxes respectively. IG#’s represent internal seed stock designations. Data was obtained from the PiMS database.

(XLS)

File S3 Gene expression is altered in *abh1* mutant plants. Data represents the smallest fold change in gene expression between *abh1* and wild-type from two biological replicates. Microarray data used for this analysis is publicly available in the NCBI GEO database (<http://www.ncbi.nlm.nih.gov/projects/geo/>, GSE7112) and was originally published by Kuhn et al., 2008 [48].

(XLS)

Acknowledgments

We thank Sarah Ellison, Taylor Cohen and Scott Curran for plant cultivation and sample collection. Some seed stocks included in this study were generously provided by Catherine Chan and Joseph Ecker before being deposited in stock centers.

Author Contributions

Conceived and designed the experiments: SCM RM IB DS JW JIS MLG JFH. Performed the experiments: SCM GA CS DMC JBM NS JFH JW. Analyzed the data: SCM GA CS IB DS JW JIS MLG JFH. Contributed reagents/materials/analysis tools: SCM JW RM IB DS JIS MLG JFH. Wrote the paper: SCM GA JFH IB DS MLG JIS JBM JW.

References

- Marschner H (1995) Mineral Nutrition of Higher Plants. Ed 2. San Diego: Academic Press.
- Baxter I (2009) Ionomics: studying the social network of mineral nutrients. *Curr Opin Plant Biol* 12: 381–386.
- Baxter I (2010) Ionomics: The functional genomics of elements. *Briefings Funct Genomics* 9: 149–156.
- Salt DE, Baxter I, Lahner B (2008) Ionomics and the study of the plant ionome. *Annu Rev Plant Biol* 59: 709–733.
- Chen Z, Watanabe T, Shimano T, Okazaki K, Osaki M, et al. (2008) Rapid characterization of plant mutants with an altered ion-profile: a case study using *Lotus japonicus*. *New Phytol* 181: 795–801.
- Eide DJ, Clark S, Nair TM, Gehl M, Gribskov M, et al. (2005) Characterization of the yeast ionome: a genome-wide analysis of nutrient mineral and trace element homeostasis in *Saccharomyces cerevisiae*. *Genome Biol* 6: R77.
- Lahner B, Gong J, Mahmoudian M, Smith EL, Abid KB, et al. (2003) Genomic scale profiling of nutrient and trace elements in *Arabidopsis thaliana*. *Nat Biotechnol* 21: 1215–1221.
- Danku JMC, Gumaelius L, Baxter I, Salt DE (2009) A high-throughput method for *Saccharomyces cerevisiae* (yeast) ionomics. *J Anal At Spectrom* 24: 103.
- Kobayashi Y, Koyama H (2002) QTL analysis of Al tolerance in recombinant inbred lines of *Arabidopsis thaliana*. *Plant Cell Physiol* 43: 1526–1533.
- Bentsink L, Yuan K, Koornneef M, Vreugdenhil D (2003) The genetics of phytate and phosphate accumulation in seeds and leaves of *Arabidopsis thaliana*, using natural variation. *Theor Appl Genet* 106: 1234–1243.
- Hoekenga OA, Vision TJ, Shaff JE, Monforte AJ, Lee GP, et al. (2003) Identification and characterization of aluminum tolerance loci in *Arabidopsis* (*Landsberg erecta* x *Columbia*) by quantitative trait locus mapping. A physiologically simple but genetically complex trait. *Plant Physiol* 132: 936–948.
- Payne KA, Bowen HC, Hammond JP, Hampton CR, Lynn JR, et al. (2004) Natural genetic variation in caesium (Cs) accumulation by *Arabidopsis thaliana*. *New Phytol* 162: 535–548.
- Vreugdenhil D, Aarts MGM, Koornneef M, Nelissen H, Ernst WHO (2004) Natural variation and QTL analysis for cationic mineral content in seeds of *Arabidopsis thaliana*. *Plant, Cell Environ* 27: 828–839.
- Buescher E, Achberger T, Amusan I, Giannini A, Ochsensfeld C, et al. (2010) Natural genetic variation in selected populations of *Arabidopsis thaliana* is associated with ionic differences. *PLoS One* 5: e11081.
- Conn SJ, Berninger P, Broadley MR, Gilliham M (2012) Exploiting natural variation to uncover candidate genes that control element accumulation in *Arabidopsis thaliana*. *New Phytol* 193: 859–866.
- Baxter I, Hermans C, Lahner B, Yakubova E, Tikhonova M, et al. (2012) Biodiversity of mineral nutrient and trace element accumulation in *Arabidopsis thaliana*. *PLoS One* 7: e35121.
- Ghandilyan A, Ilk N, Hanhart C, Mbengue M, Barboza L, et al. (2009) A strong effect of growth medium and organ type on the identification of QTLs for phytate and mineral concentrations in three *Arabidopsis thaliana* RIL populations. *J Exp Bot* 60: 1409–1425.
- Baxter I, Brazelton JN, Yu D, Huang YS, Lahner B, et al. (2010) A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter *AtHKT1;1*. *PLoS Genet* 6: e1001193.
- Baxter I, Muthukumar B, Park HC, Buchner P, Lahner B, et al. (2008) Variation in molybdenum content across broadly distributed populations of *Arabidopsis thaliana* is controlled by a mitochondrial molybdenum transporter (*MOT1*). *PLoS Genet* 4: e1000004.
- Loudet O, Saliba-Colombani V, Camilleri C, Calenge F, Gaudon V, et al. (2007) Natural variation for sulfate content in *Arabidopsis thaliana* is highly controlled by *APR2*. *Nat Genet* 39: 896–900.
- Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, et al. (2006) Natural variants of *AtHKT1* enhance Na⁺ accumulation in two wild populations of *Arabidopsis*. *PLoS Genet* 2: e210.
- Morrissey J, Baxter IR, Lee J, Li L, Lahner B, et al. (2009) The ferroportin metal efflux proteins function in iron and cobalt homeostasis in *Arabidopsis*. *Plant Cell* 21: 3326–3338.
- Tian H, Baxter IR, Lahner B, Reinders A, Salt DE, et al. (2010) *Arabidopsis* *NPCC6/NaKR1* is a phloem mobile metal binding protein necessary for phloem function and root meristem maintenance. *Plant Cell* 22: 3963–3979.
- Garcia-Oliveira AL, Tan L, Fu Y, Sun C (2009) Genetic identification of quantitative trait loci for contents of mineral nutrients in rice grain. *J Integr Plant Biol* 51: 84–92.
- Munns R, James RA, Xu B, Athman A, Conn SJ, et al. (2012) Wheat grain yield on saline soils is improved by an ancestral Na⁺ transporter gene. *Nat Biotechnol* 30: 360–364.
- Sánchez-Rodríguez E, Rubio-Wilhelmi M, Cervilla LM, Blasco B, Rios JJ, et al. (2010) Study of the ionome and uptake fluxes in cherry tomato plants under moderate water stress conditions. *Plant Soil* 335: 339–347.
- Sorić R, Ledencan T, Zdunić Z, Jambrović A, Brkić I, et al. (2011) Quantitative trait loci for metal accumulation in maize leaf. *Maydica* 56: 323–329.
- Ernst WH, Nelissen HJ, Ten Bookum WM (2000) Combination toxicology of metal-enriched soils: physiological responses of a Zn- and Cd-resistant ecotype of *Silene vulgaris* on polymetallic soils. *Environ Exp Bot* 43: 55–71.
- Ghandilyan A, Barboza L, Tisé S, Granier C, Reymond M, et al. (2009) Genetic analysis identifies quantitative trait loci controlling rosette mineral concentrations in *Arabidopsis thaliana* under drought. *New Phytol* 184: 180–192.
- Prinzenberg AE, Barbier H, Salt DE, Stich B, Reymond M (2010) Relationships between growth, growth response to nutrient supply, and ion content using a recombinant inbred line population in *Arabidopsis*. *Plant Physiol* 154: 1361–1371.
- Baxter IR, Vitek O, Lahner B, Muthukumar B, Borghi M, et al. (2008) The leaf ionome as a multivariable system to detect a plant's physiological status. *Proc Natl Acad Sci U S A* 105: 12081–12086.
- Baxter I, Ouzzani M, Orcun S, Kennedy B, Jandhyala SS, et al. (2007) Purdue ionomics information management system. An integrated functional genomics platform. *Plant Physiol* 143: 600–611.
- Zhu JK, Liu J, Xiong L (1998) Genetic analysis of salt tolerance in *Arabidopsis*. Evidence for a critical role of potassium nutrition. *Plant Cell* 10: 1181–1191.
- Huang G-T, Ma S-L, Bai L-P, Zhang L, Ma H, et al. (2012) Signal transduction during cold, salt, and drought stresses in plants. *Mol Biol Rep* 39: 969–987.
- Zhu JK (2003) Regulation of ion homeostasis under salt stress. *Curr Opin Plant Biol* 6: 441–445.
- Mahajan S, Pandey GK, Tuteja N (2008) Calcium- and salt-stress signaling in plants: shedding light on SOS pathway. *Arch Biochem Biophys* 471: 146–158.
- Liu J, Ishitani M, Halfter U, Kim CS, Zhu JK (2000) The *Arabidopsis thaliana* *SOS2* gene encodes a protein kinase that is required for salt tolerance. *Proc Natl Acad Sci U S A* 97: 3730–3734.
- Quintero FJ, Ohta M, Shi H, Zhu J-K, Pardo JM (2002) Reconstitution in yeast of the *Arabidopsis* SOS signaling pathway for Na⁺ homeostasis. *Proc Natl Acad Sci U S A* 99: 9061–9066.
- Qiu Q-S, Guo Y, Dietrich MA, Schumaker KS, Zhu J-K (2002) Regulation of *SOS1*, a plasma membrane Na⁺/H⁺ exchanger in *Arabidopsis thaliana*, by *SOS2* and *SOS3*. *Proc Natl Acad Sci U S A* 99: 8436–8441.
- Shi H, Ishitani M, Kim C, Zhu JK (2000) The *Arabidopsis thaliana* salt tolerance gene *SOS1* encodes a putative Na⁺/H⁺ antiporter. *Proc Natl Acad Sci U S A* 97: 6896–6901.
- Qiu Q-S, Guo Y, Quintero FJ, Pardo JM, Schumaker KS, et al. (2004) Regulation of vacuolar Na⁺/H⁺ exchange in *Arabidopsis thaliana* by the salt-overly-sensitive (*SOS*) pathway. *J Biol Chem* 279: 207–215.
- Shi H, Quintero FJ, Pardo JM, Zhu J-K (2002) The putative plasma membrane Na⁺/H⁺ antiporter *SOS1* controls long-distance Na⁺ transport in plants. *Plant Cell* 14: 465–477.
- Hugouvieux V, Kwak JM, Schroeder JI (2001) An mRNA cap binding protein, *ABH1*, modulates early abscisic acid signal transduction in *Arabidopsis*. *Cell* 106: 477–487.
- Hugouvieux V, Murata Y, Young JJ, Kwak JM, Mackesy DZ, et al. (2002) Localization, ion channel regulation, and genetic interactions during abscisic acid signaling of the nuclear mRNA cap-binding protein, *ABH1*. *Plant Physiol* 130: 1276–1287.
- Izaurrealde E, Stepinski J, Darzynkiewicz E, Mattaj IW (1992) A cap binding protein that may mediate nuclear export of RNA polymerase II-transcribed RNAs. *J Cell Biol* 118: 1287–1295.
- Hamm J, Mattaj IW (1990) Monomethylated cap structures facilitate RNA export from the nucleus. *Cell* 63: 109–118.
- Balatsos NAA, Nilsson P, Mazza C, Cusack S, Virtanen A (2006) Inhibition of mRNA deadenylation by the nuclear cap binding complex (CBC). *J Biol Chem* 281: 4517–4522.
- Kuhn JM, Hugouvieux V, Schroeder JI (2007) mRNA cap binding proteins: effects on abscisic acid signal transduction, mRNA processing, and microarray analyses. *Curr Top Microbiol Immunol* 326: 139–150.
- Raczynska KD, Simpson CG, Ciesiolka A, Szew L, Lewandowska D, et al. (2010) Involvement of the nuclear cap-binding protein complex in alternative splicing in *Arabidopsis thaliana*. *Nucleic Acids Res* 38: 265–278.
- Chen X (2008) A silencing safeguard: links between RNA silencing and mRNA processing in *Arabidopsis*. *Dev Cell* 14: 811–812.

51. Gregory BD, O'Malley RC, Lister R, Urich MA, Tonti-Filippini J, et al. (2008) A link between RNA metabolism and silencing affecting Arabidopsis development. *Dev Cell* 14: 854–866.
52. Laubinger S, Sachsenberg T, Zeller G, Busch W, Lohmann JU, et al. (2008) Dual roles of the nuclear cap-binding complex and SERRATE in pre-mRNA splicing and microRNA processing in Arabidopsis thaliana. *Proc Natl Acad Sci U S A* 105: 8795–8800.
53. Kim S, Yang J-Y, Xu J, Jang I-C, Prigge MJ, et al. (2008) Two cap-binding proteins CBP20 and CBP80 are involved in processing primary MicroRNAs. *Plant Cell Physiol* 49: 1634–1644.
54. Bush MS, Hutchins AP, Jones AME, Naldrett MJ, Jarmolowski A, et al. (2009) Selective recruitment of proteins to 5' cap complexes during the growth cycle in Arabidopsis. *Plant J* 59: 400–412.
55. Colmenero-Flores JM, Martínez G, Gamba G, Vázquez N, Iglesias DJ, et al. (2007) Identification and functional characterization of cation-chloride cotransporters in plants. *Plant J* 50: 278–292.
56. Grennan AK (2006) Genevestigator. Facilitating web-based gene-expression analysis. *Plant Physiol* 141: 1164–1166.
57. Russell JM (2000) Sodium-potassium-chloride cotransport. *Physiol Rev* 80: 211–276.
58. Kong X-Q, Gao X-H, Sun W, An J, Zhao Y-X, et al. (2011) Cloning and functional characterization of a cation-chloride cotransporter gene OsCCC1. *Plant Mol Biol* 75: 567–578.
59. Gollery M, Harper J, Cushman J, Mittler T, Girke T, et al. (2006) What makes species unique? The contribution of proteins with obscure features. *Genome Biol* 7: R57.
60. Gollery M, Harper J, Cushman J, Mittler T, Mittler R (2007) POFs: what we don't know can hurt us. *Trends Plant Sci* 12: 492–496.
61. Krogh A, Larsson B, Von Heijne G, Sonnhammer EL (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 305: 567–580.
62. Mäser P, Thomine S, Schroeder JI, Ward JM, Hirschi K, et al. (2001) Phylogenetic relationships within cation transporter families of Arabidopsis. *Plant Physiol* 126: 1646–1667.
63. Leng Q, Mercier RW, Yao W, Berkowitz GA (1999) Cloning and first functional characterization of a plant cyclic nucleotide-gated cation channel. *Plant Physiol* 121: 753–761.
64. Leng Q, Mercier RW, Hua B-G, Fromm H, Berkowitz GA (2002) Electrophysiological analysis of cloned cyclic nucleotide-gated ion channels. *Plant Physiol* 128: 400–410.
65. Ali R, Ma W, Lemtiri-Chlieh F, Tsalts D, Leng Q, et al. (2007) Death don't have no mercy and neither does calcium: Arabidopsis CYCLIC NUCLEOTIDE GATED CHANNEL2 and innate immunity. *Plant Cell* 19: 1081–1095.
66. Chan CWM, Schorrak LM, Smith RK, Bent AF, Sussman MR (2003) A cyclic nucleotide-gated ion channel, CNGC2, is crucial for plant development and adaptation to calcium stress. *Plant Physiol* 132: 728–731.
67. Clough SJ, Fengler KA, Yu IC, Lippok B, Smith RK, et al. (2000) The Arabidopsis *dnd1* “defense, no death” gene encodes a mutated cyclic nucleotide-gated ion channel. *Proc Natl Acad Sci U S A* 97: 9323–9328.
68. Gobert A, Park G, Amtmann A, Sanders D, Maathuis FJM (2006) Arabidopsis thaliana cyclic nucleotide gated channel 3 forms a non-selective ion transporter involved in germination and cation transport. *J Exp Bot* 57: 791–800.
69. Ma W, Ali R, Berkowitz GA (2006) Characterization of plant phenotypes associated with loss-of-function of AtCNGC1, a plant cyclic nucleotide gated cation channel. *Plant Physiol Biochem* 44: 494–505.
70. Guo KM, Babourina O, Christopher DA, Borsic T, Rengel Z (2010) The cyclic nucleotide-gated channel AtCNGC10 transports Ca²⁺ and Mg²⁺ in Arabidopsis. *Physiol Plant* 139: 303–312.
71. Frietsch S, Wang Y-F, Sladek C, Poulsen LR, Romanowsky SM, et al. (2007) A cyclic nucleotide-gated channel is essential for polarized tip growth of pollen. *Proc Natl Acad Sci U S A* 104: 14531–14536.
72. Baxter I, Dilkes BP (2012) Elemental profiles reflect plant adaptations to the environment. *Science* 336: 1661–1663.
73. Mäser P, Eckelman B, Vaidyanathan R, Horie T, Fairbairn DJ, et al. (2002) Altered shoot/root Na⁺ distribution and bifurcating salt sensitivity in Arabidopsis by genetic disruption of the Na⁺ transporter AtHKT1. *FEBS Lett* 531: 157–161.
74. Salt DE (2004) Update on plant ionomics. *Plant Physiol* 136: 2451–2456.
75. Nordborg M, Hu TT, Ishino Y, Jhaveri J, Toomajian C, et al. (2005) The pattern of polymorphism in Arabidopsis thaliana. *PLoS Biol* 3: e196.
76. Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, et al. (2003) Genome-wide insertional mutagenesis of Arabidopsis thaliana. *Science* 301: 653–657.
77. Sessions A, Burke E, Presting G, Aux G, Mcelver J, et al. (2002) A high-throughput Arabidopsis reverse genetics system. *Plant Cell* 14: 2985–2994.
78. Woody ST, Austin-Phillips S, Amasino RM, Krysan PJ (2007) The WiscDsLox T-DNA collection: an Arabidopsis community resource generated by using an improved high-throughput T-DNA sequencing pipeline. *J Plant Res* 120: 157–165.
79. Sundaresan V, Springer P, Volpe T, Haward S, Jones JD, et al. (1995) Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements. *Genes Dev* 9: 1797–1810.
80. Tissier AF, Marillonnet S, Klimyuk V, Patel K, Torres MA, et al. (1999) Multiple independent defective suppressor-mutator transposon insertions in Arabidopsis: a tool for functional genomics. *Plant Cell* 11: 1841–1852.
81. Krysan PJ, Young JC, Tax F, Sussman MR (1996) Identification of transferred DNA insertions within Arabidopsis genes involved in signal transduction and ion transport. *Proc Natl Acad Sci U S A* 93: 8145–8150.