

Mice with Deficient BK Channel Function Show Impaired Prepulse Inhibition and Spatial Learning, but Normal Working and Spatial Reference Memory

Marei Typlt¹✉, Magdalena Mirkowski¹✉, Erin Azzopardi¹, Lukas Ruettiger², Peter Ruth³, Susanne Schmid¹*

1 Department of Anatomy and Cell Biology, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, **2** Hearing Research Center, Universität Tübingen, Tübingen, Germany, **3** Pharmakologie, Toxikologie und Klinische Pharmazie, Institut für Pharmazie, Universität Tübingen, Tübingen, Germany

Abstract

Genetic variations in the large-conductance, voltage- and calcium activated potassium channels (BK channels) have been recently implicated in mental retardation, autism and schizophrenia which all come along with severe cognitive impairments. In the present study we investigate the effects of functional BK channel deletion on cognition using a genetic mouse model with a knock-out of the gene for the pore forming α -subunit of the channel. We tested the F1 generation of a hybrid SV129/C57BL6 mouse line in which the *slc1* gene was deleted in both parent strains. We first evaluated hearing and motor function to establish the suitability of this model for cognitive testing. Auditory brain stem responses to click stimuli showed no threshold differences between knockout mice and their wild-type littermates. Despite of muscular tremor, reduced grip force, and impaired gait, knockout mice exhibited normal locomotion. These findings allowed for testing of sensorimotor gating using the acoustic startle reflex, as well as of working memory, spatial learning and memory in the Y-maze and the Morris water maze, respectively. Prepulse inhibition on the first day of testing was normal, but the knockout mice did not improve over the days of testing as their wild-type littermates did. Spontaneous alternation in the y-maze was normal as well, suggesting that the BK channel knock-out does not impair working memory. In the Morris water maze knock-out mice showed significantly slower acquisition of the task, but normal storage memory once the task was learned. Thus, we propose a crucial role of the BK channels in learning, but not in memory storage or recollection.

Citation: Typlt M, Mirkowski M, Azzopardi E, Ruettiger L, Ruth P, et al. (2013) Mice with Deficient BK Channel Function Show Impaired Prepulse Inhibition and Spatial Learning, but Normal Working and Spatial Reference Memory. PLoS ONE 8(11): e81270. doi:10.1371/journal.pone.0081270

Editor: Valentin Ceña, Universidad de Castilla-La Mancha, Spain

Received: June 10, 2013; **Accepted:** October 10, 2013; **Published:** November 26, 2013

Copyright: © 2013 Typlt et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Canadian Institute for Health Research (CIHR) MOP#114870 to S. Schmid Deutsche Forschungsgemeinschaft (DFG) scholarship to M.Typlt Ontario Ministry of Training, Colleges and Universities scholarship to E. Azzopardi. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Susanne.Schmid@schulich.uwo.ca

✉ These authors contributed equally to this work.

Introduction

The large-conductance, voltage- and calcium-activated potassium channels (BK channels) are expressed throughout the mammalian nervous system [1–3] and can be found at neuronal soma and processes, as well as on presynaptic terminals [1,4,5]. Activated by membrane depolarization and/or micromolar concentrations of intracellular calcium, BK channels drive the membrane potential towards the potassium equilibrium potential and therefore re- and hyperpolarize the neuron [6–8]. By limiting the duration of action potentials, they regulate the general excitability of neurons [9–11], as well as the transmitter release at presynaptic terminals [8,12–15].

Over the last several years evidence has accumulated that alterations in the expression and function of BK channels are involved in a number of mental disorders. So far, functional alterations of these channels have been linked to mental retardation [16,17], autism [18], and schizophrenia (for review see [19]) which all involve sensorimotor, as well as cognitive impairments.

In order to explore the role of BK channels in cognitive processes we tested a mouse model lacking functional BK channels. Previous studies found that BK channel knock-out mice with a 129SVJ inbred background have impaired hearing [20–22] and motor function [23], which would impact most of the standard behavioral tests. However, another study in mice with a FVB/NJ inbred background suggests strain specific differences in the hearing ability of BK channel knock-out mice [24]. We here used the F1 generation of a hybrid SV129/C57BL6 background. Several studies indicate that using a hybrid or mixed background in F1 or F2 generation is most effective to contrast the gene knock-out to natural occurring variations in behaviour due to genetic background and environmental factors [25–27].

After confirming normal hearing and locomotion, we first tested sensorimotor gating. Sensorimotor gating is a pre-attentive process that filters sensory information for salient stimuli within the environmental noise. In all patient groups linked with an alteration in BK channel expression disruptions in sensorimotor gating have been described (fragile X syndrome: [28], autism: [29,30], and schizophrenia: [31–33]). Therefore, we hypothesized that senso-

rimotor gating will be impaired in the BK channel knockout mice. We further tested the BK knockout mice in the Y-maze to assess their working memory, as well as in the Morris water maze to assess hippocampus dependent learning and memory. Since BK channels are abundantly expressed in the hippocampus [2,3,34], we expected to find deficits specifically in these tasks.

Methods

Animals and animal care

We used mice of the F1 generation of a hybrid SV129/C57BL6 line with deficient BK channel function. BK channel function was abolished by deleting the *slol* gene which encodes the pore forming channel protein (α -subunit; for details see [23]). Heterozygous C57BL6 mice were paired with heterozygous SV129 mice. Exclusively mice of the respective F1 generation were tested in order to avoid effects of inbreeding. Also, preliminary data had suggested that these hybrid mice do not show hearing deficits as previously described in other genetic models with impaired BK channel function [20,21]. In total, we tested 20 wild-type (WT; 10/10 male/female), 17 heterozygote ($BK\alpha^{+/-}$; 10/7), and 19 homozygote knock-out mice ($BK\alpha^{-/-}$; 11/8), but different, randomly selected, subsets of these animals were used for the single tests. The hearing test was performed on a separate set of animals (for details see respective method sections). The animals were litter- and/or age matched. We only tested mice at ages from 3 to 7 months to avoid effects of aging.

All mice were generated and genotyped at the Pharmaceutical Institute, University of Tübingen, Germany. In order to ensure proper tracking each mouse got an individual ear-tag when collecting the DNA sample for the genotyping. The mice were shipped to Canada at the age of 1.5-3 months and subsequently allowed to acclimate for 2 weeks before behavioural testing started. Mice were group housed with mixed genetic background within groups, with a 12 hour light-dark cycle and with ad libitum food and water. Testing only occurred during the light cycle from 7am to 7pm. If animals were tested repeatedly over several days, the time of testing for each animal was kept constant.

All procedures were in accordance with the ethical guidelines of the Canadian Council on Animal Care (CCAC) and approved by the University of Western Ontario Animal Use Subcommittee.

Hearing measurements/auditory brainstem responses (ABR)

Hearing measurements were conducted at the University of Tübingen, Germany, and were in accordance with institutional and national guidelines following approval by the University of Tübingen, Veterinary Care Unit, and the Animal Care and Ethics Committee of the regional board of the Federal State Government of Baden-Württemberg, Germany.

Auditory brainstem responses (ABR) were recorded from 10 WT (5/5 male/female) and 8 $BK\alpha^{-/-}$ mice (6/2), as described previously [35]. Each mouse was tested at several time points between the age of 4–15 weeks (9 ± 3.5 weeks) to evaluate the development of the hearing thresholds. Briefly, recordings were performed while mice were under ketamine (75 mg/kg body weight) and xylazine (5 mg/kg body weight) anesthesia. ABRs to free field click (100 μ s), and pure tone (3 ms, 1 ms ramp) acoustic stimuli were recorded using sub-dermal silver wire electrodes at the ear (active), the vertex (reference) and the back (ground) of the animals. Signals were amplified (50-100-fold), band-pass filtered (200 Hz high-pass and 5 kHz low-pass), and averaged for 64-256 repetitions at each sound pressure level (SPL) presented (0-100 dB

in steps of 5 dB). Hearing threshold was defined as the lowest SPL that produced a potential visually distinct from background noise.

Grip force test

In order to assess muscular strength, the grip force test was used. The front limb strength of 16 animals of each genotype (WT: 9 males/ 7 females, $BK\alpha^{+/-}$: 10/6, $BK\alpha^{-/-}$: 9/7) was measured in 5 trials per animal. The animals were between 7 and 9 months (34 ± 2 weeks) old at the time of testing. A force gauge (Grip Strength Meter, Columbus Instruments, Columbus, Ohio, USA) was connected to a metal grid which allowed the animal to hold on. The animal was held by the base of its tail and lowered towards the grid until it gripped it with both front limbs. Then the animal was gently pulled back until it lost the grip. Between trials the animals were allowed to recover in their home cages for about 10 minutes. For subsequent analyses the results from the 5 trials were averaged for each animal.

Cat walk

To evaluate gait 11 WT (6 males/5 females), 11 $BK\alpha^{+/-}$ (7/4), and 10 $BK\alpha^{-/-}$ mice (7/3) aged 7 to 9 month (33 ± 2 weeks) were tested on the CatWalk (Noldus Information Technology Inc, Leesburg, VA, USA). Animals traversed a transparent walkway through which they were video-taped. Animals were allowed to walk back and forth until a minimum of 5 consecutive steps could be recorded. Using the CatWalkTM software (Version 7.0, Noldus Information Technology Inc, Leesburg, VA, USA) the paw prints of the animals were tracked. We analyzed stride length and paw print area of each individual paw, as well as inter-limb coordination using the regularity index (RI) as introduced by Hamers et al. [36]. The regularity index grades the inter-limb coordination as the ratio between the number of normal step sequence patterns and the total number of paw placements. A RI of 100% indicates perfect inter-limb coordination.

Open field locomotor activity

General locomotor activity was measured in 16 mice of each genotype (WT: 9 males/ 7 females, $BK\alpha^{+/-}$: 10/6, $BK\alpha^{-/-}$: 9/7) at the age of 6 to 9 month (30 ± 2.5 weeks). Each animal was placed in a square open field box (Versamax Animal Activity Monitor, AccuScan Instruments, Columbus, OH, USA) for 2 hours. During the whole time of testing the room was moderately lit (200 lux). Using the VersMaxTM software (AccuScan Instruments) we analyzed the distance traveled during the 2 hours, the number of rearing movements, as well as the time spent in the center versus the time spent at the margins as a measure of anxiety.

Acoustic startle reflex/prepulse inhibition (PPI)

Prepulse inhibition of the acoustic startle reflex was used to measure sensorimotor gating. 18 WT (10 males/8 females), 17 $BK\alpha^{+/-}$ (10/7), and 19 $BK\alpha^{-/-}$ mice (9/10) were tested as described previously [37,38]. Only animals aged 4-6 months (21.5 ± 2 weeks) were tested to avoid possible effects of age induced hearing loss. Testing was conducted in sound attenuated startle boxes from MED Associates (MED-ASR-PRO1, St Albans, VT, USA) using the associated software for stimulus presentation and recordings (Startle Reflex, Version 6.0, MED Associates, Inc.).

Sound was presented by two loud speakers in each startle box. The first speaker presented a constant background noise (65 dB SPL white noise), whereas the second speaker presented the startle and prepulse stimuli. The sound was calibrated with the sound pressure level measurement package provided by MED Associates

(ANL-929A-PC) placing the microphone into the animal holder at approximately the same distance to the speaker as the animals' head during the test sessions. To record the startle response, the animals were placed into a tube like plastic animal holder which was then mounted on top of a piezo driven transducer which translated vertical movements into a voltage signal. The voltage signal in response to a given stimulus was amplified, digitalized and stored on a computer. We started the recording 50 ms before any stimulus was given and recorded for a total of 500 ms per stimulus.

Before the actual testing the animals were acclimatized to the startle boxes for 3 days to reduce anxiety. During these 5 minute long acclimation phases the background noise was presented. On the third day, these 5 minutes were followed by a short input/output (I/O) test to determine the appropriate gain setting for the transducer signal for each individual animal. The I/O test consisted of 12 acoustic stimuli with increasing intensity (65–120 dB SPL, 20 ms duration). Once set, the gain was kept constant for all recordings of a given animal. In order to reduce variability we tested each animal on 5 consecutive days using the following protocol.

Animals were first acclimatized to the startle boxes as during the acclimation days, and then habituated to the startle stimulus to avoid habituation effects during the prepulse testing. For habituation, the startle stimulus (20 ms, 105 dB SPL white noise) was presented 100 times with varying inter-trial intervals (10–20 seconds). The prepulse subsequent test consisted of 7 different trial conditions which were each presented 10 times in a pseudo-randomized order and at varying inter-trial intervals (10–20 seconds). The trial conditions included startle alone trials to determine the baseline startle, and trials with one of two types of acoustic prepulses (4 ms, 75 and 85 dB SPL, white noise) leading the startle pulse by 10, 30, or 100 ms. Unfortunately, animals sometimes startled to the prepulse of 85 dB SPL, thus we omitted these trials from further analysis, since effects of muscle fatigue could not be excluded.

For each trial the startle amplitude was calculated as the difference between the maximum positive and negative displacement of the transducer induced by the startling animal. Prepulse inhibition was calculated for each animal and each trial condition as $PPI (\%) = (1 - \text{average startle amplitude to pulse with prepulse} / \text{average startle amplitude to pulse only}) * 100$.

Y-maze

In order to assess working memory and exploration behavior 15 mice of each genotype (WT: 9 males/ 6 females, $BK\alpha^{+/-}$: 10/5, $BK\alpha^{-/-}$: 7/8) aged between 8 and 10 month (36.5 ± 2 weeks) were tested in the Y-maze. The custom-made maze consisted of three white plastic arms separated by a 120° angle. Each arm was 30 cm long, 10 cm wide, and restricted by 20 cm high walls. The room in which the test was performed was moderately lit (200 lux). The animals were placed in the center of the maze and were allowed to explore freely for 5 minutes, during which the number and order of arm entries was recorded. An arm entry was defined as the animal placing all four paws into a given arm. Spontaneous alternation was calculated as the ratio of number of triads (sequence of 3 consecutive arm entries visiting all 3 arms) and total arm entries.

The strength of the spontaneous alteration task is that it requires no prior training, nor food or water restriction. The animals naturally tend to explore new environments and therefore prefer to investigate a new arm of the maze rather than returning to the previously explored one. In order to achieve this, the animals need to recall which arm they visited last (for review see: [39–41]).

Water maze

To assess spatial reference learning and memory we tested 14 mice of each genotype (WT: 9 males/ 4 females, $BK\alpha^{+/-}$: 10/4, $BK\alpha^{-/-}$: 7/7) in the Morris water maze. The animals were between 10 and 12 month (46 ± 3 weeks) old at the time of testing. We used a circular pool with a diameter of 1.5 meters and 60 cm high walls. The water was 45 cm deep and at room temperature (24°C). The pool was divided into equally sized quadrants in each of which a squared platform (10×10 cm) could be positioned in the middle. The platform was transparent and about 1.5 cm below the water surface to avoid that the animals can see it. Permanent visual cues on the room walls allowed for spatial orientation. A camera mounted above the pool recorded every trial and animals were tracked using the ANY-maze™ software (Version 4.98, Stoelting Co, Wood Dale, IL, USA)

Each animal underwent four training days with four trials a day (10–15 minutes inter-trial interval). The platform position was assigned randomly, but was counterbalanced for the genotypes. Once assigned, the platform position for a given animal was not changed between trials and days. However, the starting position of the animal was changed between every trial (first trial randomized, for the following trials the position moved clockwise between the quadrants). If the mouse did not find the platform within 90 seconds it was guided to the platform on the first trial. The animals were allowed to sit on the platform for 10 to 15 second and were then moved back into their home cages. For every trial we analyzed the time it took the animal to reach the platform. On the fifth day we ran a 60 second probe trial without the platform and recorded the time the animals spent in the quadrant where the platform previously was hidden.

Statistics

Data analyses and graphical display were done with Microsoft Excel (Version 14.0.6129.5000, Microsoft Corp.), Graphpad (for graphical display, Prism 6.01, Graphpad Software, LaJolla, CA, USA) and SPSS (for statistical analysis, Version 20.0.0, IBM Corp.). Data are expressed as mean \pm S.E. We also displayed the standard error in the graphs as measure of precision of the sample mean.

Depending on the paradigm, a 2-Way-ANOVA, a MANOVA, or a repeated measurement ANOVA with genotype and gender as between-subjects factors was performed to compare groups.

For repeated measurement ANOVAs the Mauchly test was used to judge if the data violated the sphericity assumption. In case of a violation the degrees of freedom were corrected using the Greenhouse-Geisser (if $\epsilon < .75$) or the Huynh-Feldt method (if $\epsilon > .75$). For post hoc analyses the Sidak t-test was performed. Differences were considered statistically significant when p-values were smaller than 0.05. In the figures significance levels between genotypes were indicated as followed: * $p < .05$, ** $p < .01$, *** $p < .001$.

Results

We tested the F1 generation of homozygous $BK\alpha$ knockout mice ($BK\alpha^{-/-}$), as well as their heterozygous ($BK\alpha^{+/-}$), and wildtype (WT) littermates for their hearing ability and motor function. Subsequently, we assessed their cognitive functions using a variety of tests.

All mice had a normal development of weight, from 15 g body weight at the age of 4 weeks up to 30 g at 15 weeks. There was no significant difference in the absolute weight and weight gain between the WT and the $BK\alpha^{-/-}$ mice. We also did not find a gender effect for the given age range tested (see Fig. S1).




Figure 1. Hearing thresholds measured with auditory brainstem response. (A) Click thresholds of WT and BK α ^{-/-} mice. Mean and S.D. (filled symbols). Single animal thresholds are shown as smaller, open symbols beside either group mean. The difference in threshold (3 dB) was statistically not significant. (B) Pure tone thresholds (mean and S.D.) as function of stimulus frequency for WT (black circles) and BK α ^{-/-} mice (grey squares). Threshold curves were just significantly different, based on significant elevated thresholds at 32 kHz (*:p' = 0.048, 1-sided t-test without alpha correction) and a drop out of threshold for BK α ^{-/-} mice at 45 kHz (star: p = 0.0262, Fisher Exact Probability Test). doi:10.1371/journal.pone.0081270.g001

Hearing

Hearing function was measured by ABR thresholds. Thresholds for click evoked ABRs were not statistically different for WT (19.8 ± 5.8 dB SPL) and BK α ^{-/-} mice (23.3 ± 6.2 dB SPL, $p = 0.13$, Fig.1A); and within the tested age range (4 to 15 weeks) the ABR thresholds did not change significantly (see Fig. S2). Consistent with the thresholds for the (broadband) click stimulus, the thresholds for frequency specific pure-tone stimuli were similar for WT and BK α ^{-/-} mice over the low and middle frequency range of hearing (4–22 kHz, Fig.1B). At frequencies of 32 kHz and above the hearing thresholds of the BK α ^{-/-} mice were significantly elevated and not even detectable at 45 kHz (2-way ANOVA: $p = 0.044$, 45 kHz: $p = 0.0262$). Despite this high frequency hearing loss, a progression of hearing thresholds upon aging could not be found within the frequency range of best hearing (8–16 kHz, also addressed with the click stimulus, see Fig.S2).

Thus, we concluded that cognitive tests that require normal hearing function can be performed with this animal model, as long as testing occurs in the low and/or mid frequency range.

Motor Functions

BK channels are abundantly expressed in the cerebellum and other motor areas in the brain. Therefore, deficient BK channel function might affect motor function. Since almost all cognitive tests rely on motor behavior and/or locomotion, we next tested these in our mice.

All BK α ^{-/-} mice had a noticeable muscle tremor, whereas their BK α ^{+/-} and WT littermates moved normally. Apart from the just visual evidence, we were able to detect and measure this tremor during the startle testing. A two-way ANOVA revealed that the voltage signal on the movement sensitive platform during the periods before the acoustic stimulation was significantly different for the genotypes (Fig.2A, $F(2,50) = 393.24$, $p < .001$, no gender effect, nor genotype x gender interaction). The BK α ^{-/-} mice had a significant higher noise (43.0 ± 1.2 mV) compared to the WT (34.3 ± 1.1 mV, $p < .001$) and BK α ^{+/-} mice (37.07 ± 1.17 mV, $p = .003$). This increased noise was not due to initiated movements of the mice during this brief period since that would have resulted in much larger signal amplitudes. All the recordings were visually inspected to exclude trials with




Figure 2. Muscle functions. (A) Muscular tremor was measured employing data recorded during the startle testing. During periods of no acoustic stimulation the noise of the recorded signal was significantly higher in the BK α ^{-/-} mice. (B) In addition to the tremor the BK α ^{-/-} mice also showed significant reduced muscle strength judging by their front limb grip force. doi:10.1371/journal.pone.0081270.g002




Figure 3. Gait strait analysis. (A) Examples of the paw print and their timing for WT (left) and BK $\alpha^{-/-}$ mice (right). The bars indicating the timing of each single paw on the floor (from left to right: left hind limb, left front limb, right front limb, right hind limb). (B) Stride length for all paws is reduced in the BK $\alpha^{-/-}$ mice, but (C) no significant differences were found for the paw print areas. (D) The inter-limb coordination as measured with the regularity index in reduced in BK $\alpha^{-/-}$ mice. The dashed line shows a regularity index of 100% indicating perfect inter-limb coordination. doi:10.1371/journal.pone.0081270.g003

spontaneous movement. Thus, the increased noise seems to reflect the muscular tremor seen in the BK $\alpha^{-/-}$ mice.

In order to assess the muscular strength of the mice we used the grip force test (Fig.2B). A two-way ANOVA reported a significant difference between the genotypes ($F(2,42) = 119.03, p < .001$). BK $\alpha^{+/-}$ mice (0.38 ± 0.01 lbf) showed similar strength as WT mice (0.41 ± 0.01 lbf, $p = .361$), whereas BK $\alpha^{-/-}$ mice showed significant lower muscular strength compared to both, WT ($p < .001$) and BK $\alpha^{+/-}$ mice ($p < .001$). With a grip force of on average 0.17 ± 0.01 lbf they had less than half the muscle strength of their

WT littermates. There was no gender effect, nor a genotype x gender interaction for the grip force.

In order to evaluate the ability of the mice to walk properly, they were tested on the cat walk. We analyzed stride length and paw print area of each individual paw, as well as inter-limb coordination. An example of paw prints and the respective timing of the paws is given in Fig.3A for both WT and BK $\alpha^{-/-}$ mice. A MANOVA on all parameters captured from the catwalk, reported a highly significant difference between genotypes ($\lambda_{Wilks} = .082, F(18,36) = 5.00, p < .001$, no gender effect and no genotype x gender interaction).




Figure 4. Locomotion. (A) BK $\alpha^{-/-}$ mice show normal locomotion in the open field box as well as (B) normal rearing activity. doi:10.1371/journal.pone.0081270.g004

Most prominent was the difference between the genotypes in the stride length (Fig.3B). The effect was consistent throughout all four limbs, but we will describe this effect here using the left front limb as an example. For the left limb stride length a follow-up two-way ANOVA reported significant differences for the genotype ($F(2,26) = 32.6, p < .001$). The stride length of the BK $\alpha^{-/-}$ mice was about 30% shorter than that of WT and BK $\alpha^{+/-}$ mice, with an average stride length of 5407 ± 150 mm as opposed to 7815 ± 220 mm ($p < .001$) and 8219 ± 330 mm ($p < 0.001$), respectively. In contrast, the paw print area did not differ between genotypes for any of the paws (Fig.3C).

In order to grade the inter-limb coordination, the regularity index (RI) was calculated. Whereas 81% (9/11) of the WT animals were rated as having a perfect RI, only 45% (5/11) of the BK $\alpha^{+/-}$ mice and only 27% (3/11) of BK $\alpha^{-/-}$ mice were rated perfect. Indeed, the average regularity index for the genotypes differed significantly ($F(2,26) = 4.35, p = .023$, Fig.3D). With a $RI_{BK\alpha^{-/-}} = 91.5 \pm 2.2\%$ the BK $\alpha^{-/-}$ mice showed significantly less regularity in their walking pattern than the WT mice which had an average $RI_{WT} = 99.3 \pm 0.5\%$ ($p = .037$). With a $RI = 95.84 \pm 2.2\%$ the BK $\alpha^{+/-}$ mice showed intermediate inter-limb coordination, but it was not significantly different to either the WT ($p = .416$), or the BK $\alpha^{-/-}$ mice ($p = .507$).

Despite the BK $\alpha^{-/-}$ mice showing tremor, reduced muscle strength, and gait disturbances, they exhibited normal locomotor

behavior (Fig.4). A two hour test in the open field box yielded no significant differences between the genotypes (MANOVA, $\lambda_{Wilks} = .928, F(6,80) = 0.64, p = .801$, no gender effect, nor genotype x gender interaction). On average the BK $\alpha^{-/-}$ mice traveled the same distance (989 ± 127 cm) with the same speed (9.8 ± 0.7 cm/s) and showed similar rearing activity (30.1 ± 8.8) as their WT (distance = 840 ± 76 cm, speed = 9.0 ± 0.2 cm/s, rearings = 24.1 ± 5.6), and BK $\alpha^{+/-}$ litter mates (distance = 1050 ± 200 cm, speed = 9.4 ± 0.3 cm/s, rearings = 42.9 ± 11.3). There was also no significant difference between genotypes for the respective parameters within the first 5 minutes of the open field test. To further assess locomotion, we also measured the swim speed of the mice in the Morris water maze. We could not detect any differences in swim speed between genotypes ($F(2,36) = 1.20, p = .314$, no gender effect, nor genotype x gender interaction, data not shown).

We conclude that BK $\alpha^{-/-}$ mice show normal locomotion but disrupted fine motor coordination. Therefore, cognitive tests that rely on normal locomotion can be performed with this animal model, as long as fine motor coordination is not required.

Sensorimotor gating

We originally tested prepulse inhibition over five days to reduce variability in the data. Interestingly, the repeated measurements ANOVA (inter-stimulus interval x day) did not only report a




Figure 5. Prepulse inhibition (PPI) of the acoustic startle response. (A) On the first day of testing no significant differences in prepulse inhibition were found between genotypes for all tested inter-stimulus intervals. (B) Across days the WT and BK $\alpha^{+/-}$ mice showed improvement in PPI whereas the BK $\alpha^{-/-}$ mice did not. The average PPI for all tested ISI is shown, since no effect of ISI was reported by the repeated measurements ANOVA. The right panel shows the average difference in PPI between the fifth and first day of testing. Differences were calculated for each single animal and then averaged for genotypes. doi:10.1371/journal.pone.0081270.g005




Figure 6. Spontaneous alternation in the Y-maze. (A) Spontaneous alternations are presented as percent of total arm entries. There was no difference between the genotypes. (B) The number of total arm entries is significantly increased in the BK $\alpha^{-/-}$ mice. doi:10.1371/journal.pone.0081270.g006

genotype effect ($F(2,48) = 6.04, p = .005$, no gender effect, no genotype x gender interaction), but also a significant effect for the day of testing ($F(3,922,188.275) = 7.198, p < .001$) and a day x genotype interaction ($F(7,845,188.275) = 2.155, p = .034$). Therefore, we analyzed the first day of testing separately and then additionally analyzed the development of prepulse inhibition across the 5 days of testing.

For the first day of testing the repeated measurements ANOVA (inter-stimulus interval) on prepulse inhibition did not reveal any significant differences between the genotypes ($F(2,50) = 0.338, p = .715$), nor an effect of gender, or a genotype x gender interaction (Fig.5A), suggesting that the deficit in BK channel function does not impair prepulse inhibition. However, we did find that the prepulse inhibition in WT and BK $\alpha^{+/-}$ improved over the days of testing, whereas in BK $\alpha^{-/-}$ mice it stayed constant (Fig.5B). Since the best prepulse inhibition was typically seen on the fifth day of testing, we computed the difference between the performance on day five and one for each animal to judge their improvement of PPI over days. The ANOVA confirmed that in the BK $\alpha^{-/-}$ mice prepulse inhibition improved significantly less than in their WT littermates ($F(2,49) = 5.05, p = .010$, no gender effect, no genotype x gender interaction). While in the WT mice prepulse inhibition on day five was $34.0 \pm 6.1\%$ higher, it was only improved by $5.3 \pm 6.7\%$ in the BK $\alpha^{-/-}$ mice.

Spontaneous Alternation

We next tested our animals for spatial working memory using the spontaneous alternation task in the Y-maze. A two-way ANOVA reported no significant differences between the genotypes ($F(2,39) = 0.04, p = .960$, Fig.6A). All genotypes showed on average 52% alternation, suggesting that there is no impairment in spatial working memory in the BK $\alpha^{-/-}$ mice. However, note that there was a significant difference in the number of total arm entries between the genotypes ($F(2,39) = 17.13, p < .001$, Fig.6B). The BK $\alpha^{-/-}$ mice showed twice as many arm entries (22.1 ± 1.8) as the WT ($11.3 \pm 0.7, p < .001$) and BK $\alpha^{+/-}$ mice ($13.3 \pm 1.5, p < .001$), and therefore spent less time in a arm per entry. It could be observed that they did not explore the arms completely. This possibly indicates a different exploratory behaviour strategy. There was no gender effect, nor a genotype x gender interaction for spontaneous alterations or number of arm entries.

Spatial learning and memory

Finally, we tested the animals for spatial reference learning and memory in the Morris water maze. The animals underwent four

training days with four trials a day. As mentioned previously, there was no difference in the swim speed for the genotypes. However, repeated measurements ANOVA (day x trials) for the time to reach the platform reported significant differences between genotypes ($F(2,36) = 11.33, p < .001$, no gender effect, nor genotype x gender interaction). The ANOVA also found a main effect for the day ($F(3,108) = 57.65, p < .001$) and for the trial ($F(3,108) = 5.306, p = .002$), but there was no interaction with genotype for neither of them, nor a day x trial x genotype interaction. The BK $\alpha^{-/-}$ mice took longer to reach the platform throughout all trials (Fig.7A, B). On average, it took the BK $\alpha^{-/-}$ mice 38.8 ± 2.8 seconds to get to the platform, while the WT and BK $\alpha^{+/-}$ mice reached the platform in 22 seconds (WT = 22.1 ± 2.9 s, $p < .001$, BK $\alpha^{+/-}$ = 22.6 ± 3.1 s, $p < .001$).

When analyzing the performance within days, the BK $\alpha^{-/-}$ mice show less improvement over the trials, especially on days one to three. However, they did improve over the days; it just took them longer to learn the task than their WT and BK $\alpha^{+/-}$ littermates. Indeed, the exponential fits to the data yielded a longer time constant for the BK $\alpha^{-/-}$ mice ($\tau = 2.72$ days) compared to the WT ($\tau = 0.80$ days) and BK $\alpha^{+/-}$ mice data ($\tau = 0.70$ days, Fig.7B). Eventually they do seem to catch up on day four where they did perform as well as WT and BK $\alpha^{+/-}$ mice in the last trial.

Once the task was learned, the BK $\alpha^{-/-}$ mice did remember the localization of the platform as well as their WT and BK $\alpha^{+/-}$ littermates. On the probe day all mice spent a significantly higher amount of time in the quadrant where previously the platform was hidden compared to the other quadrants (Fig.7C). A MANOVA on the time in the target quadrant, as well as the time in the neighboring and opposite quadrants did not find a genotype effect ($\lambda_{\text{Wilks}} = .698, F(6,66) = 2.17, p = .057$), nor a gender effect, or a genotype x gender interaction. The WT and BK $\alpha^{-/-}$ mice spent an equal amount of time in the quadrant where previously the platform was hidden ($51.6 \pm 4.2\%$ and $56.2 \pm 3.9\%$ respectively). Thus, we conclude that the BK $\alpha^{-/-}$ mice do not have spatial memory deficits.

In summary, the present study suggests that a knockout of the α -subunit of the BK channel results in deficits in learning. The spatial working memory and the spatial reference memory seem not to be affected.

Discussion

In the present study we used for the first time a genetic model of BK channel deficiency to test the role of BK channels in cognition. We used mice with a hybrid SV129/C57BL6 background which




Figure 7. Spatial learning and memory in the Morris water maze. (A) The average escape latency improved across the training trails in all genotypes, but especially on days one to three, the BK $\alpha^{-/-}$ mice improve less within the training sessions. Note that on the last training trail on day 4 they caught up and performed as well as the WT mice. (B) Escape latency as a function of days of experience with the task. The average escape latency for each day was fitted with a one component exponential function for each genotype and time constants were derived from the resulting functions. The BK $\alpha^{-/-}$ mice have a longer time constant than the WT and the BK $\alpha^{+/-}$ mice. (C) Performance on the probe day. All mice spend a significant higher amount of time in the quadrant where previously the platform was hidden (target, solid dark grey) than in any other quadrant. The dashed line indicates 25% chance criterion.
doi:10.1371/journal.pone.0081270.g007

are not hearing impaired and show normal locomotion. We found that the ablation of functional BK channels disrupts prepulse inhibition learning and slows down the acquisition of new tasks, but does not affect memory.

Auditory processing and motor function in BK channel knockout mice

We first verified that our hybrid SV129/C57BL6 BK $\alpha^{-/-}$ mice could perform all tasks required for the cognitive tests that we employed. One of the major concerns was their hearing. Rüttiger et al. [20] showed that BK channels are critical for the survival of high frequency responding outer hair cells in the basal cochlea and that a deletion of the pore forming α -subunit of the BK channel leads to progressive hearing loss with age starting around eight weeks at which the high frequency range seems to be especially affected. In contrast, Pyott et al. [24] did not find any age-dependent changes in auditory thresholds. Whereas Rüttiger et al. [20] tested a 129SVJ inbred mouse line, Pyott et al. [24] worked with a FVB/NJ inbred mice. Thus, the hearing phenotype of BK α subunit deficient mice seems to be strain specific. Further, Pyott et al. [24] suggested that the BK channels are only required under extreme levels of hair cell activity where they serve to protect normal hearing.

In the present study we used a hybrid SV129/C57BL6 mouse line and we could not find any age depending reduction of hearing thresholds for the click stimuli, which were later used in the acoustic startle reflex experiments. However, we still see an effect of the BK channel function loss on the high frequency range hearing thresholds. This high frequency hearing loss should not affect the acoustic startle reflex experiments since the click stimuli used there only include low to mid range frequencies. BK channels can not only alter hearing thresholds, but also temporal coding of sounds [21,22] which could prevent the detection of the short (4 ms) prepulse in the acoustic startle experiments. However, Kurt et al. [22] showed that mice with a conditional BK channel knockout for the inner hair cells surprisingly have improved discrimination of temporally modulated sounds. Thus, the BK channel knockout is unlikely to have affected the detection of the prepulse in our experiments.

Another major concern with general BK channel knockout mice is their motor function. BK channels are abundantly expressed in the cerebellum [1,3,23] and loss of BK channel function has been shown to lead to cerebellar ataxia with intentional tremor, abnormal gait, and motor coordination [23]. We mainly confirmed most of these findings in the mouse model used here. However, despite tremor, impaired muscle strength, and gait disturbances the mice exhibited normal locomotor behavior, for both walking and swimming, making it suitable for cognitive tests involving locomotion, but not fine motor coordination.

BK channels in sensorimotor gating

BK channels deficiency has been implicated in mental retardation/ fragile X syndrome [16,17], autism [18], and schizophrenia (for review see [19]). Impaired prepulse inhibition is one common symptom of these mental disorders (mental retardation: [28], autism: [29,30], schizophrenia: [31–33]). Thus, we expected to find deficits in prepulse inhibition in the BK channel knockout mice. However, prepulse inhibition in our BK channel knockout mice was not significantly different to that of their wild-type littermates on the first day of testing. Prepulse inhibition was quite low for both the wild-type and the knockout mice with only about 20-30%, which has been reported before for different mice strains (compare [42,43]).

Interestingly, we found a deficit in prepulse inhibition learning. Wild-type mice showed an increase in prepulse inhibition across the five days of testing to an average of around 60% whereas the knockout mice did not substantially improve. This effect cannot be attributed to differences in the baseline startle, since both wild-type and knockout mice had similar baseline startle amplitudes. A floor effect in the knockout mice can be further excluded because the absolute startle amplitude in the wild-type mice was smaller than in the knockout mice in the presence of a prepulse, and all startle amplitudes were at least twice the noise level and well detectable (data not shown). Experience-dependent increase in prepulse inhibition has been described before and has been attributed to an associative learning process which links the prepulse with the startle stimulus in a timely manner [42,43]. However, it is unclear where and how exactly this effect is mediated. Our data suggests a critical role for the BK channels in mechanisms mediating prepulse inhibition learning.

BK channels in learning and memory

Another common symptom of the mental disorders linked with BK channel deficiency is cognitive disruption. BK channels are expressed almost throughout the entire central nervous system, but the expression is particularly high in the hippocampus [2,3,34]. Thus, it is likely that hippocampus dependent tasks are impaired if BK channel function is disrupted. Indeed, it took BK channel knockout mice longer to learn the Morris water maze task. Matthews and Disterhoft [44] reported a comparable effect on trace eyeblink conditioning when blocking BK channels in the hippocampus with intracranial injections of paxilline. The animals that had received drug injections learned the association of the auditory stimulus with the air puff to the eye considerably slower than the control animals. Sausbier et al. [23] also reported slower acquisition of the conditioned eyeblink reflex in global BK channel knockout mice (129SVJ inbred), but they attributed it to a cerebellar learning deficit. In line with these findings is also the deficit in prepulse inhibition learning in the BK channel knockout mice which is attributed to an associated learning process as discussed above. Taken together, these studies indicate a crucial role for the BK channels in synaptic mechanisms underlying learning. This might be an evolutionary highly conserved mechanism, as a calcium dependent potassium current also has been linked to associative learning in molluscs [45].

How exactly the BK channels are involved in learning is not fully understood yet. BK channels mediate the fast after-hyperpolarization and thus regulate intrinsic neuronal excitability [7,10,46,47]. Blocking the BK channels with either paxilline or iberiotoxin increases the firing rate to step currents in *in vitro* whole-cell recordings [10], as well as the *in vivo* spontaneous firing

rate of hippocampal CA1 [48]. Additionally, BK channels have been shown to regulate transmitter release at the presynaptic terminals [8,12,13]. Blocking the BK channels enhances the probability of transmitter release at CA3-CA3 synapses in the hippocampus [12]. Together those effects lead to an increased neuronal noise which reduces the dynamic range for neurons to code information on a firing rate base in BK channel deficient systems. However, BK channels are expressed abundantly in the central nervous system, it is therefore possible that other structures beside the hippocampus are involved in mediating the described effects. In particular, the substantia nigra, the habenula, and striatum also have been shown to be involved in associative learning and they express BK channels [49,50].

Interestingly, memory, once it is established seems not to be affected by the BK channel knock-out. Both, working memory (which was measured with the y-maze spontaneous alternation task), as well as the spatial reference memory (which was measured on the probe day in the Morris water maze), were comparable between the knock-out and the wild-type mice.

Conclusion

The present study shows that BK channel knock-out mice do not improve prepulse inhibition of startle over days and are slower in learning the Morris water maze task, suggesting a critical role for the BK channels in mechanisms underlying associative learning. BK channels may therefore become an increasingly interesting target for the development of new drugs for enhancing cognitive function.

Supporting Information

Figure S1 Weight gain in males (blue) versus females (red) and wild-types (dots) versus knock-out (squares) mice. No difference in body weight gain was observed between any of these groups during 4 to 15 weeks of age. (TIF)

Figure S2 ABR hearing thresholds for WT (black circles) and BK α -/- mice (grey squares) for click stimuli did not change significantly with age. Animals measured on more than one time point are connected with lines. (TIF)

Author Contributions

Conceived and designed the experiments: MT EA LR PR SS. Performed the experiments: MT EA MM LR. Analyzed the data: MT MM EA SS LR. Contributed reagents/materials/analysis tools: LR SS PR. Wrote the paper: MT MM SS. Created mouse line: PR.

References

- Knaus HG, Schwarzer C, Koch RO, Eberhart A, Kaczorowski GJ, et al. (1996) Distribution of high-conductance Ca(2+)-activated K+ channels in rat brain: targeting to axons and nerve terminals. *J Neurosci* 16: 955–963.
- Wanner SG, Koch RO, Koschak A, Trieb M, Garcia ML, et al. (1999) High-Conductance Calcium-Activated Potassium Channels in Rat Brain: Pharmacology, Distribution, and Subunit Composition. *Biochemistry (Mosc)* 38: 5392–5400. doi:10.1021/bi983040c.
- Sausbier U, Sausbier M, Sailer C, Arntz C, Knaus H-G, et al. (2006) Ca2+-activated K+ channels of the BK-type in the mouse brain. *Histochem Cell Biol* 125: 725–741. doi:10.1007/s00418-005-0124-7.
- Misonou H, Menegola M, Buchwalder L, Park EW, Meredith A, et al. (2006) Immunolocalization of the Ca2+-activated K+ channel Slo1 in axons and nerve terminals of mammalian brain and cultured neurons. *J Comp Neurol* 496: 289–302. doi:10.1002/cne.20931.
- Sailer CA, Kaufmann WA, Kogler M, Chen L, Sausbier U, et al. (2006) Immunolocalization of BK channels in hippocampal pyramidal neurons. *Eur J Neurosci* 24: 442–454. doi:10.1111/j.1460-9568.2006.04936.x.
- Vergara C, Latorre R, Marrion NV, Adelman JP (1998) Calcium-activated potassium channels. *Curr Opin Neurobiol* 8: 321–329. doi:10.1016/S0959-4388(98)80056-1.
- Poolos NP, Johnston D (1999) Calcium-Activated Potassium Conductances Contribute to Action Potential Repolarization at the Soma But Not the Dendrites of Hippocampal CA1 Pyramidal Neurons. *J Neurosci* 19: 5205–5212.
- Hu H, Shao L-R, Chavoshy S, Gu N, Trieb M, et al. (2001) Presynaptic Ca2+-Activated K+ Channels in Glutamatergic Hippocampal Terminals and Their Role in Spike Repolarization and Regulation of Transmitter Release. *J Neurosci* 21: 9585–9597.
- Shao L-R, Halvorsrud R, Borg-Graham L, Storm JF (1999) The Role of BK-Type Ca2+-Dependent K+ Channels in Spike Broadening During Repetitive Firing in Rat Hippocampal Pyramidal Cells. *J Physiol* 521: 135–146. doi:10.1111/j.1469-7793.1999.00135.x.
- Nelson AB, Krispel CM, Sekirnjak C, du Lac S (2003) Long-Lasting Increases in Intrinsic Excitability Triggered by Inhibition. *Neuron* 40: 609–620. doi:10.1016/S0896-6273(03)00641-X.

11. Brenner R, Chen QH, Vilaythong A, Toney GM, Noebels JL, et al. (2005) BK channel $\beta 4$ subunit reduces dentate gyrus excitability and protects against temporal lobe seizures. *Nat Neurosci* 8: 1752–1759. doi:10.1038/nn1573.
12. Raffaelli G, Saviane C, Mohajerani MH, Pedarzani P, Cherubini E, et al. (2004) BK potassium channels control transmitter release at CA3–CA3 synapses in the rat hippocampus. *J Physiol J Physiol* 557, 557: 147, 147–157, 157. doi:10.1113/jphysiol.2004.062661, 10.1113/jphysiol.2004.062661.
13. Wang Z-W (2008) Regulation of Synaptic Transmission by Presynaptic CaMKII and BK Channels. *Mol Neurobiol* 38: 153–166. doi:10.1007/s12035-008-8039-7.
14. Robitaille R, Garcia ML, Kaczorowski GJ, Chariton MP (1993) Functional colocalization of calcium and calcium-gated potassium channels in control of transmitter release. *Neuron* 11: 645–655. doi:10.1016/0896-6273(93)90076-4.
15. Robitaille R, Charlton MP (1992) Presynaptic calcium signals and transmitter release are modulated by calcium-activated potassium channels. *J Neurosci* 12: 297–305.
16. Higgins JJ (2008) Dysregulation of large-conductance Ca²⁺-activated K channel expression in nonsyndromal mental retardation due to a cereblon p.R419X mutation. *Neurogenetics* 9: 219–223.
17. Deng P-Y, Rotman Z, Blundon JA, Cho Y, Cui J, et al. (2013) FMRP Regulates Neurotransmitter Release and Synaptic Information Transmission by Modulating Action Potential Duration via BK Channels. *Neuron* 77: 696–711. doi:10.1016/j.neuron.2012.12.018.
18. Laumonier PD, Roger PD, Guérin MD, Molinari PD, M'Rad MD, et al. (2006) Association of a Functional Deficit of the BKCa Channel, a Synaptic Regulator of Neuronal Excitability, With Autism and Mental Retardation. *Am J Psychiatry* 163: 1622–1629.
19. Zhang L, Li X, Zhou R, Xing G (2006) Possible role of potassium channel, big K in etiology of Schizophrenia. *Med Hypotheses* 67: 41–43. doi:10.1016/j.mehy.2005.09.055.
20. Rüttiger L, Sausbier M, Zimmermann U, Winter H, Braig C, et al. (2004) Deletion of the Ca²⁺-Activated Potassium (BK) A-Subunit but Not the BK β 1-Subunit Leads to Progressive Hearing Loss. *Proc Natl Acad Sci U S A* 101: 12922–12927. doi:10.1073/pnas.0402660101.
21. Oliver D, Taberner AM, Thurm H, Sausbier M, Arntz C, et al. (2006) The Role of BKCa Channels in Electrical Signal Encoding in the Mammalian Auditory Periphery. *J Neurosci* 26: 6181–6189. doi:10.1523/JNEUROSCI.1047-06.2006.
22. Kurt S, Sausbier M, Rüttiger L, Brandt N, Moeller CK, et al. (2012) Critical role for cochlear hair cell BK channels for coding the temporal structure and dynamic range of auditory information for central auditory processing. *FASEB J*. doi:10.1096/fj.11-200535.
23. Sausbier M, Hu H, Arntz C, Feil S, Kamm S, et al. (2004) Cerebellar ataxia and Purkinje cell dysfunction caused by Ca²⁺-activated K⁺ channel deficiency. *Proc Natl Acad Sci U S A* 101: 9474–9478. doi:10.1073/pnas.0401702101.
24. Pyott SJ, Meredith AL, Fodor AA, Vázquez AE, Yamoah EN, et al. (2007) Cochlear Function in Mice Lacking the BK Channel α , β 1, or β 4 Subunits. *J Biol Chem* 282: 3312–3324. doi:10.1074/jbc.M608726200.
25. Gerlai R (1996) Gene-targeting studies of mammalian behavior: is it the mutation or the background genotype? *Trends Neurosci* 19: 177–181. doi:10.1016/S0166-2236(96)20020-7.
26. Wolfer DP, Lipp H-P (2000) Dissecting the behaviour of transgenic mice: is it the mutation, the genetic background, or the environment? *Exp Physiol* 85: 627–634.
27. Wolfer DP, Crusio WE, Lipp H-P (2002) Knockout mice: simple solutions to the problems of genetic background and flanking genes. *Trends Neurosci* 25: 336–340. doi:10.1016/S0166-2236(02)02192-6.
28. Frankland PW, Wang Y, Rosner B, Shimizu T, Balleine BW, et al. (2004) Sensorimotor gating abnormalities in young males with fragile X syndrome and Fmr1-knockout mice. *Mol Psychiatry* 9: 417–425. doi:10.1038/sj.mp.4001432.
29. McAlonan GM, Daly E, Kumari V, Critchley HD, Amelvoort T van, et al. (2002) Brain anatomy and sensorimotor gating in Asperger's syndrome. *Brain* 125: 1594–1606. doi:10.1093/brain/awf150.
30. Perry W, Minassian A, Lopez B, Maron L, Lincoln A (2007) Sensorimotor Gating Deficits in Adults with Autism. *Biol Psychiatry* 61: 482–486. doi:10.1016/j.biopsych.2005.09.025.
31. Geyer MA, Braff DL (1982) Habituation of the blink reflex in normals and schizophrenic patients. *Psychophysiology* 19: 1–6.
32. Ludewig K, Geyer MA, Vollenweider FX (2003) Deficits in prepulse inhibition and habituation in never-medicated, first-episode schizophrenia. *Biol Psychiatry* 54: 121–128. doi:10.1016/S0006-3223(02)01925-X.
33. Takahashi H, Iwase M, Ishii R, Ohi K, Fukumoto M, et al. (2008) Impaired prepulse inhibition and habituation of acoustic startle response in Japanese patients with schizophrenia. *Neurosci Res* 62: 187–194. doi:10.1016/j.neures.2008.08.006.
34. Tseng-Crank J, Foster CD, Krause JD, Mertz R, Godinot N, et al. (1994) Cloning, expression, and distribution of functionally distinct Ca²⁺-activated K⁺ channel isoforms from human brain. *Neuron* 13: 1315–1330. doi:10.1016/0896-6273(94)90418-9.
35. Engel J, Braig C, Rüttiger L, Kuhn S, Zimmermann U, et al. (2006) Two classes of outer hair cells along the tonotopic axis of the cochlea. *Neuroscience* 143: 837–849. doi:10.1016/j.neuroscience.2006.08.060.
36. Hamers FPT, Lankhorst AJ, van Laar TJ, Veldhuis WB, Gispen WH (2001) Automated Quantitative Gait Analysis During Overground Locomotion in the Rat: Its Application to Spinal Cord Contusion and Transection Injuries. *J Neurotrauma* 18: 187–201. doi:10.1089/08977150150502613.
37. Geyer MA, Swerdlow NR (2001) Measurement of Startle Response, Prepulse Inhibition, and Habituation. *Current Protocols in Neuroscience*. John Wiley & Sons, Inc.
38. Valsamis B, Schmid S (2011) Habituation and prepulse inhibition of acoustic startle in rodents. *J Vis Exp JoVE*: e3446. doi:10.3791/3446.
39. Richman C (1986) Spontaneous alternation behavior in animals: A review. *Curr Psychol* 5: 358–391.
40. Lalonde R (2002) The neurobiological basis of spontaneous alternation. *Neurosci Biobehav Rev* 26: 91–104. doi:10.1016/S0149-7634(01)00041-0.
41. Hughes RN (2004) The value of spontaneous alternation behavior (SAB) as a test of retention in pharmacological investigations of memory. *Neurosci Biobehav Rev* 28: 497–505. doi:10.1016/j.neubiorev.2004.06.006.
42. Plappert CF, Pilz PK., Schnitzler H-U (2004) Factors governing prepulse inhibition and prepulse facilitation of the acoustic startle response in mice. *Behav Brain Res* 152: 403–412. doi:10.1016/j.bbr.2003.10.025.
43. Plappert CF, Kuhn S, Schnitzler HU, Pilz PKD (2006) Experience Increases the Prepulse Inhibition of the Acoustic Startle Response in Mice. *Behav Neurosci* 120: 16–23.
44. Matthews EA, Disterhoft JF (2009) Blocking the BK channel impedes acquisition of trace eyeblink conditioning. *Learn Mem* 16: 106–109. doi:10.1101/lm.1289809.
45. Alkon DL, Sakakibara M, Forman R, Harrigan J, Lederhendler I, et al. (1985) Reduction of two voltage-dependent K⁺ currents mediates retention of a learned association. *Behav Neural Biol* 44: 278–300. doi:10.1016/S0163-1047(85)90296-1.
46. Alger BE, Nicoll RA (1980) Epileptiform burst afterhyperpolarization: calcium-dependent potassium potential in hippocampal CA1 pyramidal cells. *Science* 210: 1122–1124. doi:10.1126/science.7444438.
47. Disterhoft JF, Oh MM (2006) Learning, aging and intrinsic neuronal plasticity. *Trends Neurosci* 29: 587–599.
48. Matthews EA, Weible AP, Shah S, Disterhoft JF (2008) The BK-mediated fAHP is modulated by learning a hippocampus-dependent task. *Proc Natl Acad Sci* 105: 15154–15159. doi:10.1073/pnas.0805855105.
49. Matsumoto M, Hikosaka O (2007) Lateral habenula as a source of negative reward signals in dopamine neurons. *Nature* 447: 1111–1115. doi:10.1038/nature05860.
50. Matsumoto M, Hikosaka O (2009) Two types of dopamine neuron distinctly convey positive and negative motivational signals. *Nature* 459: 837–841. doi:10.1038/nature08028.g