

Genetic Diversity of Food-Isolated *Salmonella* Strains through Pulsed Field Gel Electrophoresis (PFGE) and Enterobacterial Repetitive Intergenic Consensus (ERIC-PCR)

Imen Fendri^{1,2*}, Amal Ben Hassena¹, Noel Grosset², Mohamed Barkallah¹, Lamia Khannous¹, Victoria Chuat³, Michel Gautier², Radhouane Gdoura¹

1 Unité de recherche Toxicologie - Microbiologie Environnementale et Santé, Faculté des Sciences de Sfax, Université de Sfax, Sfax, Tunisia, **2** Laboratoire de Microbiologie, Département agroalimentaire Agrocampus Ouest, Rennes, France, **3** INRA, CIRMBIA, UMR 1253 Science et technologie du lait et de l'œuf (STLO), Agrocampus ouest, Rennes, France

Abstract

All over the world, the incidence of *Salmonella* spp contamination on different food sources like broilers, clams and cow milk has increased rapidly in recent years. The multifaceted properties of *Salmonella* serovars allow the microorganism to grow and multiply in various food matrices, even under adverse conditions. Therefore, methods are needed to detect and trace this pathogen along the entire food supply network. In the present work, PFGE and ERIC-PCR were used to subtype 45 *Salmonella* isolates belonging to different serovars and derived from different food origins. Among these isolates, *S. Enteritidis* and *S. Kentucky* were found to be the most predominant serovars. The Discrimination Index obtained by ERIC-PCR (0.85) was slightly below the acceptable confidence value. The best discriminatory ability was observed when PFGE typing method was used alone (DI=0.94) or combined with ERIC-PCR (DI=0.93). A wide variety of profiles was observed between the different serovars using PFGE or/and ERIC-PCR. This diversity is particularly important when the sample origins are varied and even within the same sampling origin.

Citation: Fendri I, Ben Hassena A, Grosset N, Barkallah M, Khannous L, et al. (2013) Genetic Diversity of Food-Isolated *Salmonella* Strains through Pulsed Field Gel Electrophoresis (PFGE) and Enterobacterial Repetitive Intergenic Consensus (ERIC-PCR). PLoS ONE 8(12): e81315. doi:10.1371/journal.pone.0081315

Editor: A. Mark Ibekwe, U. S. Salinity Lab, United States of America

Received: September 6, 2013; **Accepted:** October 21, 2013; **Published:** December 3, 2013

Copyright: © 2013 Fendri, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work received financial support from "Ministère de l'enseignement supérieur et de la recherche scientifique" granted to the "Unité de Recherche de Toxicologie, Microbiologie Environnementale et Santé". The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: imen.fendri@yahoo.fr

Introduction

Salmonellosis is a major health problem worldwide and accounts for high morbidity rates. Infection with *Salmonella enterica* occurs mainly through the consumption of contaminated food, and the estimated annual number of human infections is greater than 93.8 million cases, with 155,000 deaths per year worldwide [1]. Many *Salmonella* serovar infections result in diarrheal diseases, bacteraemia and extraintestinal focal infections in infants and more serious complications among the elderly and immunocompromised adults [2]. The pathogenicity [3] and the increase of antimicrobial resistance in *Salmonella* have been recognized as the ultimate causes. Filter feeding organisms such as clams harvested from contaminated waters are known to concentrate high levels of *Salmonella* serovars leading to a high incidence of this pathogen on seafood [4]. This is also the case in broilers [5]. However, information on milk contamination is scarce.

Different phenotypic and biochemical characteristics have been previously used for the epidemiological investigation of *Salmonella* [6,7]. Beyond the phenotypic characterization, a reliable genetic level discriminatory method is required. In fact, molecular typing methods that rely on DNA sequence differences are essential for

the epidemiological study of pathogenic *Salmonella* serovars [8]. Bacterial housekeeping genes were widely used for molecular typing and were based on polymorphisms analysis in defined genetic loci in the bacterial genome by the PCR amplification and sequencing of the PCR products [9].

Randomly Amplified Polymorphic DNA (RAPD) and Enterobacterial Repetitive Intergenic Consensus (ERIC) fingerprinting were recently used to differentiate *Salmonella* serovars in seafood and human origins [10–12]. An earlier study compared four molecular typing methods for the differentiation of *Salmonella* spp (RAPD, ERIC, Ribotyping PCR and Single Strand Conformation Polymorphism (SSCP)) and observed ERIC-PCR to be the most efficient [11]. ERIC is a short interspersed repetitive consensus sequence originally found in *E. coli* and *Salmonella* and ERIC-PCR uses outward facing primers complementary to each end of the repeat in a PCR [13]. On the other hand, the application of pulsed-field gel electrophoresis (PFGE) has been proved to be useful for the discrimination and epidemiological characterization of *Salmonella enterica* strains [14–16].

In the present study, the serotyped *Salmonella* isolated from Tunisian clams, broilers and milk were subjected to DNA based fingerprinting using PFGE and ERIC-PCR in order to

characterize *Salmonella* isolates collected from different origins and to define the relationships between them.

Materials and Methods

Salmonella isolation and serotyping

All the samples were analyzed according to the International Organization for Standardization Method 6579 (ISO). Isolation and biochemical identification were carried out according to standard laboratory methods. Suspected *Salmonella* colonies were screened using real time PCR. 1 ml of isolates culture was centrifuged at 13,000 g for 20 min. The pellet was resuspended in 200 µl sterile water. The total volume was extracted by Quick-gDNA MiniPrep D3006 Kit (Zymo Research, CA, USA) as recommended by the manufacturer. Extracted DNA was resuspended in 50 µl of elution buffer and stored at -20°C until subsequent analysis [17]. Real-time PCR was performed on the CFX96™ real-time PCR cycler (Biorad). Amplification reactions were carried out at a final volume of 25 µl containing 0.2 µM of each primer (Table 1), 12.5 µl of 2× SYBR® Permixon Ex Taq™ Tli RNaseH Plus (TaKaRa) and 1 µl of genomic DNA. PCR amplification was conducted by incubating the samples at 95 °C for 3 s, followed by 40 cycles of 5 s at 95 °C and 30 s at 60 °C. Primers amplifying *invA* gene were used as previously described [18]. A single confirmed *Salmonella* isolate from each positive sample was serotyped according to the Kauffman–White scheme using commercial antisera [19]. Serotyping was carried out at the National Centre of Enteropathogenic Bacteria, Pasteur Institute, Tunis.

Salmonella isolates selection

Forty five *S. enterica* isolates were selected for further molecular typing. These isolates were obtained from i) a total tissue of 7 clams from the sampling station M2 located at the Gulf of Gabes (Southern Mediterranean). Sampling was carried out manually by randomly picking up clams off the coast. The sampling process was supervised by the Commissariat Régional du Développement Agricole de Mednine (CRDA). ii) 28 samples derived from intestine, carcass and liver of sexed chickens (HubbardJV) collected at a meat processing industry situated in the region of Sfax and iii) 10 samples of cow milk collected from farms (Sfax, Tunisia) after the consent of their owners. Permission to use these




Figure 1. Representative PFGE fingerprint of differences between *Salmonella* isolates on 1 per cent agarose gel. M, lambda DNA marker; lane 1 to 14: O, P, Q, K, L, M, T, L1, L2, L3, L4, L5, L9 and L10.

doi:10.1371/journal.pone.0081315.g001

animal parts was obtained from the slaughterhouse (Sfax, Tunisia) to use these animal parts. The origin of each isolate and its appropriated serotype are given in Table 1.

Pulsed-field gel electrophoresis (PFGE)

Pulsed-field gel electrophoresis was performed according to the one-day (24 to 28 h) standardized laboratory protocol for the molecular subtyping of *Salmonella* by PFGE (Pulse-Net, Centers for Disease Control and Prevention, Atlanta, Ga.), (Centers for Disease Control and Prevention, 2005), and as described by Ribot et al. [20] with minor changes. Cells were suspended in a wash buffer (10 mM Tris-HCl pH 7.6, 1 M NaCl) and then lysed in a lysis buffer (10 mM Tris-HCl, 750 mM EDTA pH 9, 10% of N-Lauroylsarcosine (Sigma Aldrich, France) and 14 mg/ml proteinase K (Eurobio, France). Genomic DNA was prepared by embedding *Salmonella* isolates cells in agarose plugs (Invitrogen,

Table 1. Distribution of *Salmonella* isolates derived from broilers, clams and cow milk.

<i>Salmonella</i> serovars	Origin	Number of strains	Total	Frequency (%)
Enteritidis	Broiler Intestine	5	19	42.2
	Broiler Liver	7		
	Broiler Carcass	5		
	Clam	1		
Kentucky	Broiler Intestine	6	18	40.0
	Broiler Liver	4		
	Cow Milk	8		
Anatum	Cow Milk	2	2	4.4
London	Clam	2	2	4.4
Irenea	Clam	2	2	4.4
Poona	Clam	1	1	2.2
Brancaster	Clam	1	1	2.2

doi:10.1371/journal.pone.0081315.t001




Figure 2. PFGE Dendrogram showing the relationship between *Salmonella* isolates. The similarities between strains were evaluated using the Dice coefficient and the UPGMA clustering method. Genetic similarity between samples in duplicate is 80%. doi:10.1371/journal.pone.0081315.g002

France). XbaI (New England Biolabs, UK) was used to digest the DNA for each isolate. Electrophoresis was performed in a Biorad Chef-DR®-II system in 0.5X Tris-Borate-EDTA (TBE) extended-range buffer (Eurobio, France) with recirculation at 14°C. The following settings were used for DNA migration: Step I with an initial switch time of 20 sec, a final switch time of 45 sec, a gradient of 6 V/cm and 9 h of electrophoresis; Step II with initial switch time 5 sec, final switch time 15 sec, a gradient of 6 V/cm and 10 h of electrophoresis. Three lambda markers (lambda DNA cI857 and 1 Sam7, GelSyringe™, New England Biolabs, UK) were included on each gel. Following electrophoresis, the gel was stained with Gel-Red 3X in a 0.1 M NaCl solution (FluoProber, Interchim), visualized under UV light and then photographed.

ERIC-PCR for *Salmonella* isolates

For ERIC-PCR, the primers ERIC-1R (5'-ATGTAAGCTC-CTGGGGATTAC-3') and ERIC2 (5'AAGTAAGTGACTG-GGGTGAGCG-3') (Sigma Aldrich) [13,21] were used with some changes. The PCR was performed in a 50 µL solution containing 1 µM of each primer, 5 µL of 10X PCR buffer, 250 µM dNTPs, 3 mM MgCl₂, and 3.0 U of Taq DNA polymerase (Sigma Aldrich). The PCR conditions were one cycle at 95°C for 10 min, followed by 4 cycles of 5 min at 94°C, 5 min at 40°C and 5 min at 72°C and then followed by 30 cycles of 1 min at 94°C, 1 min at 55°C, and 2 min at 72°C, and the last extension at 72°C for 10 min. A 10 µl aliquot of each amplification reaction was analyzed using electrophoresis on a 2% agarose gel and run in a 1X TBE buffer, pH 8.3. The gel was stained with Gel-Red 3X in a 0.1 M NaCl solution (FluoProber, Interchim) and photographed.




Figure 3. Representative ERIC-PCR fingerprint of different between *Salmonella* isolates on 2 per cent agarose gel. M: 1000 bp DNA marker; lane 1 to 13: F34', F35', C31', C32', C33', C34', C36', L1, L2, L3, L4, L5 and L9.
doi:10.1371/journal.pone.0081315.g003

A 1000 base pair Smart Ladder (Eurogentec, France) was included on the gel as a marker.

Data Analysis

The banding patterns from PFGE and ERIC-PCR analysis were analyzed with BioNumerics Software version 6.5 (Applied-Maths, Ghent, Belgium). The similarities between strains were calculated using the Dice coefficient with an optimization of 1%. The dendrograms were obtained by means of the Unweighted Pair Group Method with Arithmetic Average (UPMGA) clustering algorithm. Numerical index of discriminatory ability of PFGE, ERIC-PCR and combined typing methods were calculated by applying Simpson's Index of Diversity equation as previously described by Nath et al. [12].

Ethical statement

None of the authors of this paper has a financial or personal relationship with other people or organizations that could inappropriately influence or bias the content of this work.

Results

Serotyping of *Salmonella* isolates

In this work, 45 *Salmonella enterica* isolates were recovered from different origins (Table 1). After serotyping, nine distinct serovars were identified, among which two were dominant: *Salmonella* Enteritidis (n=19) and *Salmonella* Kentucky (n=18). Minor serovars included *Salmonella* Anatum (n=2), *Salmonella* Irenea (n=2) and *Salmonella* London (n=2). Only one isolate was recovered for each of *S. Poona* and *S. Brancaster* (Table 1).

Verification of *Salmonella* isolates by real time PCR

All isolated strains were serotyped as *Salmonella* strains. To confirm this, all strains were analyzed by real time PCR using primers previously described as *invA* gene specific [22]. PCR results obtained in this study indicated that all *Salmonella* strains tested by PCR were positive for the presence of a 284 bp fragment of the *invA* gene.

PFGE typing results

Our PFGE observations raise the question of whether *Salmonella* isolates from different food origins are phylogenetically related or

comprise multiple lineages. Results show that most serovars, including *S. Enteritidis* and *S. Kentucky*, which are antigenically identical to each other, were assigned to multiple pulsotype profiles (Figure 1) and clusters (Figure 2). It is important to note that when three lambda markers were included on each gel, the genetic similarity between them was 80% (data not shown). Therefore we considered that two isolates presenting more than 80% of similarity were the same.

PFGE of XbaI-digested genomic DNA from 45 *Salmonella* isolates showed 13 different macrorestriction profiles or clusters (P1 to P13), while the remaining 5 isolates were unclustered. The latter belonged to serovars *S. Enteritidis* (I16 and O), *S. Kentucky* (I21), *S. Poona* (T) and *S. Brancaster* (K) (Figure 2). Six isolates were distributed among 3 clusters with two belonging to *S. Irenea* (P5), two isolates belonging to *S. Anatum* (P8) and two belonging to *S. London* (P10). Each cluster was composed of isolates derived from one origin, clam for P5 and P10 and milk for P8.

Heterogeneity was observed within the two major serovars, *S. Enteritidis* and *S. Kentucky*. The first serovar with 19 isolates was distributed among 2 unclustered isolates and 17 assigned to 6 PFGE clusters designated P6, P7, P9, P11, P12 and P13. About 95% of *Salmonella* isolates belonging to these clusters were derived from broiler samples (Figure 2) with 7 isolates from liver, 5 from intestine and 5 from total carcass.

The *S. Kentucky* serovar with 18 isolates was distributed among only one unclustered (I21) isolate and 17 assigned to 4 PFGE clusters designated P1, P2, P3 and P4 (Figure 2). Eight clustered isolates belonging to this serovar were derived from milk and nine from broiler samples (intestine or liver).

The discrimination index (DI) of PFGE in this analysis was found to be 0.94.

ERIC-PCR typing analysis

Two 1000 base pair Smart Ladders were included on the gel as a marker. The genetic similarity between them was 80% (data not shown). The ERIC-PCR of 45 *Salmonella* isolates yielded different patterns consisting of 3–9 bands (Figure 3). All the serovars were grouped into 8 clusters (E1 to E8) while the remaining 6 isolates were unclustered. The ungrouped isolates belonged to serovars *S. Enteritidis* (I32'), *S. Kentucky* (I20'), *S. Poona* (T), *S. Brancaster* (K) and *S. Anatum* (L9 and L10) (Figure 4). Clustering based on fragment profiles grouped *S. Enteritidis* serovar into only two clusters (E4 and E8) (Figure 4). The second major group, *S. Kentucky* serovar, was distributed among only one unclustered isolate and 17 assigned to 4 ERIC clusters designated E1, E2, E6 and E7 (Figure 4).

Finally, *S. Irenea* (E5) and *S. London* (E3) clusters contained two *Salmonella* isolates each. The discrimination index (DI) of ERIC-PCR typing in this analysis was found to be 0.85.

Composite analysis of PFGE and ERIC-PCR

Data from the two molecular typing methods were subjected to a composite analysis to determine whether a better clustering of the serovars could be obtained. Clustering based on fragment profiles grouped the serovars into 10 clusters (C1–C10) (Figure 5). The 19 isolates of *S. Enteritidis* were grouped into 4 clusters (C7, C8, C9 and C10), while the remaining 5 isolates were distinct from each other. Except I21, all isolates belonging to *S. Kentucky* serovar were differentiated into 3 groups (C4, C5 and C6). *S. Anatum* (C1), *S. London* (C2) and *S. Irenea* (C3) were still grouped into two isolates per cluster. The combined PFGE-ERIC-PCR patterns allowed a DI of 0.93.




Figure 4. Genetic similarities of *Salmonella* strains isolated from clams, broilers and milk based on ERIC-PCR patterns. The dendrogram was generated by BioNumerics Software with the bande-matching coefficient of Dice and the UPGMA clustering. Genetic similarity between samples in duplicate is 80%. doi:10.1371/journal.pone.0081315.g004

Discussion

Salmonellosis is one of the most common causes of foodborne infection worldwide. *Salmonella spp.* can be isolated from different origins such as raw meat and poultry products as well as milk and milk based products [23].

This work focuses on the assessment of two molecular methods (PFGE and ERIC-PCR) for inter and intraserovar strains differentiation. These techniques were evaluated both alone and in combination for typing *Salmonella* isolates. The methods applied in our study are among those methods used for epidemiologic analysis of important zoonotic bacterial pathogens [24].

Salmonella enterica isolates isolated in this work have avian (broiler), sea products (clam) and bovine (milk) sources. All over the world, the most often isolated serovar is *S. Enteritidis* [25]. However, in this study, 42% versus 40% of the 45 isolates

belonged to *S. Enteritidis* and *S. Kentucky* serovars, respectively, suggesting the emergence of *S. Kentucky* serovar in Tunisia.

All isolated strains were serotyped and confirmed as *Salmonella* strains using *invA* gene. The invasion gene *invA* is essential for full virulence in *Salmonella* and it is thought to trigger the internalization required for the invasion of deeper tissues [22].

The PFGE of XbaI-digested genomic DNA from 45 *Salmonella* isolates showed 13 different macrorestriction profiles or clusters (P1 to P13). Heterogeneity was mainly observed within the two major serovars, *S. Enteritidis* and *S. Kentucky*. The distribution of isolates in clusters was done independently of the origin of the broiler samples. In this study, only one isolate belonging to *S. Enteritidis* serovar (O) was obtained from clam and no one from milk. In an earlier investigation, researchers showed that among the 58 seafood associated *Salmonella* serovars, nine were observed but no one belonged to *S. Enteritidis* serovar [2]. However,




Figure 5. Dendrogram showing the percentage of similarity between typable *Salmonella* isolates generated from composite fingerprinting. Genetic similarity between samples in duplicate is 80%. doi:10.1371/journal.pone.0081315.g005

Salmonella Enteritidis serovar has already been reported to survive and grow in fermented milks [26–27].

It is important to note that using the PFGE molecular typing method, heterogeneity was observed within *salmonella* belonging to the same serovar. Serological relatedness did not show correlation with genetic relatedness as previously reported in many studies [28–30]. The discrimination index (DI) of PFGE in this analysis was found to be 0.94. This typing method presented so that a very high discriminatory power (if complying with the conventional 5% level of acceptable probability where a $DI > 0.95$ is desired [31].

The ERIC-PCR of the 45 *Salmonella* isolates also yielded different patterns grouped into 8 clusters (E1 to E8). In this case, the isolates distribution in clusters was done independently of the origin of the broiler samples.

Inter-serovars heterogeneity was observed even within the two *S. Anatum* isolates which were clustered in the PFGE dendrogram. This result proves the effectiveness of this molecular typing method and shows that ERIC-PCR could be useful for subtyping *Salmonella* serovars, where ubiquitous and similar PFGE patterns occur. Similar results were previously reported in many studies

when ERIC-PCR was compared to other molecular typing methods [10–12]. Besides, the accuracy, simplicity and lower cost of ERIC PCR compared to other typing methods enhance its usefulness for *Salmonella* serovars analysis and it has been successfully used for typing many entero-bacteria [32–34].

Recently, a collection of 57 *Salmonella* Kentucky isolates was analyzed by Turki and others using plasmid profiling, PFGE, ribotyping, ERIC-PCR fingerprinting, and Random Amplification of Polymorphic DNA [35]. The authors showed a discriminatory index of 0.647 for PFGE versus 0.903 for ERIC-PCR. However, in the present work, results show that PFGE is more discriminative than ERIC-PCR to differentiate even intraserovars isolates. In fact, the DI of ERIC-PCR analysis was found to be 0.85. As regards discriminatory power alone, this datum shows that the DI obtained by ERIC-PCR is slightly below the acceptable confidence value for interpreting the discrimination level. Thus, PFGE typing method (with a DI of 0.94) is more discriminatory than ERIC-PCR which is insufficient in this case as a single typing method. Our result is in agreement with that of other workers who

reported that PFGE is one of the most reliable techniques for discriminating different serotypes of *Salmonella* [29,36].

Turki and others also indicate that a single method cannot be relied upon for discriminating between *S. Kentucky* strains, and a combination of typing methods such ERIC2 and RAPD2 allows further discrimination [35]. Data from the two molecular typing methods used in the present report were subjected to a composite analysis to determine whether a better serovars clustering could be obtained. The combined PFGE-ERIC-PCR patterns allowed a DI of 0.93 close to that obtained by PFGE and higher than that obtained by ERIC-PCR. In statistical terms, this would provide 93% confidence in the ability to accurately discriminate between two unrelated strains. Results of the combined analysis were highly discriminatory and thus more efficient as reported by Shariat et al. [30].

In the present work, the *Salmonella* search was positive in broiler intestine, liver and carcass. During the slaughter, this pathogen can contaminate carcasses and meat, resulting in a source of food borne illness [37]. Results show different profile patterns between 3 isolates derived from broiler carcass (C31', C32' and C33') suggesting that a vertical transfer could occur. Bacterial contamination can be then spread to millions of chicks within few days via horizontal transfer. On the other hand, *S. Kentucky* isolates belonging to cluster C5 (Figure 5) derived from two different food borne origins and presented the same profile pattern. Finally, isolates derived from clams were always different from those derived from broiler and/or milk which is a further argument in favor of the diversity of *Salmonella* strains contaminating foods.

References

- Hendriksen RS, Vieira AR, Karlsmose S, Wong DMA, Jensen AB, et al. (2011) Global monitoring of *Salmonella* serovar distribution from the World Health Organization Global Foodborne Infections Network Country Data Bank: results of quality assured laboratories from 2001 to 2007. *Foodborne Path Dis* 8: 887–900.
- Bhowmick PB, Srikanth S, Devegowda D, Shekar M, Darshane-Ruwandepika HA, et al. (2012) Serotyping and molecular characterization for study of genetic diversity among seafood associated non typhoidal *Salmonella* serovars. *Indian J Med Res* 135: 371–381.
- Shah DH, Zhou X, Kim HY, Call DR, Guard J (2012) Transposon mutagenesis of *Salmonella enterica* serovar Enteritidis identifies genes that contribute to invasiveness in human and chicken cells and survival in egg albumen. *Infect Immun* 80: 4203–4215.
- Kumar R, Lalitha KV (2012) Digoxigenin-labeled probe-based colony blotting assay for rapid quantification of *Salmonella* serovars in seafood and water. *JAOAC Int* 6: 1652–1655.
- Alali WQ, Hofacre CL, Mathis GF, Faltys G (2013) Effect of essential oil compound on shedding and colonization of *Salmonella enterica* serovar Heidelberg in broilers. *Poult Sci* 92: 836–841.
- Akoachere JFTK, Tanih NF, Ndip LM, Ndip RN (2009) Phenotypic Characterization of *Salmonella Typhimurium* Isolates from Food-animals and Abattoir Drains in Buea, Cameroon. *J Health Popul Nutr* 27: 612–618.
- Kwon YK, Kim A, Kang MS, Her M, Jung BY, et al. (2010) Prevalence and characterization of *Salmonella Gallinarum* in the chicken in Korea during 2000 to 2008. *Poultry Sci* 89: 236–242.
- Pallavi S, Foley SL, Nayak R, Kwon YM (2013) Massively parallel sequencing of enriched target amplicons for high-resolution genotyping of *Salmonella* serovars. *Mol Cell Probes* 27: 80–85.
- Sukhnanand S, Alcaine S, Warnick LD, Su WL, Hof J, et al. (2005) DNA sequence-based subtyping and evolutionary analysis of selected *Salmonella enterica* serotypes. *J Clin Microbiol* 43: 3688–3698.
- Kumar R, Surendran PK, Thampuran N (2008) Molecular fingerprinting of *Salmonella enterica subsp. enterica Typhimurium* and *Salmonella enterica subsp. enterica Derby* isolated from tropical seafood in South India. *Mol Biotechnol* 40: 95–100.
- Lim H, Lee KH, Hong CH, Bahk GJ, Choi WS (2005) Comparison of four molecular typing methods for the differentiation of *Salmonella spp.* *Int J Food Microbiol* 105: 411–418.
- Nath G, Pushpa M, Gulati AK (2010) ERIC PCR and RAPD based fingerprinting of *Salmonella Typhi* strains isolated over a period of two decades. *Infect Genet Evol* 10: 530–536.
- Versalovic J, Koeuth T, LeLupsbi JR (1991) Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res* 19: 6823–6831.
- Stevens A, Kerouanton A, Marault M, Millemann Y, Brisabois A, et al. (2008) Epidemiological analysis of *Salmonella enterica* from beef sampled in the slaughterhouse and retailers in Dakar (Senegal) using pulsed-field gel electrophoresis and antibiotic susceptibility testing. *Int J Food Microbiol* 123: 191–197.
- Barua H, Biswas PK, Olsen KE, Shill SK, Christensen JP. (2013) Molecular characterization of motile serovars of *Salmonella enterica* from broiler and commercial broiler poultry farms in Bangladesh. *Plos One* 8: e57811.
- Zou W, Chen HC, Hise KB, Tang H, Foley SL, et al. (2013) Meta-analysis of pulsed-field gel electrophoresis fingerprints based on a constructed *Salmonella* database. *PLoS One* 8: e59224.
- Barkallah M, Fendri I, Dhieb A, Gharbi Y, Greub G, et al. (2013) First detection of *Wadliia chondrophila* in Africa using SYBR Green real-time PCR on veterinary samples. *Vet Microbiol* 164: 101–107.
- Münch S, Wemerv U, Kinne J, Joseph M, Braun P, et al. (2013) Comparing the presence of different genes in *Salmonella* subspecies I–IV and development of a diagnostic multiplex PCR method for identification of *Salmonella* subspecies. *Berl Munch Tierarztl Wochenschr* 126: 16–24.
- Grimont PAD, Weill FX (2007) Antigenic formulae of the *Salmonella* serovars, 9th ed. Institut Pasteur., Paris, p. 166.
- Ribot EM, Fair MA, Gautom R, Cameron DN, Hunter SB, et al. (2006) Standardization of pulsed-field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathog Dis* 3:59–67.
- Sangdee A, Natphosuk S, Srisathan A, Sangdee K (2013) Development of SCAR Primers Based on a Repetitive DNA Fingerprint for *Escherichia coli* Detection. *J Microbiol* 51: 31–35.
- Torpdahl M, Skov MN, Sandvang D, Baggesen DL (2005) Genotypic characterization of *Salmonella* by multilocus sequence typing, pulsed-field gel electrophoresis and amplified fragment length polymorphism. *J Microbiol Methods* 63: 173–184.
- Gorman R, Adley CC (2004) Characterization of *Salmonella enterica* serotype Typhimurium isolates from human, food and animal sources in the Republic of Ireland. *J Clin Microbiol* 42: 2314–2316.
- Morshed R, Peighambari SM (2010) Drug resistance, plasmid profile and random amplified polymorphic DNA analysis of Iranian isolates of *Salmonella* Enteritidis. *New Microbiol* 33: 47–56.
- Ben Aissa R, Al-Gallas N, Troudi H, Belhadj N, Belhadj A (2007) Trends in *Salmonella enterica* serotypes isolated from human, food, animal, and environment in Tunisia, 1994–2004. *J Infect* 55: 324–339.
- Feresu SB, Nyathi H (1990) Fate of pathogenic and non-pathogenic *Escherichia coli* strains in two fermented milk products. *J Appl Bacteriol* 69: 814–821.
- Mufandaedza J, Viljoen SB, Feresu SB, Gadaga TH (2006) Antimicrobial properties of lactic acid bacteria and yeast-LAB cultures isolated from traditional

Conclusions

In the current work, PFGE and ERIC-PCR were used for subtyping *Salmonella* isolates belonging to different serovars. An analysis of the two typing methods indicated that some of the *Salmonella* isolates were indistinguishable and/or highly related. These isolates were thus grouped into clusters whose number is higher when PFGE typing method was used. The dendrograms showed that PFGE and ERIC-PCR differentiated isolates into grouping that correlated with serovars. A wide variety of profiles was observed between the different serovars. This diversity is particularly important when the sample origins are varied, and even within the same sampling origin.

Acknowledgments

The authors are grateful to Pr. Néji Gharsallah from the University of Sfax for his fruitful discussions. We also thank Dr. Idriss Ben Naila from the Veterinary Center of Sfax for his help.

Author Contributions

Conceived and designed the experiments: IF. Performed the experiments: IF ABH NG MB LK VC. Analyzed the data: IF VC MG RG. Contributed reagents/materials/analysis tools: NG VC. Wrote the paper: IF LK MG RG.

- fermented milk against pathogenic *Escherichia coli* and *Salmonella enteritidis* strains. *Int J Food Microbiol* 108: 147–152.
28. Arguello H, Carvajal A, Naharro G, Arcos M, Rodicio MR, et al. (2013) Serotyping and genotyping of *Salmonella* in slaughter pigs, from farm to cutting plant, with a focus on the slaughter process. *Int J Food Microbiol* 161: 44–52.
 29. Chen MH, Hwang WZ, Wang SW, Shih YC, Tsen HY (2011) Pulsed field electrophoresis (PFGE) analysis for multidrug resistant *Salmonella enterica* serovar Schwarzengrund isolates collected in six years (2000_2005) from retail chicken meat in Taiwan. *Food Microbiol* 28: 399–405.
 30. Shariat N, DiMarzio MJ, Yin S, Dettinger L, Sandt CH, et al. (2013) The combination of CRISPR-MVLST and PFGE provides increased discriminatory power for differentiating human clinical isolates of *Salmonella enterica* subsp. *Enterica* serovar *Enteritidis*. *Food Microbiol* 34: 164–173.
 31. Struelens MJ (1996) Consensus guidelines for appropriate use and evaluation of microbial epidemiologic typing systems. *Clin Microbiol Infect* 2: 2–11.
 32. Adzitey F, Rahmat Ali GR, Huda N, Cogan T, Corry J (2013) Prevalence, antibiotic resistance and genetic diversity of *Listeria monocytogenes* isolated from ducks, their rearing and processing environments in Penang, Malaysia. *Food Control* 32: 607–614.
 33. Rantsiou K, Alessandria V, Coccolin L (2012) Prevalence of Shiga toxin-producing *Escherichia coli* in food products of animal origin as determined by molecular methods. *Int J Food Microbiol* 154: 37–43.
 34. Yuan W, Chai TJ, Miao ZN (2010) ERIC-PCR identification of the spread of airborne *Escherichia coli* in pig houses. *Sci Total Environ* 408: 1446–1450.
 35. Abbassi-Ghozzi I, Jaouani A, Aissa RB, Martinez-Urtaza J, Boudabous A, et al. (2011) Antimicrobial resistance and molecular analysis of non-typhoidal *Salmonella* isolates from human in Tunisia. *Pathol Biol* 59: 207–212.
 36. Turki Y, Mehri I, Fhoula I, Hassen A, Ouzari H (2013) Comparison of five molecular subtyping methods for differentiation of *Salmonella* Kentucky isolates in Tunisia. *World J Microbiol Biotechnol* DOI 10.1007/s11274-013-1414-1.
 37. Centers of Disease Control (2010) *Salmonella* Montevideo infections associated with salami products made with contaminated imported black and red pepper_United States, July 2009-April 2010. *MMWR*, 59: 1647–1650.