

# Aberrant Interference of Auditory Negative Words on Attention in Patients with Schizophrenia

Norichika Iwashiro<sup>1\*</sup>, Noriaki Yahata<sup>1,2</sup>, Yu Kawamuro<sup>3</sup>, Kiyoto Kasai<sup>1,4</sup>, Hidenori Yamasue<sup>1\*</sup>

**1** Department of Neuropsychiatry, Graduate School of Medicine, The University of Tokyo, Bunkyo-ku, Tokyo, Japan, **2** Global Center of Excellence (COE) Program, The University of Tokyo, Bunkyo-ku, Tokyo, Japan, **3** Takada-Nishishiro Hospital, Jyoetsu-shi, Niigata, Japan, **4** JST, National Bioscience Database Center (NBDC), Chiyoda-ku, Tokyo, Japan

## Abstract

Previous research suggests that deficits in attention-emotion interaction are implicated in schizophrenia symptoms. Although disruption in auditory processing is crucial in the pathophysiology of schizophrenia, deficits in interaction between emotional processing of auditorily presented language stimuli and auditory attention have not yet been clarified. To address this issue, the current study used a dichotic listening task to examine 22 patients with schizophrenia and 24 age-, sex-, parental socioeconomic background-, handedness-, dexterous ear-, and intelligence quotient-matched healthy controls. The participants completed a word recognition task on the attended side in which a word with emotionally valenced content (negative/positive/neutral) was presented to one ear and a different neutral word was presented to the other ear. Participants selectively attended to either ear. In the control subjects, presentation of negative but not positive word stimuli provoked a significantly prolonged reaction time compared with presentation of neutral word stimuli. This interference effect for negative words existed whether or not subjects directed attention to the negative words. This interference effect was significantly smaller in the patients with schizophrenia than in the healthy controls. Furthermore, the smaller interference effect was significantly correlated with severe positive symptoms and delusional behavior in the patients with schizophrenia. The present findings suggest that aberrant interaction between semantic processing of negative emotional content and auditory attention plays a role in production of positive symptoms in schizophrenia. (224 words)

**Citation:** Iwashiro N, Yahata N, Kawamuro Y, Kasai K, Yamasue H (2013) Aberrant Interference of Auditory Negative Words on Attention in Patients with Schizophrenia. *PLoS ONE* 8(12): e83201. doi:10.1371/journal.pone.0083201

**Editor:** Joel Snyder, UNLV, United States of America

**Received:** July 3, 2013; **Accepted:** October 31, 2013; **Published:** December 23, 2013

**Copyright:** © 2013 Iwashiro et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** Part of this study was supported by KAKENHI (22689034 to H.Y.), the Global Center of Excellence (COE) Program "Comprehensive Center of Education and Research for Chemical Biology of the Diseases" (N.Y.) and National Bioscience Database Center (NBDC) of Japan Science and Technology Agency (JST). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: iwashiro-ty@umin.ac.jp (NI); yamasue-ty@umin.ac.jp (HY)

## Introduction

Disconnection between different elements of brain networks or cognitive domains characterizes the schizophrenia diathesis [1,2,3]. For example, deficits in attention-emotion interaction as well as those in each domain have repeatedly been investigated in patients with schizophrenia [4,5,6,7,8,9,10,11]. Previous studies have shown that presentations of emotionally negative stimuli such as angry face or prosody interfere with attention and lead to slower response in healthy subjects [12,13,14,15]. Patients with schizophrenia show significant deviations in this interference effect of emotionally negative stimuli [4,7,9].

Dysfunctional language-related information processing such as aberrant semantic processing plays a role in the pathophysiology of schizophrenia [16,17,18], particularly in the formation of positive symptoms such as delusions [19,20,21]. Therefore, visually presenting language stimuli has often been utilized to examine the interaction between attention and emotional processing in patients with schizophrenia (reviewed in [22]). However, disruption in auditory information processing is considered crucial in the pathophysiology of schizophrenia, with an emphasis on auditory hallucinations in diagnostic criteria [23,24] and with various cognitive deficits associated with auditory

processing [25,26,27]. Taken together, investigating the interaction between emotional processing of auditorily presented language stimuli and attention on auditory perception would bring novel and significant findings in the pathophysiology of psychotic symptoms in schizophrenia.

Previous studies [12,13] investigated interaction between auditory spatial attention and emotional processing utilizing dichotic listening paradigm and showed prolonged reaction time when angry prosody presented in left ear but only when this side was to-be-attended in healthy subjects. The current dichotic listening study utilized semantically emotional words instead of emotional prosody to approach the pathophysiology of aberrant semantic processing in schizophrenia. Based on the previous prosody studies, we predicted that 1) response time would be longer when negative but not positive words were auditorily presented than when only neutral ones were presented in healthy subjects; 2) subjects with schizophrenia would show abnormality in this interference effect of auditory negative words compared with healthy subjects; and 3) the abnormality in interference effect of auditory negative words would be associated with severity of psychotic symptoms in patients with schizophrenia.

## Materials and Methods

### Ethics statement

The ethics committee of Takada-Nishishiro Hospital approved this study. Written informed consent was obtained from all participants after they were given a complete explanation of the study as approved by the ethics committee. For participants below the age of 20 years, we obtained written informed consent from his or her parents as well as from the participants themselves. All participants belong to patient group were determined by their psychiatrist in charge to have the mental and intellectual capacity to give written informed consent prior to participation in the study. If a participant or their parents could not understand these issues during explanation, s/he was not asked to participate in this research.

### Participants

Forty-six Japanese adults participated in this study. Of these, 22 in- and out-patients with schizophrenia were recruited from the Takada-Nishishiro Hospital, Japan. Diagnosis was confirmed as schizophrenia with the Structured Clinical Interview for DSM-IV Axis I Disorder Clinical Version [24]. Psychiatric symptoms were evaluated using the Positive and Negative Syndrome Scale (PANSS) [28] just before the current psychological experiment. Delusional behavior scores were calculated from subscales of the PANSS. The calculation algorithm was P1 (delusion) +P5 (grandiosity) +P6 (suspiciousness) +G9 (unusual thought content) [29,30,31]. All clinical evaluations were conducted by a psychiatrist (N.I.) fully trained to maintain reliability and consistency in diagnoses and evaluation of symptom severity. Twenty of the 22 patients were categorized as paranoid subtype; the other two were categorized as disorganized subtype. Intelligence quotient (IQ) was evaluated using the Japanese version of the National Adult Reading Test [32,33,34]. Based on previous studies, dexterous ear was decided using four question items (Appendix S1) [35,36,37].

Twenty-four healthy subjects were employed as controls. The control group was matched to the subjects with schizophrenia in age, sex ratio, parental socioeconomic status (SES) [38], handedness [39], dexterous ear, and estimated premorbid IQ. The controls were screened for neuropsychiatric disorders using the Structured Clinical Interview for DSM-IV Axis I Disorder Non-Patient Edition [23,40].

Exclusion criteria for both groups were current or past neurological or audiological illness, traumatic brain injury with any known cognitive consequences or loss of consciousness for more than 5 minutes, history of electroconvulsive therapy, and substance abuse or addiction. To screen for participants with potential auditory abnormalities, we tested hearing ability in both ears. The test comprised a 1000 Hz tone at 30 dB and a 4000 Hz tone at 40 dB tone, as measured by audiometer (Minato; <http://www.minato-med.co.jp/products/inspect/amc1.html>). This test was taken from the Japanese Ordinance on Industrial Safety and Health No. 44 (Ordinance of the Ministry of Health, Labour and Welfare) (e.g. <https://www.rouki.or.jp/furukawa/modules/mydownloads/jyouhou/ryoukin.pdf>). An additional exclusion criterion for the control group was history of psychiatric disease in self or a family history of Axis I disorder in a first-degree relative. The ethics committee of Takada-Nishishiro Hospital approved this study. After a complete explanation of the study, written informed consent was obtained from every individual (Table 1).

### Experimental procedure

We selected three word categories of emotional valence (20 different negative/60 different neutral/20 different positive words) from the Affective Norms for English Words (ANEW; [41]) and translated them to Japanese (Appendix S2). All the words were nouns and consisted of four morae in Japanese (Appendix S3). For validation, 20 healthy adults (10 male/10 female) not participating in the current experiment rated these words in terms of their valence, arousal, and dominance by the same 9-point Likert scale as in the ANEW [41]. Word frequency was quoted from a Japanese database [42], and number of orthographic neighbours was calculated from a Japanese thesaurus [43]. The current experiment used three types of word lists that significantly differed only in emotional valence (negative/neutral/positive) (ANOVA;  $p < 0.001$ ), but were equal in all other language features such as arousal ( $p = 0.15$ ), dominance ( $p = 0.72$ ), word frequency ( $p = 0.25$ ), and number of orthographic neighbours ( $p = 0.22$ ) (ref. [41,44]). All the words were transformed to auditory stimuli pronounced by synthesized voice in neutral prosody using VoiceText® (HOYA; <http://voicetext.jp/>). We then processed each auditory stimulus to equalize the duration of the utterances into 700 milliseconds, which were spoken in a naturalistic way using Sound Forge Pro 10® (Sony Creative Software; <http://www.sonycreativesoftware.com/soundforgepro>). The amplitude (negative; positive; neutral, mean/SD/range =  $-77.6/6.3/-90.3$  to  $-67.4; -79.6/7.9/-66.8$  to  $-90.3; -77.4/7.0/-65.2$  to  $-90.3$  dB) and pitch (negative; positive; neutral, mean/SD/range =  $925.7/640.2/288.5-2679.0; 870.4/538.4/374.2-1925.7; 1035.3/646.8/295.7-2468.4$  Hz) did not significantly differ among the negative, positive, and neutral word categories ( $F[2,89] = 0.69, p = 0.50$  and  $F[2,97] = 0.62, p = 0.54$  respectively). In addition, the mean acoustic energies (RMS) of the two words, which were presented simultaneously in a dichotic paradigm, were always matched using Mitsyn® (WLH) and CoolEdit Pro® (Syntrillium Software Corporation), according to previous studies [12,13,45,46].

The task design was developed based on that in a previous study (ref. [12]), although that study used emotional prosody in contrast to emotional content of words. Participants responded by pressing a button while listening to a pair of words. To manipulate voluntary attention orthogonally to emotional words, we used a dichotic listening paradigm in which stimuli matched for mean acoustic energy and duration were presented simultaneously to both ears (negative/neutral/positive and neutral or neutral and negative/neutral/positive, on the right and left sides, respectively) in random order (Figure 1). Each stimulus condition (negative-neutral, neutral-negative, positive-neutral, neutral-positive, and neutral-neutral, on the right-left sides, respectively) had 40 trials. The subjects attended at the right ear for half of the 40 trials and the left ear for the other half, for each condition. Subjects were asked to attend at the right or left ear in a pseudo-random order. This information was given by instructions shown on the screen before the presentation of the auditory stimuli. Each word appeared four times in the experimental session and all the trials were presented in random order. Participants were required to selectively attend to either the left or right ear, and select the word heard in the attended ear from four presented words on the screen as quickly as possible. For each event, the time taken to select one word by pressing a button after the auditory stimuli was presented and the number of correct responses were recorded automatically in E-prime 2.0 (<http://www.ipsjapan.com/EPRIME.htm>) as response time (RT) and correct ratio (CR), respectively.

**Table 1.** Subject characteristics and symptom scores.

	Patients with Schizophrenia (n = 22)		Controls (n = 24)		T-test	
	Mean	SD	Mean	SD	T-value	P-value
Age (Range)	31.6 (20–40)	5.2	29.9 (18–39)	5.7	1.1	0.28
Male/Female	11/11		11/13		0.08 ( $\chi^2$ )	0.78 ( $\chi^2$ )
SES <sup>a</sup>	3.9	1.0	2.7	0.8	4.7	0.00
Parental SES	3.0	0.8	3.1	0.7	−0.4	0.70
Years of education	13.5	1.8	14.4	1.4	1.72	0.09
IQ (JART25) <sup>b</sup>	96.7	10.0	99.3	8.1	−1.0	0.34
Handedness <sup>c</sup>	89.5	34.5	98.0	4.8	−1.2	0.24
Dexterous ear (right/mixed/left)	18/1/3		16/5/3		2.7 ( $\chi^2$ )	0.26 ( $\chi^2$ )
Neuroleptic dose <sup>d</sup> (mg/day)	709.4	641.8				
Onset of illness (years)	24.3	6.3				
Duration of illness (months)	104.1	80.5				
PANSS <sup>e</sup> Positive symptoms	17.0	2.7				
PANSS <sup>e</sup> Negative symptoms	23.5	4.6				
PANSS <sup>e</sup> General psychopathology	42.5	6.9				
PANSS <sup>e</sup> Delusional behavior	9.9	1.9				

<sup>a</sup>Socioeconomic status, assessed using the Hollingshead scale. Higher scores indicate lower status;

<sup>b</sup>Estimated from scores on the Japanese Adult Reading Test;

<sup>c</sup>Assessed using the Edinburgh Inventory. >0 indicates right-handed;

<sup>d</sup>Based on chlorpromazine equivalents;

<sup>e</sup>Positive and Negative Syndrome Scale.

doi:10.1371/journal.pone.0083201.t001

## Statistical analysis

**Effects of emotional stimuli, their laterality, and attention in healthy subjects and patients with schizophrenia.** First, to verify previous findings that response to negative stimuli was different from the response to neutral or positive stimuli in healthy subjects [12,14], we examined CR and RT for negative-neutral, positive-neutral, and neutral-neutral word pairs in the healthy participants and the patients with schizophrenia. We performed four repeated measures ANOVAs with two within-subject factors (Stimulus-Type: Negative-Neutral/Neutral-Negative/Positive-Neutral/Neutral-Positive/Neutral-Neutral; Attention-Side: right/left). Two had CR as the dependent variable (one for healthy subjects and one for patients with schizophrenia) and two had RT as the dependent variable (one for healthy subjects and one for patients with schizophrenia). The statistical model was based on that in a previous study (ref. [12]). The threshold for statistical significance was set at  $p < 0.05$ .

**Comparisons between subjects with schizophrenia and healthy controls.** Analyses for comparisons between patients with schizophrenia and controls were confined to an index, which revealed significant interference of emotional valence in healthy subjects (i.e. RT, see Results). Based on prior findings describing a general slowness in RT in patients with schizophrenia [47,48,49], we treated interference indices as dependent variables in the main analysis. Interference indices allow a more specific assessment of the effect of emotional valence on inhibitory processes by minimizing the effect of general slowness. They were calculated by a widely employed formula:


$$\text{Interference index} = (\text{RT for negative/positive and neutral}) \\ - (\text{RT for neutral and neutral}).$$

Thus, the condition of neutral and neutral word pairs served as baseline [7,50].

A repeated measures ANOVA with one between-subject factor (Diagnosis: schizophrenia/control) and two within-subject factors (Stimulus-Type: Negative-Neutral/Neutral-Negative/Positive-Neutral/Neutral-Positive; Attention-Side: right/left) with interference indices as dependent variables was performed. If a significant Diagnosis-by-Stimulus-Type or Diagnosis-by-Stimulus-Type-by-Attention-Side was found, follow-up analyses using repeated measures ANOVAs separately for each Stimulus-Type were performed. The threshold for statistical significance was set at  $p < 0.05$ .

**Correlations between interference indices and clinical measurements.** Associations between interference indices showing significant group differences and positive symptoms, negative symptoms and general psychopathology on the PANSS, and delusional behavior scores were further tested with Spearman's rank correlation in the patient group. Selective attention to threatening information may lead an individual to form paranoid delusions about the environment [48,51,52]. Thus we hypothesized that abnormal interference indices for negative-neutral word pairs would correlate with the severity of positive symptoms and delusional behavior in schizophrenia. Therefore, statistical significance was set at  $p < 0.05$  for these correlations. In contrast, because no theoretical hypothesis about the correlations between severity of negative symptoms, general psychopathology, and interference indices exists, statistical significance was set at  $p < 0.025$  after Bonferroni correction for two correlations (one condition  $\times$  two clinical measurements).

Additionally, correlations between interference indices and potential confounding factors, including age, self SES, parental SES, age of onset, duration of illness, and dose of neuroleptics, were tested separately in each group using Spearman's rank



**Figure 1. Experimental paradigm.** Participants were asked to selectively attend to either the right- or left-sided voice stimulus within a pair. Then, they were required to select the word heard at the attended ear from four presented candidate words on the screen as quickly as possible. Orthogonally to the task demands, voices could have a semantically different emotional valence, either neutral on both sides, negative (or positive) on the attended side and neutral on the other side, or vice versa, neutral on the attended side and negative (or positive) on the other side.  
doi:10.1371/journal.pone.0083201.g001

correlation. In contrast, because no theoretical hypothesis about the correlations between the other clinical measurements and interference indices exists, the threshold for statistical significance was set at  $p < 0.0056$  (Bonferroni correction for nine correlations [six for schizophrenia group {one condition  $\times$  six clinical measures}; three for control group {one condition  $\times$  three clinical measures}]).

## Results

### Effects of emotional stimuli, their laterality, and attention in healthy controls and patients with schizophrenia

For RT, the repeated measures ANOVA showed a significant main effect of Stimulus-Type ( $F[4,92] = 3.89$ ,  $p = 0.006$ ) with no significant interaction ( $F[4,92] = 1.93$ ,  $p = 0.11$ ). Post-hoc paired t-tests revealed that RT for Negative-Neutral and Neutral-Negative (right-left side) ( $t[23] = 3.25$ ,  $p = 0.0035$ ;  $t[23] = 2.40$ ,  $p = 0.025$ , respectively) were significantly longer than that for Neutral-Neutral word pairs, irrespective of direction of attention in healthy controls (Table 2, Figure 2). The same result was obtained if one extreme outlier with the maximum RT for Negative-Neutral stimuli was excluded (main ANOVA,  $p = 0.03$ ; post hoc paired t-test,  $p = 0.00048$ ). In contrast, no significant effects were found in schizophrenia group ( $p > 0.16$ ). Repeated measures ANOVAs showed no significant main effects or interactions for CR

( $p > 0.64$ ). A significant interaction between Attention-Side and Stimulus-Type was found in the patient group ( $F[4,84] = 4.92$ ,  $p = 0.001$ ). Post-hoc paired t-tests revealed that the CRs for Neutral-Negative ( $t[21] = 2.51$ ,  $p = 0.02$ ), Positive-Neutral ( $t[21] = 2.53$ ,  $p = 0.02$ ) and Neutral-Positive ( $t[21] = 2.59$ ,  $p = 0.017$ ) were higher than CR for Neutral-Neutral word pairs when patients attended to the right ear. The CRs for Negative-Neutral ( $t[21] = 2.75$ ,  $p = 0.012$ ) and Neutral-Neutral ( $t[21] = 3.18$ ,  $p = 0.005$ ) were higher than CR for Neutral-Negative word pairs when patients attended at left ear.

### Comparisons between subjects with schizophrenia and controls


The repeated measures ANOVA showed a significant main effect of stimulus-type ( $F[3,132] = 2.69$ ,  $p = 0.049$ ) and interaction between Diagnosis and Stimulus-Type ( $F[3,132] = 2.83$ ,  $p = 0.041$ ), and no other effect and interaction ( $p > 0.35$ ). Post-hoc paired t-tests revealed that interference indices for Negative-Neutral ( $t[45] = 2.47$ ,  $p = 0.018$ ), Neutral-Negative ( $t[45] = 2.30$ ,  $p = 0.026$ ) and Positive-Neutral ( $t[45] = 2.14$ ,  $p = 0.038$ ) (right-left side) were significantly longer than that for Neutral-Positive (right-left side) word pairs, irrespective of direction of attention. For Negative-Neutral (right-left side) stimulus-type, the post-hoc repeated measures ANOVA showed a significant main effect of Diagnosis ( $F[1,44] = 5.39$ ,  $p = 0.025$ ). For other stimulus types, there was no significant main effect of Diagnosis or interactions. This analysis demonstrated that the interference index in patients with schizophrenia was significantly smaller than that in controls when negative words were presented to the right ear, irrespective of attention side (Table 3, Figure 3). The statistical conclusions remained the same when the one left-handed subject in the schizophrenia group was excluded and when analysis of covariance including dexterous ear as a covariate was employed. Furthermore, if extreme outliers (i.e. one patients with the minimum interference index and one control with the maximum index for Negative-Neutral stimuli) were excluded, there remains a tendency of the same statistical conclusion (main ANOVA,  $p = 0.070$ ; post hoc ANOVA,  $p = 0.066$ ).

### Correlational analysis

Because no main effect of Attention-Side and no interaction between it and any other factors were found, means of the interference indices for participants' attending on right- and left-ear were used. The interference index for negative-neutral pairs when negative words were presented to the right ear irrespective of attention side significantly negatively correlated with the PANSS positive symptoms severity ( $\rho = -0.43$ ,  $p = 0.044$ ) and delusional behavior ( $\rho = -0.45$ ,  $p = 0.035$ ) (Figure 4), but was not correlated with PANSS negative symptoms ( $p = 0.39 > 0.05/2$ ) or general psychopathology ( $p = 0.14 > 0.05/2$ ). There was no significant correlation between the interference index and potential confounds in each group ( $p \geq 0.10 > 0.05/9$ ).

## Discussion

To our knowledge, the present study is the first to demonstrate that response time was longer when emotionally negative but not positive words were auditorily presented than when only neutral ones were presented. Furthermore, for the first time, a significantly smaller interference effect of negative emotional words on auditory attention was demonstrated in patients with schizophrenia compared with healthy controls, while interference of positive emotional words was not significantly different between patients and controls. Furthermore, reduced interference of negative



**Figure 2. Effect of auditory negative words on response time in healthy subjects and patients with schizophrenia.** Response times for Negative-Neutral, Neutral-Negative stimuli on the right-left side were significantly longer than that for Neutral-Neutral word pairs, irrespective of direction of attention in the healthy controls ( $p=0.0035$ ,  $p=0.025$ , respectively). In contrast, response time for Positive-Neutral ( $p=0.11$ ), Neutral-Positive ( $p=0.12$ ) word pairs on the right-left side was not significantly longer than that for Neutral-Neutral word pairs. doi:10.1371/journal.pone.0083201.g002


**Table 2. Reaction time (msec).**

Healthy subjects (n = 24)				Paired t-test (post-hoc)				Patients with schizophrenia (n = 22)					
Stimulus-Type	Attention-Side		Attention-Side		Relative to each stimulus-type				Stimulus-Type		Attention-Side		
	Mean	SD	Mean	SD	Neu-Neg	Pos-Neu	Neu-Pos	Neu-Neu	Mean	SD	Mean	SD	
Neg-Neu	2125.0	633.9	R	2178.9	T-value	2.16	1.83	1.89	3.25	3004.4	1471.1	3048.9	1394.7
			L	2071.1									
Neu-Neg	2020.2	572.0	R	1988.3	T-value	0.16	0.48	2.4	3092.2	1624.3	3092.4	1890.9	
			L	2052.1									P-value
Pos-Neu	2012.2	600.9	R	2049.0	T-value	0.33	1.67	3062.8	1551.4	3066.3	1599.2		
			L	1975.4								P-value	0.74
Neu-Pos	1998.7	543.1	R	1966.8	T-value	1.63	2932.6	1473.3	2999.5	1587.8			
			L	2030.6							P-value	0.12	2865.7
Neu-Neu	1913.7	435.7	R	1884.7	T-value	3017.8	1640.9	3021.8	1708.0				
			L	1942.8						3013.7	1593.0		

ANOVA						
Effect	F	P	Effect Size <sup>a</sup>	F	P	Effect Size <sup>a</sup>
Stimulus-Type	3.89	0.01	0.41	1.71	0.16	0.28
Attention-Side	0.00	0.99	0.00	0.44	0.52	0.14
Stimulus-Type × Attention-Side	1.93	0.11	0.29	0.19	0.94	0.10

Abbreviations: Neg-Neu/Neu-Neg/Pos-Neu/Neu-Pos/Neu-Neu, Negative-Neutral/Neutral-Negative/Positive-Neutral/Neutral-Positive/Neutral-Neutral word pair (right ear-left ear). R/L, Right/Left; \*Statistically significant; <sup>a</sup>Cohen defines fs of 0.1, 0.25, 0.4 as small, medium, large, respectively. doi:10.1371/journal.pone.0083201.t002



**Figure 3. Aberrant interference of auditory negative words in patients with schizophrenia.** The interference index in patients with schizophrenia was significantly smaller compared with healthy controls when negative words were presented at the right ear, irrespective of attention side ( $p = 0.025$ ). Because no main effect of Attention-Side and no interaction between Attention-Side and any other factors were found, the means of the interference indices for attending to the right and left ear are presented. doi:10.1371/journal.pone.0083201.g003


**Table 3.** Comparisons of Interference indices between the patients and controls.

Patients with schizophrenia (n = 22)				Controls (n = 24)				Repeated measures ANOVA (post-hoc)	
Stimulus-Type	Attention-Side			Stimulus-Type	Attention-Side			Group	P
	Mean (msec)	SD	Mean (msec)		SD	Mean (msec)	SD		
Neg-Neu	-13.4	69.9	R 27.1	211.3	66.9	294.3	341.5	F	5.39
			L -53.9						
Neu-Neg	74.4	67.7	R 70.6	106.5	64.8	103.7	277.0	F	0.12
			L 78.2						
Pos-Neu	45.0	68.7	R 44.5	98.5	65.8	164.4	328.9	F	0.32
			L 45.6						
Neu-Pos	-85.2	65.6	R -22.3	85.0	62.9	82.2	400.9	F	3.51
			L -148.0						

ANOVA			
Effect	F	P	Effect Size <sup>a</sup>
Stimulus-Type	2.69	0.05	0.25
Attention-Side	0.89	0.35	0.14
Group	2.19	0.15	0.22
Stimulus-Type × Attention-Side	0.47	0.70	0.11
Attention-Side × Group	0.03	0.86	0.03
Stimulus-Type × Attention-Side × Group	0.55	0.65	0.11

Abbreviations: Neg-Neu/Neu-Neg/Pos-Neu/Neu-Pos, Negative-Neutral/Neutral-Negative/Positive-Neutral/Neutral-Positive word pair (right ear-left ear).  
 R/L, Right/Left;  
 \*Statistically significant;  
<sup>a</sup>Cohen defines fs of 0.1, 0.25, 0.4 as small, medium, large, respectively.  
 doi:10.1371/journal.pone.0083201.t003



**Figure 4. Relationships between the aberrant interference and positive symptoms, delusional behavior.** Scatter plots depict correlations between the interference index for negative-right/neutral-left word pairs irrespective of attention side and severity of positive symptoms in the Positive and Negative Syndrome Scale (PANSS) ( $\rho = -0.43$ ,  $p = 0.044$ ) (a) and severity of delusional behavior ( $\rho = -0.45$ ,  $p = 0.035$ ) (b) in the patients with schizophrenia.  
doi:10.1371/journal.pone.0083201.g004

auditory words significantly correlated with severe positive symptoms and delusional behavior in patients.

We demonstrated that auditorily presented negative words were semantically processed whether they were or were not the focus of attention in healthy subjects. This interference effect of implicitly processed negative emotional stimuli has been observed, similar to visually presented negative words [15,53]. The current results are consistent with previous findings that detection of potential threats, such as angry faces [14] or angry prosody [12], occurs even when they are not initially the focus of attention.

With respect to the laterality effect for the attended ear, our findings are partially inconsistent with those of previous dichotic listening studies [12,13,46,54,55]. For instance, in a paper by Hugdahl et al., the authors used simple consonant–vowel syllables and found that subjects tended to answer the sound from the right ear more often than that from left ear when subjects were not forced attention to either of ear [55]. However, the current study forced subjects to attend right or left ear. The previous studies that also forced subjects to attend right or left ear similarly failed to find laterality effect on CR [12,13,46]. Although one previous study [54] reported laterality effect on CR utilizing the similar experimental design as these studies [12,13,46] in participants predominantly consists of males. The current study as well as the previous studies [12,13,46] included a number of female subjects. It has been suggested that the emotional process is less lateralized in female than male (reviewed in [56]). Unlike previous studies [12,13,46,54], we also did not find significant laterality effects on the RT. Several differences between the current and previous dichotic listening tasks might explain this discrepancy in the findings. First, our study investigated the effect of semantic contents of emotional words on auditory attention, while these previous studies investigated that of emotional prosody [12,13,46,54]. The processes underlying the semantic aspects of verbal stimuli may not be affected by attention-side so much as by the processing of prosody. Second, in the current study, we required the subjects to select target words from four words shown

on a screen, while in previous studies, the subjects were simply asked to judge the gender of a voice by pressing a button [12,13,46,54]. Thus, the current study was more demanding in terms of reading speed and working memory than were these previous studies. IQ and years of education did not differ between the patient and control groups. Furthermore, we employed interference indices, which treat RTs for neutral-neutral stimuli as a baseline. This allowed us to minimize the individual differences in these factors that were unrelated to emotional valence [7,50], although the possibility that these factors confounded the laterality effect of attention-side cannot be completely ruled out. We observed small to moderate effect sizes for interaction between Attention-Side and Stimulus-Type on RT in the healthy subjects, effect of Attention-Side on RT in the patients with schizophrenia and interference indices across groups (Table 2 and 3). It is also possible that we did not find laterality effects due to insufficient statistical power for these indices.

In patients with schizophrenia, this interference effect of auditory negative words was significantly smaller than in healthy individuals. Most previous studies using an emotional Stroop task and investigating attention-emotion interaction by visually presenting emotional words have shown increased interference of negative words in patients with schizophrenia compared with healthy controls [22,57]. Another study demonstrated that patients with schizophrenia were unable to select nonsalient over salient visually presented stimuli [58]. The apparent discrepancy between the current and previous studies might derive from the difference in the modality used in the task, with emotional words presented auditorily in the current study. Indeed, significantly less interference of negative emotional stimuli in other modalities such as face, prosody, or odour have been shown in patients with schizophrenia compared with healthy controls [9,59,60]. A previous study reported prolonged RTs for emotional prosody in patients with schizophrenia compared with healthy controls. However, this study did not employ interference indices, despite the observation of prolonged RTs (which were observed even for neutral stimuli) in

patients with schizophrenia [54]. Another possible explanation for the discrepancy is that less interference in patients with schizophrenia was attributable to general deficits in emotional recognition [61,62], although the reduced interference was not observed to positive words in either ear or negative words in left ear.

The smaller interference effect when negative words were presented to the right ear might reflect functional and structural left-lateralized abnormalities in brain regions related to language processing, including the inferior frontal gyrus and superior temporal gyrus in patients with schizophrenia (reviewed in [63]). Although it has been suggested that emotional semantic stimuli are predominantly processed in right prefrontal areas [64,65], semantic processing itself is thought to be dominant in the left hemisphere [66]. In particular, left lateralized abnormalities have been shown in brain activities elicited by semantic contents of auditorily presented word stimuli [67,68,69,70]. Deficits in these brain regions have been associated with the formation of psychotic symptoms, especially delusions, through abnormal semantic processing [19,20,21,29,31]. Although speculative, the current correlation between less interference of negative words and severe psychotic symptoms and delusional behavior infers that less interference of negative words when they were presented to the right ear resulted from deficits in the left language-related brain regions in patients with schizophrenia. However, neuroimaging studies are needed to examine the neural substrates related to the smaller interference effect of auditory negative words in patients with schizophrenia. Our finding that reduced interference correlated with severe positive symptoms in patients with schizophrenia might be related to the altered sensorimotor gating [71] for negative word stimuli. Sensorimotor gating is defined as the pre-attentive ability of the brain to modulate its sensitivity to an incoming stimulus, and is hypothesized to be a protective mechanism that prevents sensory overload of higher brain functions by filtering out the irrelevant sensory input [71]. Deficits in sensorimotor gating could lead to perceptual and attentional impairments associated with schizophrenia [71,72] and the associated positive symptoms [73].

We must address the methodological limitations of our study. First, the sample size was relatively small. Results should be

replicated in future studies with larger sample sizes, although previous studies investigating interaction between visual emotional words and attention in similar sample sizes [4,7,47]. Second, participants included patients with chronic schizophrenia taking antipsychotic medications. Interference indices did not show any significant correlation with dose of neuroleptics or duration of illness in the study participants. However, effects of chronic illness (reviewed in [74]) and antipsychotic medications (reviewed in [75]) on the present findings cannot be totally ruled out. Future studies should employ patients with first episode schizophrenia to minimize effects of chronicity and medications.

In summary, the present study showed that the interference effect of negative words on auditory attention was abnormally smaller in patients with schizophrenia compared with healthy subjects, and was correlated with severe positive symptoms and delusional behavior. The current findings indicate that abnormality in the interaction between semantic-based emotional processing and auditory attention plays a role in the pathophysiology of psychotic symptoms such as delusions in schizophrenia.

## Supporting Information

**Appendix S1 Four question items used to decide the dextrous ear.** These items were based on those in previous literature (Brybaert, 1994; Coren, 1993; Polemikos and Palaeliou, 2000). (DOCX)

**Appendix S2 Words list in original English from the Affective Norms for English Words (ANEW; Bradley and Lang, 1999).** (In alphabetical order). (DOCX)

**Appendix S3 Japanese words list to show the Japanese four morae.** (In alphabetical order). (DOCX)

## Author Contributions

Conceived and designed the experiments: NI NY HY. Performed the experiments: NI YK. Analyzed the data: NI HY. Wrote the paper: NI HY KK.

## References

- Ellison-Wright I, Bullmore E (2009) Meta-analysis of diffusion tensor imaging studies in schizophrenia. *Schizophr Res* 108: 3–10.
- Friston KJ (1998) The disconnection hypothesis. *Schizophr Res* 30: 115–125.
- Jung WH, Jang JH, Shin NY, Kim SN, Choi CH, et al. (2012) Regional brain atrophy and functional disconnection in Broca's area in individuals at ultra-high risk for psychosis and schizophrenia. *PLoS One* 7: e51975.
- Besnier N, Kaladjian A, Mazzola-Pomicino P, Adida M, Fakra E, et al. (2011) Differential Responses to Emotional Interference in Paranoid Schizophrenia and Bipolar Mania. *Psychopathology* 44: 1–11.
- Dickinson D, Ramsey ME, Gold JM (2007) Overlooking the obvious: a meta-analytic comparison of digit symbol coding tasks and other cognitive measures in schizophrenia. *Arch Gen Psychiatry* 64: 532–542.
- Irani F, Seligman S, Kamath V, Kohler C, Gur RC (2012) A meta-analysis of emotion perception and functional outcomes in schizophrenia. *Schizophr Res* 137: 203–211.
- Roux P, Christophe A, Passerieux C (2010) The emotional paradox Dissociation between explicit and implicit processing of emotional prosody in schizophrenia. *Neuropsychologia* 48: 3642–3649.
- Saykin AJ, Shtasel DL, Gur RE, Kester DB, Mozley LH, et al. (1994) Neuropsychological deficits in neuroleptic naive patients with first-episode schizophrenia. *Arch Gen Psychiatry* 51: 124–131.
- Schneider F, Koch K, Reske M, Kellermann T, Seiferth N, et al. (2006) Interaction of negative olfactory stimulation and working memory in schizophrenia patients: Development and evaluation of a behavioral neuroimaging task. *Psychiatry Res* 144: 123–130.
- Tandon R, Nasrallah HA, Keshavan MS (2009) Schizophrenia, "just the facts" 4. Clinical features and conceptualization. *Schizophr Res* 110: 1–23.
- Taylor SF, Kang J, Brege IS, Tso IF, Hosanagar A, et al. (2012) Meta-Analysis of Functional Neuroimaging Studies of Emotion Perception and Experience in Schizophrenia. *Biol Psychiatry* 71: 136–145.
- Grandjean D, Sander D, Pourtois G, Schwartz S, Seghier ML, et al. (2005) The voices of wrath: brain responses to angry prosody in meaningless speech. *Nat Neurosci* 8: 145–146.
- Sander D, Grandjean D, Pourtois G, Schwartz S, Seghier ML, et al. (2005) Emotion and attention interactions in social cognition: brain regions involved in processing anger prosody. *Neuroimage* 28: 848–858.
- Vuilleumier P, Armony JL, Driver J, Dolan RJ (2001) Effects of attention and emotion on face processing in the human brain: an event-related fMRI study. *Neuron* 30: 829–841.
- Carraro L, Castelli L, Macchiella C (2011) The automatic conservative: ideology-based attentional asymmetries in the processing of valenced information. *PLoS One* 6: e26456.
- DeLisi LE (2001) Speech disorder in schizophrenia: review of the literature and exploration of its relation to the uniquely human capacity for language. *Schizophr Bull* 27: 481–496.
- Mitchell RLC, Crow TJ (2005) Right hemisphere language functions and schizophrenia: the forgotten hemisphere? *Brain* 128: 963–978.
- Wang K, Cheung EF, Gong QY, Chan RC (2011) Semantic processing disturbance in patients with schizophrenia: a meta-analysis of the N400 component. *PLoS One* 6: e25435.
- Francis AN, Seidman LJ, Jabbar GA, Meshulam-Gately R, Thermenos HW, et al. (2012) Alterations in brain structures underlying language function in young adults at high familial risk for schizophrenia. *Schizophr Res* 141: 65–71.
- Iwashiro N, Suga M, Takano Y, Inoue H, Natsubori T, et al. (2012) Localized gray matter volume reductions in the pars triangularis of the inferior frontal



- gyrus in individuals at clinical high-risk for psychosis and first episode for schizophrenia. *Schizophr Res* 137: 124–131.
21. Takahashi T, Wood SJ, Yung AR, Soulsby B, McGorry PD, et al. (2009) Progressive Gray Matter Reduction of the Superior Temporal Gyrus During Transition to Psychosis. *Arch Gen Psychiatry* 66: 366–376.
  22. MacLeod CM, MacDonald PA (2000) Interdimensional interference in the Stroop effect: uncovering the cognitive and neural anatomy of attention. *Trends Cogn Sci* 4: 383–391.
  23. American Psychiatric Association (1994) *Diagnostic and Statistical Manual of Mental Disorders*, 4th ed. Washington DC: American Psychiatric Press.
  24. First MB, Spitzer RL, Gibbon M, Williams JBW (1997) Structured clinical interview for DSM-IV axis I disorders: clinical version (SCID-CV). Washington, DC: American Psychiatric Press.
  25. Fisher M, Holland C, Merzenich MM, Vinogradov S (2009) Using neuroplasticity-based auditory training to improve verbal memory in schizophrenia. *Am J Psychiatry* 166: 805–811.
  26. Kasai K, Shenton ME, Salisbury DF, Hirayasu Y, Lee CU, et al. (2003) Progressive decrease of left superior temporal gyrus gray matter volume in patients with first-episode schizophrenia. *Am J Psychiatry* 160: 156–164.
  27. Miyanishi T, Sumiyoshi T, Higuchi Y, Seo T, Suzuki M (2013) LORETA current source density for duration mismatch negativity and neuropsychological assessment in early schizophrenia. *PLoS One* 8: e61152.
  28. Kay SR, Fiszbein A, Opler LA (1987) The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull* 13: 261–276.
  29. Suga M, Yamasue H, Abe O, Yamasaki S, Yamada H, et al. (2010) Reduced gray matter volume of Brodmann's Area 45 is associated with severe psychotic symptoms in patients with schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 260: 465–473.
  30. Sumich A, Chitnis XA, Fannon DG, O'Ceallaigh S, Doku VC, et al. (2005) Unreality symptoms and volumetric measures of Heschl's gyrus and planum temporal in first-episode psychosis. *Biol Psychiatry* 57: 947–950.
  31. Yamasaki S, Yamasue H, Abe O, Yamada H, Iwanami A, et al. (2007) Reduced planum temporale volume and delusional behavior in patients with schizophrenia. *Eur Arch Psychiatry Clin Neurosci* 257: 318–324.
  32. Matsuoka K, Kim Y (2006) Japanese Adult Reading Test (JART). Shinkou-Igaku publishers, Tokyo.
  33. Matsuoka K, Uno M, Kasai K, Koyama K, Kim Y (2006) Estimation of premorbid IQ in individuals with Alzheimer's disease using Japanese ideographic script (Kanji) compound words: Japanese version of National Adult Reading Test. *Psychiatry and Clin Neurosci* 60: 332–339.
  34. Uetsuki M, Matsuoka K, Kim Y, Araki T, Suga M, et al. (2006) Estimation of premorbid IQ by JART in schizophrenia. *Seishin Igaku (Clinical Psychiatry)* 48: 15–22.
  35. Brysbaert M (1994) Lateral preferences and visual field asymmetries: appearances may have been overstated. *Cortex* 30: 413–429.
  36. Coren S (1993) The lateral preference inventory for measurement of handedness, footedness, eyedness, and earedness—norms for young-adults. *Bull Psychonomic Soci* 31: 1–3.
  37. Polemikos N, Palacliou C (2000) Sidedness preference as an index of organization of laterality. *Perceptual and Motor Skills* 91: 1083–1090.
  38. Hollingshead AB (1957) Two-factor index of social position. New Haven CT: Yale University Press.
  39. Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9: 97–113.
  40. First MB, Spitzer RL, Gibbon M, Williams JBW (1997) Structured clinical interview for DSM-IV axis I disorders, non-patient ed. New York: Biometrics Research Department, New York State Psychiatric Institute.
  41. Bradley MM, Lang PJ (1999) Affective norms for English words (ANEW): Instruction manual and affective ratings. The Center for Research in Psychophysiology, University of Florida.
  42. Ikehara S, Miyazaki M, Shirai S, Yokoo A, Nakaiwa H, et al. (1999) *Goi-Taikai - A Japanese Lexicon CDROM*. Tokyo: Iwanami Shoten.
  43. Ikegami A, Kaneda H, Sugisaki K, Suzuki T, Nakajima T, et al. (1998) *Daijisen*. In: Matsumura A, editor. Tokyo: Shogakukan.
  44. Gaillard R, Del Cul A, Naccache L, Vinckier F, Cohen L, et al. (2006) Nonconscious semantic processing of emotional words modulates conscious access. *Proc Natl Acad Sci U S A* 103: 7524–7529.
  45. Belin P, Zatorre RJ, Lafaille P, Ahad P, Pike B (2000) Voice-selective areas in human auditory cortex. *Nature* 403: 309–312.
  46. Aue T, Cuny C, Sander D, Grandjean D (2011) Peripheral responses to attended and unattended angry prosody: A dichotic listening paradigm. *Psychophysiology* 48: 385–392.
  47. Demily C, Attala N, Fouldrin G, Czernecki V, Menard JF, et al. (2010) The Emotional Stroop task: A comparison between schizophrenic subjects and controls. *European Psychiatry* 25: 75–79.
  48. Fear C, Sharp H, Healy D (1996) Cognitive processes in delusional disorders. *Br J Psychiatry* 168: 61–67.
  49. Kinderman P (1994) Attention bias, persecutory delusions and the self-concept. *Br J Med Psychol* 67: 53–66.
  50. Rao NP, Arasappa R, Reddy NN, Venkatasubramanian G, Reddy YC (2010) Emotional interference in obsessive-compulsive disorder: a neuropsychological study using optimized emotional Stroop test. *Psychiatry Res* 180: 99–104.
  51. Ullmann LP, Krasner L (1969) *A Psychological Approach to Abnormal Behaviour*. Englewood Cliffs: Prentice-Hall.
  52. Green MJ, Phillips ML (2004) Social threat perception and the evolution of paranoia. *Neurosci Biobehav Rev* 28: 333–342.
  53. Whalen PJ, Bush G, McNally RJ, Wilhelm S, McInerney SC, et al. (1998) The emotional counting Stroop paradigm: A functional magnetic resonance imaging probe of the anterior cingulate affective division. *Biol Psychiatry* 44: 1219–1228.
  54. Alba-Ferrara L, de Erausquin GA, Hirnstein M, Weis S, Hausmann M (2013) Emotional prosody modulates attention in schizophrenia patients with hallucinations. *Front Hum Neurosci* 7: 59.
  55. Hugdahl K, Westerhausen R, Alho K, Medvedev S, Laine M, et al. (2009) Attention and cognitive control: Unfolding the dichotic listening story. *Scand J Psychol* 50: 11–22.
  56. Wager TD, Phan KL, Liberzon I, Taylor SF (2003) Valence, gender, and lateralization of functional brain anatomy in emotion: a meta-analysis of findings from neuroimaging. *Neuroimage* 19: 513–531.
  57. Williams JM, Mathews A, MacLeod C (1996) The emotional Stroop task and psychopathology. *Psychol Bull* 120: 3–24.
  58. Hahn B, Robinson BM, Kaiser ST, Harvey AN, Beck VM, et al. (2010) Failure of schizophrenia patients to overcome salient distractors during working memory encoding. *Biol Psychiatry* 68: 603–609.
  59. Strauss GP, Llerena K, Gold JM (2011) Attentional disengagement from emotional stimuli in schizophrenia. *Schizophr Res* 131: 219–223.
  60. Park SH, Kim JJ, Kim CH, Kim JH, Lee KH (2011) Sustained attention in the context of emotional processing in patients with schizophrenia. *Psychiatry Res* 187: 18–23.
  61. Bach DR, Buxtorf K, Grandjean D, Strik WK (2009) The influence of emotion clarity on emotional prosody identification in paranoid schizophrenia. *Psychol Med* 39: 927–938.
  62. Leitman DI, Hoptman MJ, Foxe JJ, Saccette E, Wylie GR, et al. (2007) The neural substrates of impaired prosodic detection in schizophrenia and its sensorial antecedents. *Am J Psychiatry* 164: 474–482.
  63. Li X, Branch CA, DeLisi LE (2009) Language pathway abnormalities in schizophrenia: a review of fMRI and other imaging studies. *Curr Opin Psychiatry* 22: 131–139.
  64. Buchanan TW (2007) Retrieval of emotional memories. *Psychol Bull* 133: 761–779.
  65. Sass K, Habel U, Kellermann T, Mathiak K, Gauggel S, et al. (2012) The influence of positive and negative emotional associations on semantic processing in depression: An fMRI study. *Hum Brain Mapp*.
  66. Nishitani N, Schurmann M, Amunts K, Hari R (2005) Broca's region: from action to language. *Physiology (Bethesda)* 20: 60–69.
  67. Dollfus S, Razafimandimby A, Maiza O, Lebain P, Brazo P, et al. (2008) Functional deficit in the medial prefrontal cortex during a language comprehension task in patients with schizophrenia. *Schizophr Res* 99: 304–311.
  68. Koeda M, Takahashi H, Yahata N, Matsuura M, Asai K, et al. (2006) Language processing and human voice perception in schizophrenia: a functional magnetic resonance imaging study. *Biol Psychiatry* 59: 948–957.
  69. Ngan ET, Vouloomanos A, Cairo TA, Laurens KR, Bates AT, et al. (2003) Abnormal processing of speech during oddball target detection in schizophrenia. *Neuroimage* 20: 889–897.
  70. Angrilli A, Spironelli C, Elbert T, Crow TJ, Marano G, et al. (2009) Schizophrenia as failure of left hemispheric dominance for the phonological component of language. *PLoS One* 4: e4507.
  71. Braff DL, Geyer MA (1990) Sensorimotor gating and schizophrenia. Human and animal model studies. *Arch Gen Psychiatry* 47: 181–188.
  72. McGhie A, Chapman J (1961) Disorders of attention and perception in early schizophrenia. *Br J Med Psychol* 34: 103–116.
  73. Smith DM, Grant B, Fisher DJ, Borraresi G, Labelle A, et al. (2013) Auditory verbal hallucinations in schizophrenia correlate with P50 gating. *Clin Neurophysiol* 124: 1329–1335.
  74. Rajji TK, Mulsant BH (2008) Nature and course of cognitive function in late-life schizophrenia: A systematic review. *Schizophr Res* 102: 122–140.
  75. Moncrieff J, Leo J (2010) A systematic review of the effects of antipsychotic drugs on brain volume. *Psychol Med* 40: 1409–1422.