

# Evaluation of Dissolved Organic Carbon as a Soil Quality Indicator in National Monitoring Schemes

David L. Jones<sup>1\*</sup>, Paul Simfukwe<sup>1</sup>, Paul W. Hill<sup>1</sup>, Robert T. E. Mills<sup>2</sup>, Bridget A. Emmett<sup>3</sup>

**1** School of the Environment, Natural Resources & Geography, Bangor University, Bangor, Gwynedd, United Kingdom, **2** Ecole Polytechnique Fédérale de Lausanne (EPFL), School of Architecture, Civil and Environmental Engineering (ENAC), Laboratory of Ecological Systems (ECOS), Lausanne, Switzerland, **3** Centre for Ecology and Hydrology, Environment Centre Wales, Bangor, Gwynedd, United Kingdom

## Abstract

**Background:** Monitoring the properties of dissolved organic carbon (DOC) in soil water is frequently used to evaluate changes in soil quality and to explain shifts in freshwater ecosystem functioning.

**Methods:** Using >700 individual soils (0–15 cm) collected from a 209,331 km<sup>2</sup> area we evaluated the relationship between soil classification (7 major soil types) or vegetation cover (8 dominant classes, e.g. cropland, grassland, forest) and the absorbance properties (254 and 400 nm), DOC quantity and quality (SUVA, total soluble phenolics) of soil water.

**Results:** Overall, a good correlation ( $r^2=0.58$ ) was apparent between soil water absorbance and DOC concentration across the diverse range of soil types tested. In contrast, both DOC and the absorbance properties of soil water provided a poor predictor of SUVA or soluble phenolics which we used as a measure of humic substance concentration. Significant overlap in the measured ranges for UV absorbance, DOC, phenolic content and especially SUVA of soil water were apparent between the 8 vegetation and 7 soil classes. A number of significant differences, however, were apparent within these populations with total soluble phenolics giving the greatest statistical separation between both soil and vegetation groups.

**Conclusions:** We conclude that the quality of DOC rather than its quantity provides a more useful measure of soil quality in large scale surveys.

**Citation:** Jones DL, Simfukwe P, Hill PW, Mills RTE, Emmett BA (2014) Evaluation of Dissolved Organic Carbon as a Soil Quality Indicator in National Monitoring Schemes. PLoS ONE 9(3): e90882. doi:10.1371/journal.pone.0090882

**Editor:** Dafeng Hui, Tennessee State University, United States of America

**Received:** November 14, 2013; **Accepted:** February 3, 2014; **Published:** March 14, 2014

**Copyright:** © 2014 Jones et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was funded by the United Kingdom Department for Environment, Food and Rural Affairs. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: d.jones@bangor.ac.uk

## Introduction

Measurement of dissolved organic matter (DOM) concentrations have provided a sensitive indicator for assessing large scale geographical and temporal (e.g. decadal) changes in water quality [1]. In contrast, while DOM is frequently measured in soil studies, it has rarely been used for evaluating changes in soil quality in large scale (e.g. national) monitoring programmes [2]. To some extent, the DOM concentration of freshwaters should reflect that of the surrounding catchment area suggesting that an assessment of soil DOM might provide a sensitive soil quality indicator. This would also match with studies investigating the progressive increase in DOM within rivers, a widespread trend observed in both Europe and North America [3–4]. An investigation of the spatial or temporal changes in soil DOM within different soil types could therefore prove useful for explaining these trends in catchments containing a complex mixture of soils. One potential drawback, however, is that determination of DOM in soil can be very time consuming and expensive when large numbers of samples need to be processed. One potential solution is to use the spectral properties of soil water which provides a rapid proxy for estimating dissolved organic C (DOC) concentration [5–10]. For a single geographical location, strong relationships are often

apparent between DOC concentration and UV absorbance [11–13]. However, these relationships are often non-linear and the mathematical function describing the relationship can vary with season or changes in hydrological flow patterns [14–16]. Further, whilst differences in the quality and quantity of DOC are frequently reported for contrasting soil and vegetation types, whether these dissimilarities remain valid over large geographical scales remains uncertain. In this context, we undertook a national large scale survey of the quality and quantity of DOM across Great Britain to evaluate its potential for evaluating changes to soil quality in national soil quality monitoring programmes. Our first aim was to critically evaluate the relationship between direct and indirect methods for DOC quantification. Secondly, we aimed to evaluate whether different soil classes and vegetation cover types possessed unique DOM signatures in terms of both quantity and quality and whether these might be useful for explaining changes in DOM occurring at a national scale.

## Methods

### Soil Sampling

To encompass all the major soil and land use types, a total of 702 soil samples were collected throughout Great Britain in June–

July, 2007 (area 209,331 km<sup>2</sup>; ca. 300 km<sup>2</sup> sample<sup>-1</sup>) as part of the Centre for Ecology and Hydrology Countryside Survey (CS) [17]. Samples were selected, based on a stratified random sample of 1 km squares at gridpoints on a 15 km grid using the Institute of Terrestrial Ecology (ITE) Land Classification as the basis of the stratification (Fig. S1). At each grid intersection, a 1 km<sup>2</sup> sample area was selected. Within the 1 km<sup>2</sup> sample area, a 5 × 5 m<sup>2</sup> plot was randomly located and replicate 15 cm long × 4 cm diameter soil cores were collected. Topsoils (0–15 cm) were only sampled to reflect past and current (1978–present) standard practice in UK national monitoring [18–20]. This depth was originally selected by the UK Government’s Department for Environment, Food and Rural Affairs as it was hypothesized that it would show the greatest change in response to environmental perturbation (including changes in land use, climate, agronomic management or atmospheric deposition). The soil horizons sampled included H, O and A horizons with *a*, *e*, *i*, *h*, *g*, *k* and *p* sub-designations [21]. Across all land use categories, the dominant eight soil groupings (% of total) were: Brown soils (31%), Podzolic soils (15%), Surface water (SW) gley soils (18%), Peat soils (13%), Groundwater (GW) gley soils (12%), Lithomorphous soils (8%), and Pelosol soils (3%) [22]. The FAO World Reference Base Classification equivalent categories for these soil groups are presented in Table S1 and their major chemical, physical and biological properties presented in Table 1. Vegetation cover at each sampling point was classified into eight aggregated vegetation classes (AVC) with the following groupings (% of total): Cropland (15%), Tall grass and herbs (4%), Fertile grassland (19%), Infertile grassland (21%), Lowland woodland (3%), Upland woodland (8%), Moorland grassland mosaic (11%), and Heathland and bog (19%) [17]. Aggregate vegetation classes were derived by cluster analysis of the mean DECORANA scores for 100 smaller classes obtained by TWINSpan analysis of plant species data in each sample plots [37]. Further descriptions of the vegetation types can be found in Table S2.

### Soil Water Collection and Analysis

Soil water was obtained by adding artificial rainwater to the intact soil columns and collecting the leachate as detailed in [23–24]. The absorbance of the soil leachate water was measured at 254 and 400 nm on a Synergy 96 well plate spectrophotometer (BioTek Instruments Inc., Winooski, VT) using Falcon flat-bottom UV well plates (BD Biosciences, Franklin Lakes, NJ). DOC concentrations were measured with a TOC-V analyser (Shimadzu Corp., Kyoto, Japan). Total dissolved phenolics and tannins were assayed colorimetrically using the Folin-Ciocalteu reagent (F9252; Sigma-Aldrich Inc.) according to [25] using gallic acid as a standard. Specific ultraviolet absorbance (SUVA) was calculated by dividing the absorbance at 254 nm (cm<sup>-1</sup>) by the DOC concentration (mg l<sup>-1</sup>).


### Background Soil Analysis

Background soil characteristics were measured on replicate cores matching those used in the analysis above. After collection from the field, the soil was extruded from the core, roots and stones removed and the soil homogenised and dried (105 °C). Total soil C and N were analysed on an Elementar Vario-EL elemental analyser (Elementar Analysensysteme GmbH, Hanau, Germany) using the UKAS accredited method SOP3102 [23]. Bulk density was calculated as mass/volume after the removal of stones (>2 mm) and accounting for their volume [23]. Soil organic matter was determined by loss-on-ignition (LOI) by first drying soil (10 g) at 105°C and then measuring the mass loss after further heating at 375 °C for 16 h. Available soil P was measured using

**Table 1. Properties of the major soil groups.**

Soil group	Soil pH	Soil organic matter (%)	C-to-N ratio	Olsen P (mg kg <sup>-1</sup> )	Bulk density (g cm <sup>-3</sup> )	Exchangeable Ca (mmol kg <sup>-1</sup> )	Exchangeable Al (mmol kg <sup>-1</sup> )	Soil respiration (mg C cm <sup>-3</sup> h <sup>-1</sup> )	Soil respiration (g C g SOC <sup>-1</sup> h <sup>-1</sup> )
Pelosol	6.8 ± 1.4 <sup>a</sup>	7.3 ± 3.3 <sup>cd</sup>	10.7 ± 0.8 <sup>de</sup>	28 ± 22 <sup>abc</sup>	1.10 ± 0.21 <sup>a</sup>	62 ± 40 <sup>a</sup>	1.3 ± 4.8 <sup>ab</sup>	10.2 ± 26.4 <sup>bc</sup>	4.7 ± 3.4 <sup>b</sup>
Brown	6.5 ± 1.2 <sup>a</sup>	10.0 ± 12.5 <sup>d</sup>	12.2 ± 4.1 <sup>e</sup>	32 ± 27 <sup>abc</sup>	1.04 ± 0.33 <sup>a</sup>	39 ± 27 <sup>a</sup>	2.1 ± 4.8 <sup>b</sup>	8.6 ± 11.0 <sup>c</sup>	7.3 ± 5.6 <sup>a</sup>
Groundwater gley	6.6 ± 1.2 <sup>a</sup>	10.9 ± 13.5 <sup>d</sup>	13.4 ± 4.4 <sup>cde</sup>	32 ± 34 <sup>abc</sup>	0.98 ± 0.36 <sup>a</sup>	42 ± 31 <sup>a</sup>	2.3 ± 5.4 <sup>ab</sup>	13.2 ± 24.1 <sup>bc</sup>	8.1 ± 6.4 <sup>a</sup>
Surface water gley	5.9 ± 1.3 <sup>a</sup>	24.9 ± 30.0 <sup>bc</sup>	14.2 ± 5.5 <sup>cd</sup>	25 ± 28 <sup>bc</sup>	0.78 ± 0.43 <sup>b</sup>	54 ± 49 <sup>a</sup>	6.3 ± 14.6 <sup>ab</sup>	14.9 ± 19.2 <sup>bc</sup>	6.6 ± 5.0 <sup>ab</sup>
Podzolic	4.9 ± 0.9 <sup>b</sup>	28.8 ± 27.8 <sup>b</sup>	17.9 ± 7.0 <sup>b</sup>	21 ± 20 <sup>c</sup>	0.57 ± 0.33 <sup>c</sup>	20 ± 29 <sup>b</sup>	7.1 ± 8.9 <sup>ab</sup>	17.9 ± 32.9 <sup>bc</sup>	5.1 ± 3.9 <sup>b</sup>
Lithomorphous	6.0 ± 1.4 <sup>a</sup>	35.1 ± 32.8 <sup>b</sup>	16.2 ± 6.4 <sup>bc</sup>	41 ± 69 <sup>ab</sup>	0.56 ± 0.38 <sup>c</sup>	42 ± 39 <sup>a</sup>	3.4 ± 5.6 <sup>ab</sup>	20.8 ± 33.0 <sup>b</sup>	6.4 ± 5.6 <sup>ab</sup>
Peat	4.6 ± 0.6 <sup>b</sup>	75.9 ± 27.9 <sup>a</sup>	24.1 ± 8.5 <sup>a</sup>	44 ± 51 <sup>a</sup>	0.20 ± 0.21 <sup>d</sup>	46 ± 50 <sup>a</sup>	9.6 ± 38.0 <sup>a</sup>	43.1 ± 37.1 <sup>a</sup>	4.5 ± 3.8 <sup>b</sup>
ANOVA	***	***	***	**	***	***	**	***	***

Values are expressed on a dry weight basis and represent mean ± SD. \*\* and \*\*\* indicate significant differences between soil groups at the P<0.01 and P<0.001 levels respectively. Superscript letters indicate significant between soil groups at the P<0.05 level. doi:10.1371/journal.pone.0090882.t001



**Figure 1. Relationship between the absorbance of soil water**

**from 702 individual soils measured in the UV (254 nm) range and DOC concentration (Panel A), the relationship between total soluble phenolics and either solution absorbance at 400 nm (Panel B) or DOC concentration (Panel C).** The lines are linear regression for all the data in the plots (Panel A,  $r^2 = 0.579$ ;  $y = 0.387 + 0.301x$ ; Panel B,  $r^2 = 0.432$ ;  $y = 0.12 + 0.139x$ ; Panel C,  $r^2 = 0.375$ ;  $y = 0.025 + 0.068x$ ).  
doi:10.1371/journal.pone.0090882.g001

the Olsen method whereby 5 g of soil was extracted with 100 ml of 0.5 M sodium bicarbonate (pH 8.5). The P in the extract was determined colorimetrically using the molybdate blue method (880 nm) using a continuous flow analyser [23]. Soil pH was measured by equilibrating 10 g of field-moist soil with 25 ml of deionised water. Exchangeable Ca and Al were determined by shaking 5 g of soil with 25 ml of 1.0 M  $\text{NH}_4\text{Cl}$  (250 rev  $\text{min}^{-1}$ , 60 min). Subsequently, the extracts were centrifuged (5000 g, 10 min) and the supernatant recovered for analysis. Ca in the extracts was determined by atomic absorption spectrometry on a Perkin Elmer Analyst 400 Atomic Absorption Spectrometer (PerkinElmer, Waltham, MA, USA). The extracts were diluted with  $\text{LaCl}_3$  (0.5% w/v) prior to Ca determination. Al concentration in the extracts was determined using the modified catechol violet method [35]. The absorbance of the solution was measured at 580 nm using a PowerWave XS scanning microplate spectrophotometer (BioTek Instrument, Winooski, VT). Basal soil respiration was determined on one replicate core. The cores were wet to field capacity as described previously, placed in a sealed chamber (1250  $\text{cm}^3$  head space). The soils were then incubated at 10°C (average UK air temperature) for 1 h (at which linearity was known to be established following testing on selected cores which covered the range of soil types sampled). Subsequently, the head space gas was analysed for  $\text{CO}_2$  concentration using a Clarus 500 Gas Chromatograph (PerkinElmer). SR was determined as the change in  $\text{CO}_2$  concentration before and after incubation corrected for soil dry weight and soil organic matter content.


### Statistical Analysis

Linear and stepwise regression analysis was undertaken using Minitab v16 (Minitab Inc., State College, PA). When the solution DOC, absorbance, soluble phenolic and SUVA values were grouped according to soil and vegetation type the data failed normality testing (Shapiro-Wilk). Consequently, the data were log transformed, normality verified and an ANOVA performed with Tukey-pairwise comparisons ( $P < 0.05$  cut-off) using Minitab v16.

## Results

### Soil Water Properties

The relationship between the absorbance of soil water in the UV (254 nm) and visible (400 nm) range across 702 individual sites showed a strong linear correlation ( $r^2 = 0.931$ ;  $P < 0.001$ ; Fig. S2). A strong positive correlation was also observed between the DOC concentration in soil water and both absorbance at 254 nm ( $r^2 = 0.58$ ;  $P < 0.001$ ; Fig. 1) and 400 nm ( $r^2 = 0.47$ ;  $P < 0.001$ ). The prediction of DOC concentration using both absorbance values in a stepwise regression model did not result in a significantly better fit ( $r^2 = 0.61$ ). In contrast, the absorbance characteristics of soil water proved to be a less reliable predictor of total soluble phenolics (254 nm  $r^2 = 0.41$ ;  $P < 0.001$ ; 400 nm  $r^2 = 0.43$ ;  $P < 0.001$ ; Fig. 1). Similarly, DOC concentration either with or without inclusion of spectral properties into the regression model, proved to be the least reliable predictor of total soluble phenolics ( $r^2 = 0.38$ ;  $P < 0.001$ ). SUVA proved to be the worst predictor of



**Figure 2. Box plots showing the influence of soil class (panels A–C) and vegetation cover type (panels D–F) on the absorbance of soil solutions at 254 nm, their DOC and total soluble phenolic concentration from 702 individual soils sampled as part of a nationwide soil quality assessment.** The boundary of the box closest to zero indicates the 25<sup>th</sup> percentile, the line within the box marks the median, and the boundary of the box farthest from zero indicates the 75<sup>th</sup> percentile. Whiskers (error bars) above and below the box indicate the 90<sup>th</sup> and 10<sup>th</sup> percentiles respectively. Different letters indicate significant difference between individual groups ( $P < 0.05$ ). Soils are ranked in organic matter (OM) content from left (low OM) to right (high OM) while vegetation is broadly ranked according to agricultural productivity. doi:10.1371/journal.pone.0090882.g002

total soluble phenolics with either linear ( $r^2 = 0.13$ ) or non-linear models (data not presented).

### Is Soil or Vegetation a Better Predictor of DOC Quantity and Quality?

The quantity and quality of DOC as a function of different soil class and vegetation types is shown in Figure 2. Overall, there was

a large similarity in the measured ranges of DOC, absorbance at 254 nm and soluble phenolics between the different groupings for both soil and vegetation. Despite this, ANOVA revealed significant differences between groups (all  $P < 0.001$ ) with response gradients apparent when the groups were ranked according to either organic matter content (soils) or an intensification/altitude scale (vegetation) (Fig. 2). Of the parameters measured, soluble

phenolic content gave the greatest separation between groups. In contrast, SUVA revealed no significant differences between groupings for both soil ( $P=0.483$ ) and vegetation ( $P=0.819$ ; Fig. S3).

## Discussion

### Method of Soil Sampling

The sampling undertaken here formed part of a nationwide monitoring programme which is carried out each decade to assess changes in soil and water quality, landscape features and aquatic and terrestrial biodiversity [19,26]. Whilst the sampling depth (0–15 cm) is highly suited to agricultural soils which show a high degree of vertical homogeneity (due to cultivation), we acknowledge that the sampling regime may be less suited to highly stratified soils [27]. In these horizonated soils, differential amounts of O, E, A and possibly B horizons could be sampled by coring from 0–15 cm, depending on their thickness. This has led to many studies, particularly in forests, where soil quality is measured on organic and mineral horizons separately [27–28]. In our case, the historical legacy is such that the sampling protocols will not be changed for the foreseeable future (on scientific, socio-political and economic grounds) and therefore our results should be viewed in the context of this.

### DOC and Soil Quality Assessment

Overall, our results show that in large geographical scale soil quality assessments across a broad range of vegetation and soil types, soil water DOC concentrations can be rapidly and cheaply estimated from their spectral properties. This is particularly relevant as DOC in itself can be used to predict the movement of both organic and inorganic pollutants in soil and is often seen as a pollutant itself when entering freshwaters [9,29–30]. Disappointingly, however, we found that all soil and vegetation types had a wide variability in their DOC concentrations. This indicates that the quantity of DOC may not represent a sensitive indicator for monitoring the stability of ecosystems when faced with anthropogenic perturbation (e.g. land use change). This is in contrast to [24] who found different rates of N mineralization ( $\text{NH}_4^+ + \text{NO}_3^-$ ) in the same soils alongside other soil quality indicators as shown in Table 1. While SUVA is frequently used for assessing the quality of DOC in freshwaters [30], our results indicated that it was incapable of separating between land uses and soil classes over a wide geographical range and is therefore probably unsuited to large scale soil quality assessments. In contrast, total soluble phenols gave the best separation between ecosystem types. This was a surprising result as we had assumed that SUVA and soluble phenols would be highly correlated. This disagreement could be due to interference in both the determination of total phenolics (e.g.  $\text{SO}_2$ , DOC; [31–32]) or SUVA (e.g. pH,  $\text{Fe}^{3+}$ ; [12]). However, more likely it is due to the natural variability in the total phenol content of humic substances originating from different soil classes and vegetation cover types [33–34]. As expected, the quality and quantity of DOC were dependent on both vegetation and soil type. This is to some extent expected considering that certain soil types favour certain vegetation covers, however, it does indicate that 2-way stratification by soil and vegetation type may prove more useful for revealing unique DOC signatures.

With respect to the measurement of DOC quality, we acknowledge that recent methodological advancements are increasing our potential to characterise the many thousands of compounds that comprise DOC [36]. It is highly likely that these new analytical approaches will reveal compounds that are specific

to different functional soil or plant types (e.g. keystone compounds) improving our capacity to use DOC as a soil quality indicator. In addition, this analysis may also directly support the interpretation of temporal changes in other soil quality indicators (e.g. soil organic matter quality and quantity, soil biodiversity).

## Supporting Information

**Figure S1 Map of the UK showing the individual soil sampling locations used in the study.** The total land area is 209,331 km<sup>2</sup>. (TIF)

**Figure S2 Relationship between the absorbance of soil water from 702 individual soils measured in either the UV (254 nm) or visible (400 nm) range.** The line is a linear regression for all the data in the plot ( $r^2=0.931$ ;  $y = -0.01+0.123x$ ). (TIF)

**Figure S3 Box plots showing the influence of soil type (panel A) and vegetation cover (panel B) on the specific UV absorbance (SUVA) values from 702 individual soils sampled as part of a nationwide soil quality assessment.** The boundary of the box closest to zero indicates the 25<sup>th</sup> percentile, the line within the box marks the median, and the boundary of the box farthest from zero indicates the 75<sup>th</sup> percentile. Whiskers (error bars) above and below the box indicate the 90<sup>th</sup> and 10<sup>th</sup> percentiles respectively. No significant differences were apparent between treatments. (TIF)

**Table S1 Comparable classification of the UK soil groups with those in the FAO World Reference Base Classification (WRB, 2006).** (DOCX)

**Table S2 Summary descriptions of the eight aggregate vegetation classes represented. Table adapted from Smart et al. (2003).** (DOCX)

## Acknowledgments

We would like to thank Ed Rowe at the Centre for Ecology and Hydrology for help with the experiments and data handling. Countryside Survey (CS) is funded by a consortium of UK government funded bodies led by the Natural Environment Research Council (NERC) and the Department for Environment, Food and Rural Affairs (Defra), which includes the Centre for Ecology and Hydrology, Countryside Council for Wales, Forestry Commission, Natural England, the Northern Ireland Environment Agency, the Scottish Government, Scottish Natural Heritage, and the Welsh Assembly Government. The completion of CS was only possible with the support and advice of many dedicated individuals from these and other organizations. The authors would like to thank the land owners, farmers, and other land managers who gave permission for the field surveyors to collect data and samples from their land; without such cooperation scientific field studies like CS would not be possible. Additionally, the authors would especially like to acknowledge the major contribution made by the team of laboratory staff who processed all the soils and undertook the chemical analyses.

## Author Contributions

Conceived and designed the experiments: DJJ BAE. Performed the experiments: PS RTEM. Analyzed the data: PS PWH DJJ RTEM. Wrote the paper: DJJ PWH PS BAE.

## References

- Camarero L, Rogora M, Mosello T, Anderson NJ, Barbieri A, et al. (2009) Regionalisation of chemical variability in European mountain lakes. *Freshwater Biol* 54: 2452–2469.
- Filep T, Rekası M, Kincses I (2008) Effect of soil quality on dissolved organic matter (DOM) content of soils. *Cereal Res Commun* 36: 667–670.
- Evans CD, Monteith DT, Cooper DM (2005) Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environ Pollut* 137: 55–7.
- Clutterbuck B, Yallop AR (2010) Land management as a factor controlling dissolved organic carbon release from upland peat soils. 2 Changes in DOC productivity over four decades. *Sci Total Environ* 408: 6179–6191.
- Mattson JS, Smith CA, Jones TT, Gerchako SM, Epstein BD (1974) Continuous monitoring of dissolved organic-matter by UV-visible photometry. *Limnol Oceanogr* 19: 530–535.
- Dittman JA, Shanley JB, Driscoll CT, Aiken GR, Chalmers AT, et al. (2009) Ultraviolet absorbance as a proxy for total dissolved mercury in streams. *Environ Pollut* 157: 1953–1956.
- Brandstetter A, Sletten RS, Mentler A, Wenzel WW (1996) Estimating dissolved organic carbon in natural waters by UV absorbance (254 nm). *J Soil Sci Plant Nutr* 159: 605–607.
- Jaffrain J, Gerard F, Meyer M, Ranger J (2007) Assessing the quality of dissolved organic matter in forest soils using ultraviolet absorption spectrophotometry. *Soil Sci Soc Am J* 71: 1851–1858.
- Amery F, Degryse F, Cheyns K, De Troyer I, Mertens J, et al. (2008) The UV-absorbance of dissolved organic matter predicts the fivefold variation in its affinity for mobilizing Cu in an agricultural soil horizon. *Eur J Soil Sci* 59: 1087–1095.
- Hansson K, Kleja DB, Kalbitz K, Larsson H (2010) Amounts of carbon mineralised and leached as DOC during decomposition of Norway spruce needles and fine roots. *Soil Biol Biochem* 42: 178–185.
- Dilling J, Kaiser K (2002) Estimation of the hydrophobic fraction of dissolved organic matter in water samples using UV photometry. *Water Res* 36: 5037–5044.
- Weishaar JL, Aiken GR, Bergamaschi BA, Fram MS, Fujii R, et al. (2003) Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Environ Sci Technol* 37: 4702–4708.
- Tipping E, Corbishley HT, Koprivnjak JF, Lapworth DJ, Miller MP, et al. (2009) Quantification of natural DOM from UV absorption at two wavelengths. *Environ Chem* 6: 472–476.
- Wheeler JR (1977) Dissolved organic-carbon - spectral relationships in coastal waters. *Limnol Oceanogr* 22: 573–575.
- Edwards AC, Cresser MS (1987) Relationships between ultraviolet absorbency and total organic-carbon in 2 upland catchments. *Water Res* 21: 49–56.
- Izbicki JA, Pimentel IM, Johnson R, Aiken GR, Leenheer J (2007) Concentration, UV-spectroscopic characteristics and fractionation of DOC in stormflow from an urban stream, Southern California, USA. *Environ Chem* 4: 35–48.
- Emmett BA, Reynolds B, Chamberlain PM, Rowe E, Spurgeon D, et al. (2010) Countryside Survey: Soils Report from 2007. CS Technical Report No. 9/07, Centre for Ecology and Hydrology, Wallingford, UK.
- Bellamy PH, Loveland PJ, Bradley RI, Lark RM, Kirk GJD (2005) Carbon losses from all soils across England and Wales 1978–2003. *Nature* 437: 245–248.
- Norton LR, Maskell LC, Smart SS, Dunbar MJ, Emmett BA, et al. (2012) Measuring stock and change in the GB countryside for policy - Key findings and developments from the Countryside Survey 2007 field survey. *J Environ Manage* 113: 117–127.
- Reynolds B, Chamberlain PM, Poskitt J, Woods C, Scott WA, et al. (2013) Countryside Survey: National "soil change" 1978–2007 for topsoils in Great Britain—acidity, carbon, and total nitrogen status. *Vadose Zone J* 12: vjz2012.0114
- FAO (2006) Guidelines for Soil Description. Fourth Edition. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Avery BW (1990) Soils of the British Isles. Cambridge University Press, Cambridge.
- Emmett BA, Frogbrook ZL, Chamberlain PM, Griffiths R, Pickup R, et al. (2008) Countryside Survey. Soils Manual. NERC/Centre for Ecology & Hydrology, (CS Technical Report No. 3/07, CEH Project Number: C03259), 180 pp.
- Rowe EC, Emmett BA, Frogbrook ZL, Robinson DA, Hughes S (2012) Nitrogen deposition and climate effects on soil nitrogen availability: Influences of habitat type and vegetation. *Sci Total Environ* 434: 62–70.
- Velioglu YS, Mazza G, Gao L, Oomah BD (1998) Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *J Agric Food Chem* 46: 4113–4117.
- Smart SM, Clarke RT, van de Poll HM, Robertson EJ, Shield ER, et al. (2003) National-scale vegetation change across Britain: an analysis of sample-based surveillance data from the Countryside Surveys of 1990 and 1998. *J Environ Manage* 67: 239–254.
- Lawrence GB, Fernandez IJ, Richter DD, Ross DS, Hazlett PW, et al. (2013) Measuring environmental change in forest ecosystems by repeated soil sampling: a North American perspective. *J Environ Qual* 42: 623–639.
- O'Neill KP, Amacher MC, Palmer CJ (2005) Developing a national indicator of soil quality on U.S. forestlands: Methods and initial results. *Environ Monit Assess* 107: 59–80.
- Steinberg CEW, Hoss S, Kloas W, Lutz I, Meinelt T, et al. (2003) Hormonelike effects of humic substances on fish, amphibians, and invertebrates. *Environ Toxicol* 19: 409–411.
- Zhang Q, Kuang WF, Liu LY, Li KX, Wong KH, et al. (2009) Trihalomethane, haloacetonitrile, and chloral hydrate formation potentials of organic carbon fractions from sub-tropical forest soils. *J Hazard Mater* 172: 880–887.
- Matilainen A, Gjessing ET, Lahtinen T, Hed L, Bhatnagar A, et al. (2011) An overview of the methods used in the characterisation of natural organic matter (NOM) in relation to drinking water treatment. *Chemosphere* 83: 1431–1442.
- Ohno T, First PR (1998) Assessment of the Folin and Ciocalteu's method for determining soil phenolic carbon. *J Environ Qual* 27: 776–782.
- Sharma OK, Krishnan PS (1966). Colorimetric estimation of humic acid with phenol reagent of Folin and Ciocalteu. *Anal Biochem* 14: 11–17.
- Thorn KA, Folan DW, MacCarthy P (1989) Characterization of the International Humic Substances Society standard and reference fulvic and humic acids by solution state carbon-13 (<sup>13</sup>C) and hydrogen-1 (<sup>1</sup>H) nuclear magnetic resonance spectrometry. U.S. Geological Survey, Water-Resources Investigations Report 89-4196, Denver, CO, USA.
- Dougan WK, Wilson AL (1974) The absorptiometric determination of aluminium in water. A comparison of some chromogenic reagents and the development of an improved method. *Analyst* 99: 413–30.
- Osborne DM, Podgorski DC, Bronk DA, Roberts Q, Sipler RE, et al. (2013) Molecular-level characterization of reactive and refractory dissolved natural organic nitrogen compounds by atmospheric pressure photoionization coupled to Fourier transform ion cyclotron resonance mass spectrometry. *Rapid Comm. Mass Spectrom.* 27: 851–858.
- Bunce RGH, Barr CJ, Gillespie MK, Howard DC, Scott RA, et al. (1999) Vegetation of the British Countryside—the Countryside Vegetation System. Department of the Environment, London.