



Anti-Cyclic Citrullinated Peptide Antibody Is Associated with Interstitial Lung Disease in Patients with Rheumatoid Arthritis

Yufeng Yin¹✉, Di Liang¹✉, Lidan Zhao¹, Yang Li¹, Wei Liu², Yan Ren¹, Yongzhe Li¹, Xiaofeng Zeng¹, Fengchun Zhang¹, Fulin Tang¹, Guangliang Shan³, Xuan Zhang^{1*}

1 Department of Rheumatology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China,

2 Department of Radiology, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China,

3 Department of Epidemiology, Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

Abstract

Objective: Patients with rheumatoid arthritis (RA) are at risk to develop RA-associated interstitial lung disease (RA-ILD). This retrospective study aimed to investigate the potential association of the positivity of serum anti-cyclic citrullinated peptide antibody (anti-CCP2) and rheumatoid factor (RF) with RA-ILD in RA patients.

Methods: A total of 285 RA patients were recruited at the inpatient service of Peking Union Medical College Hospital in China between 2004 and 2013. Individual patients were evaluated for the evidence of ILD. The concentrations of serum anti-CCP2 and RF in individual patients were measured. The potential risk factors for ILD in RA patients were assessed by univariate and multivariate models.

Results: There were 71 RA patients with RA-ILD, accounting for 24.9% in this population. The positive rates of anti-CCP2 and RF in the patients with RA-ILD were significantly higher than that in the patients with RA-only (88.7% vs. 67.3%, $p < 0.001$; 84.5% vs. 70.6%, $p = 0.02$, respectively). Univariate and multivariate logistic regression analysis revealed that RA patients with positive serum anti-CCP2, but not RF, were associated with an increased risk of ILD (crude odds ratio [cOR] 3.83, 95% confidence interval [CI] 1.74–8.43, $p < 0.001$; adjusted odds ratio [aOR] 3.50, 95% CI 1.52–8.04, $p < 0.001$).

Conclusion: Our findings suggest that positive serum anti-CCP2, but not RF, may be associated with RA-ILD in RA patients.

Citation: Yin Y, Liang D, Zhao L, Li Y, Liu W, et al. (2014) Anti-Cyclic Citrullinated Peptide Antibody Is Associated with Interstitial Lung Disease in Patients with Rheumatoid Arthritis. PLoS ONE 9(4): e92449. doi:10.1371/journal.pone.0092449

Editor: Song Guo Zheng, Penn State University, United States of America

Received: November 22, 2013; **Accepted:** February 21, 2014; **Published:** April 17, 2014

Copyright: © 2014 Yin et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study is supported by the grants of the National Natural Science Foundation of China (30972731, 81172859, 81273312, 81302594), the National High Technology Research and Development Project of China (2011AA020111, 2012AA02A513), the National Major Scientific and Technological Special Project (2012ZX09303006-002), the Research Special Fund for Public Welfare Industry of Health (201202004), the Capital Health Research and Development of Special (2011-4001-02), and the National laboratory Special Fund (2060204). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: zxpumch2003@sina.com

✉ These authors contributed equally to this work.

Introduction

Patients with rheumatoid arthritis (RA) display high levels of autoantibodies as well as extra-articular manifestations, such as interstitial lung disease (ILD) [1–3]. The RA-related interstitial lung disease (RA-ILD) occurs in nearly 7–10% of the RA patients, and often is associated with a poor prognosis [4,5]. Therefore, the discovery of risk factors contributing to the development of ILD will be of great significance in the prevention and intervention of patients with RA-ILD.

Autoantibodies are valuable biomarkers for the diagnosis of RA and extra-articular manifestations. Antibodies against cyclic citrullinated peptides (anti-CCP2) and rheumatoid factor (RF) have been identified in patients with RA [6]. Previous studies have shown that the specificity and sensitivity of anti-CCP2 detection for the diagnosis of RA are 96–99% and 47–88% respectively,

dependent on the characteristics of the RA population [6,7]. Anti-CCP2 antibodies may be implicated in the pathogenesis of RA and are valuable for evaluating the erosive or non-erosive progression of articular injury in RA patients [8,9]. In addition, anti-CCP2 antibodies have been shown to be highly specific or independently associated with the development of extra-articular manifestations, including ischemic heart disease [10], type 1 diabetes mellitus [11], serositis [12], and subclinical atherosclerosis in patients with RA [13]. RF is another autoantibody most commonly detected in RA [1,14]. Detection of both anti-CCP2 and RF has additional values for the early diagnosis of RA, particularly for those with RA at early stage of the disease process [15].

However, the potential association of anti-CCP2 and RF with the development of ILD in RA patients remains controversial [16–18]. There is little information about whether anti-CCP2

antibodies or RF are associated with ILD in RA patients. In this study, we tested the levels of serum anti-CCP2 and RF in 285 patients with RA and analyzed the potential factors that were correlated with ILD in this population.

Methods

Ethics statement

The experimental protocol was approved by the Institute Review Board of Peking Union Medical College Hospital. All patients provide their written informed consent to participate in this study.

Study population

This study was approved by the Institute Review Board of Peking Union Medical College Hospital. A total of 285 patients with RA were recruited at the inpatient service of the Department of Rheumatology of Peking Union Medical College Hospital from January 2004 to October 2013. All patients fulfilled the criteria for the diagnosis of RA revised by the American College of Rheumatology (ACR) in 1987 [1]. Patients with RA-ILD were diagnosed by the presence of typical features in the lung by high-resolution computerized tomography (HRCT). The chest HRCT scans were evaluated by an expert radiologist in a blinded manner. According to the consensus for idiopathic interstitial pneumonias of the American Thoracic Society/European Respiratory Society (ATS/ERS) [19], the features of HRCT included irregular linear or reticular opacities, ground-glass opacities, consolidation, honeycombing, septal thickening, and traction bronchiectasis or bronchiolectasis. The disease activity of individual patients was evaluated by disease activity score in 28 joints (DAS28) [20,21]. Individuals were excluded if she/he had a history of ILD before the diagnosis of RA, other chronic lung diseases or incomplete medical record.

Clinical assessment

The demographic and clinical data of individual patients were retrospectively reviewed. These data included age, gender, disease duration, and cigarette smoking, co-existent autoimmune diseases, such as systemic lupus erythematosus (SLE), polymyositis/dermatomyositis (PM/DM), systemic sclerosis (SSc) and Sjögren's syndrome. The disease duration was defined from the onset of joint swelling and/or tenderness. Individuals with previous history of treatment with biological or general disease-modifying anti-rheumatic drugs (DMARDs) and corticosteroids for more than three months were recorded.

Blood samples were obtained from individual patients when they first visited to our institution. The positivity for serum anti-CCP2 (≥ 25 U/ml) and RF (≥ 15 IU/ml) in these patients was determined by enzyme linked immunosorbent assay (ELISA) using the specific kit (Euroimmun, Lübeck, Germany) and nephelometry method (Behring, Germany), respectively. The concentrations of serum C-reactive protein (CRP) as well as the value of erythrocyte sedimentation rate (ESR) in individual patients were routinely examined.

Statistical analyses

All values are expressed as the mean \pm SD or median with interquartile range (IQR) for normally and non-normally distributed data, respectively. The difference in individual variables between the RA-ILD and RA-only groups was analyzed by univariate model. Comparison of continuous data was performed using independent Student's *t* and Mann-Whitney non-parametric *U* test for normal and non-normal data respectively, and of

categorical data using the Pearson Chi-square test or Fisher's exact test. The stratification analyses for age and disease duration were conducted by Cochran-Mantel-Haenszel Chi-square test. The relationship of individual variables with RA-ILD was analyzed using a multivariate logistic regression model with stepwise selected variables. The collinearity of variables was assessed and explained by variance inflation factor (VIF) greater than 4.0 and tolerance less than 0.25 [22]. All statistical analyses were performed using the SPSS software (version 17.0, SPSS Inc., Chicago, IL, USA) and a two-tailed *P*-value of less than 0.05 was considered statistically significant.

Results

Patients and clinical characteristics

To determine the potential risk factors associated with ILD, a total of 285 patients with RA were recruited. Their demographic and clinical characteristics are summarized in **Table 1**. There were 71 patients with ILD, accounting for 24.9% in this population. The RA-ILD patients were significantly older and had significantly longer duration than that of the RA-only group of patients. However, there was no significant difference in the distribution of gender and in the percentages of smokers between the RA-only and RA-ILD groups of patients.

There was no significant difference in the percentages of complications and treatment history between these two groups of patients. Both groups of patients had similar levels of disease severity, the concentrations of serum CRP and ESR. There were 61 (21.4%) patients who had at least one other connective tissue disease (CTD), including 7 patients with SSc, 41 Sjögren's Syndrome, 4 PM/DM, and 16 SLE. There were 234 (82.1%), 180 (63.2%), 38 (13.3%) and 227 (79.6%) of patients receiving medication of prednisone, methotrexate, biological DMARD, or chemical DMARD, respectively (**Table 1**).

HRCT features and anti-CCP2 and RF status

In 71 patients with RA-ILD, HRCT scans of the lung revealed 48 cases with irregular line or reticular opacities, 22 cases with ground-glass attenuation, 26 cases with basal consolidation, 10 patients with honey-combing, 27 patients with septal thickening and 12 patients with traction bronchiectasis or bronchiolectasis. However, these HRCT characteristics were not significantly associated with the positivity of anti-CCP2 and RF in these patients (**Table 2**).

The relationship between RA-ILD and anti-CCP2 or RF

Characterization of the levels of serum anti-CCP2 and RF revealed that 207 (72.6%) serum samples were positive for anti-CCP2 and 211 (74.0%) were positive for RF in this population. Both the positivity rates of anti-CCP2 and RF were significantly higher ($p < 0.001$ and $p = 0.02$ respectively) in the group of RA-ILD than RA-only (**Table 1**). Cochran-Mantel-Haenszel stratification analyses indicated that the percentages of anti-CCP2 positivity were significantly associated with older age and longer disease duration (OR_{M-H} 3.3, 95% CI 1.5–7.1, $p = 0.003$ and OR_{M-H} 3.9, 95% CI 1.7–8.7, $p = 0.001$, respectively) (**Table 3**).

Univariate and multivariate logistic regression analysis

Univariate logistic regression analysis revealed that age and disease duration were risk factors for RA-ILD (crude odds ratio [cOR] 1.06, 95% CI 1.03–1.09, $p < 0.001$; and cOR 1.04, 95% CI 1.01–1.07, $p < 0.001$ respectively). Treatment with methotrexate (MTX) or other chemical DMARDs was not significantly associated with RA-ILD (cOR 0.58, 95% CI 0.34–1.01,

Table 1. The demographic and clinical characteristics of patients.

Characteristics	All patients	RA-ILD	RA-only	p-value
Patients, n (%)	285 (100)	71 (24.9)	214 (75.1)	—
Age, mean (SD), years	51.7 (13.4)	58.3 (11.2)	49.5 (13.4)	<0.001
Female, n (%)	211 (74.0)	50 (70.4)	161 (75.2)	0.44
Disease duration, median (IQR), years	5.0 (1.0–12.0)	9.0 (2.0–18.0)	4.0 (1.0–10.1)	0.003
Cigarette smoking, n (%)	59 (20.7)	18 (25.4)	41 (19.2)	0.31
Complication With other CTD*, n (%)	61 (21.4)	15 (21.1)	46 (21.5)	1.0
Laboratory test				
Anti-CCP2, n (%)	207 (72.6)	63 (88.7)	144 (67.3)	<0.001
RF, n (%)	211 (74.0)	60 (84.5)	151 (70.6)	0.02
CRP, mean (SD), mg/dL	40.2 (56.3)	42.6 (75.4)	39.4 (48.5)	0.68
ESR, mean (SD), mm/h	56.2 (34.5)	56.6 (33.2)	56.1 (35.0)	0.93
DAS28, mean (SD)	5.4 (1.7)	5.2 (1.8)	5.5 (1.7)	0.13
Treatment history, n (%)				
Prednisone	234 (82.1)	58 (81.7)	176 (82.2)	1.0
Methotrexate	180 (63.2)	38 (53.5)	142 (66.4)	0.07
Biological DMARD	38 (13.3)	6 (8.5)	32 (15.0)	0.23
Chemical DMARD	227 (79.6)	52 (73.2)	175 (81.8)	0.13

RA-ILD: rheumatoid arthritis-associated interstitial lung disease; RA-only: rheumatoid arthritis without interstitial lung disease; IQR: interquartile range; CTD: autoimmune disease; CRP: C-reactive protein; ESR: erythrocyte sedimentation rate; DAS28: disease activity score with 28 joints;

*Other CTD included systemic lupus erythematosus, polymyositis/dermatomyositis, systemic sclerosis and Sjögren's syndrome.

doi:10.1371/journal.pone.0092449.t001

$p = 0.05$, and cOR 0.61, 95% CI 0.32–1.14, $p = 0.12$; respectively). The positivity of anti-CCP2 and RF was significantly associated with ILD in RA patients (cOR 3.83; 95% CI 1.74–8.43, $p < 0.001$; and cOR 2.28; 95% CI 1.12–4.61, $p = 0.02$ respectively). Further stratification of low, moderate and high levels of RF indicated that only high levels of serum RF (≥ 364.0 IU/ml) were associated with ILD in RA patients ($p = 0.02$). After adjustment of age and disease duration together with the absence of collinearity (VIF and tolerance were 1.03 and 0.97 respectively) and other related co-variables (Table 4), the positivity of anti-CCP2 remained a risk factor for ILD in RA patients (adjusted odds ratio [aOR] 3.50; 95% CI 1.52–8.04, $p < 0.001$), regardless of the level of anti-CCP. However, after adjusting confounding factors including gender, age, disease duration, smoking, medications and other CTD complication, there was no significant association between any level of low, moderate and high positivity of RF and RA-ILD in this population (Table 4).

Discussion

The pulmonary manifestations in RA patients was firstly reported half a century ago [23] and affect 10 to 20% of RA patients, which is associated with increased mortality [24]. The ILD has been recognized as the most common complication in the lung of RA patients. In this retrospective study, we evaluated the potential factors that contributed to ILD in RA patients. We found that 24.9% RA patients suffered with ILD, which was higher than that reported [3,4]. The high rate of patients with ILD may be due to long disease duration in a large proportion of our RA patients. Indeed, we found that RA patients with ILD were significantly older and had longer duration than those patients without ILD in this population. Alternatively, the high rate of patients with ILD may stem from unique genetic background.

Some other risk factors for RA-ILD should also be considered in multivariate analysis of the associations of anti-CCP2 positivity with the RA-ILD. Firstly, a previous study has showed that ILD is a severe clinical entity presented in many

Table 2. The relationship between HRCT features and antibody status of anti-CCP2 and RF in patients with RA-ILD.

HRCT feature	Anti-CCP2 (+)	Anti-CCP2 (–)	p	RF (+)	RF (–)	p
Irregular line/reticular opacities, n (%)	43 (68.3)	5 (62.5)	0.71	39 (65.0)	9 (81.8)	0.49
Ground-glass attenuation, n (%)	21 (33.3)	1 (12.5)	0.42	20 (33.3)	2 (18.2)	0.48
Basal consolidation, n (%)	23 (36.5)	3 (37.5)	1.0	20 (33.3)	6 (54.5)	0.19
Honey-combing, n (%)	8 (12.7)	2 (25.0)	0.31	9 (15.0)	1 (9.1)	1.0
Septal thickening, n (%)	24 (38.1)	3 (37.5)	1.0	24 (40.0)	3 (27.3)	0.52
Traction bronchiectasis/bronchiolectasis, n (%)	9 (14.3)	3 (37.5)	0.13	10 (16.7)	2 (18.2)	1.0

doi:10.1371/journal.pone.0092449.t002

Table 3. Stratification analyses of the association of anti-CCP2 with ILD in RA patients.

	RA-ILD	RA-only	OR _{M-H} (CI 95%)	p
Age				
<46 years, n (%)	14 (15.1)	79 (84.9)		
47–58 years, n (%)	19 (19.4)	79 (80.6)	3.3 (1.5–7.1)	0.003
>59 years, n (%)	38 (40.4)	56 (59.6)		
Disease duration				
<1.9 years, n (%)	17 (20.5)	66 (79.5)		
2.0–9.9 years, n (%)	19 (18.6)	83 (81.4)	3.9 (1.7–8.7)	0.001
>10.0 years, n (%)	35 (35.0)	65 (65.0)		

OR_{M-H}: Mantel-Haenszel odds ratio. 95% CI: 95% confidence interval. Age and disease duration of all patients were divided into three equal groups.
doi:10.1371/journal.pone.0092449.t003

connective tissue diseases [25]. However, anti-CCP2 has been demonstrated to be extremely specific (96–98%) in patients with RA [26], but only 13.8% with SLE, 2.6% with SSc [27] and 7.5–9% with pSS [28,29]. In our study, 21.4% of patients had at least one other autoimmune disease besides RA and these patients distributed similarly in the RA-only and RA-ILD groups. Further univariate and multivariate analyses revealed that these comorbidities did not affect the significant association of the anti-CCP2 positivity with RA-ILD in this population. Secondly, we stratified patients, according to the radiologic features by HRC/T and we found that there was no significant difference in the anti-CCP2 positivity among these subgroups of patients, which was in consistent with a previous report [16]. Thirdly, MTX has been considered as the standard therapy of DMARDs for RA patients. Although MTX has been considered as a low toxic drug, MTX

treatment of patients with pre-existing RA-ILD can cause fatal complication [30]. In our study, we did not observe that treatment with MTX resulted in a fatal outcome. There was no significant difference in the anti-CCP2 positivity between the patients receiving MTX treatment and those without MTX treatment. Thus, MTX treatment may not affect the levels of serum anti-CCP2 in RA patients. Finally, we did not observe that regular smoking was associated with RA-ILD in this population of RA patients, which was in consistent with previous reports [31].

The positivity of anti-CCP2 and RF is valuable for the diagnosis of RA. We found that the percentages of patients with positive anti-CCP2 and/or RF in the RA-ILD group were significantly higher than that in the RA-only group. Further analysis revealed that the percentages of the older patients with anti-CCP2 positivity in the RA-ILD group were significantly higher than that in the RA-only group. The univariate model of logistic regression analyses indicated that the different levels of serum anti-CCP2, but only high levels of RF were associated with ILD in this population of RA patients. Further adjusting other covariates revealed that the positivity of anti-CCP2 was significantly correlated with ILD in RA patients. Our data were in consistent with recent reports in France [18] and Greece [12] and similar to that in Japan [32]. However, there was no significant difference in the positivity of anti-CCP2 between the RA patients with and without ILD in another Japanese population [16] and a Korean population [17]. The difference among these studies may come from difference in patient population, disease definition or methodology for detecting clinical parameters. The variable sensitivity and specificity of the methods for anti-CCP2 detection may also affect its value in evaluating the association with ILD. The significantly higher positivity in the RA-ILD patients indicated that the positivity of anti-CCP2 may be a good biomarker for the diagnosis of ILD in RA patients.

Table 4. Univariate and multivariate analyses of the associations of anti-CCP2 positivity with the RA-ILD.

	Univariate associations			Multivariate associations		
	β	cOR (95% CI)	p	β	aOR (95% CI)	p
Age	0.06	1.06 (1.03–1.09)	<0.001	0.05	1.06 (1.03–1.08)	<0.001
Disease duration	0.04	1.04 (1.01–1.07)	<0.001	0.04	1.04 (1.01–1.07)	0.02
Anti-CCP2						
Seropositive	1.34	3.83 (1.74–8.43)	<0.001	1.25	3.50 (1.52–8.04)	<0.001
Low positive	1.27	3.57 (1.46–8.76)	0.01	1.24	3.46 (1.35–8.89)	0.01
Moderate positive	1.54	4.67 (1.93–11.29)	0.00	1.44	4.22 (1.63–10.90)	0.003
High positive	1.20	3.32 (1.35–8.20)	0.01	1.08	2.96 (1.14–7.67)	0.03
RF						
Seropositive	0.82	2.28 (1.12–4.61)	0.02	–0.04	0.96 (0.38–2.44)	0.93
Low positive	0.74	2.10 (0.91–4.86)	0.08	0.38	1.46 (0.54–3.94)	0.45
Moderate positive	0.75	2.12 (0.93–4.82)	0.07	0.25	1.28 (0.45–3.66)	0.65
High positive	0.97	2.63 (1.16–5.93)	0.02	0.66	1.93 (0.68–5.47)	0.21

Univariate and multivariate associations of anti-CCP2 with the occurrence of RA-ILD was conducted by logistic regression with a backward stepwise model. Covariates in multivariate model included gender, age, disease duration, smoking, medication, and other CTD complication. β : partial regression coefficient; DMARDs: disease-modifying antirheumatic drugs; Anti-CCP2: anti-citrullinated peptide antibody; cOR: crude odds ratio; aOR: adjusted odds ratio; 95% CI: 95% confidence interval. Three equal groups of low, moderate and high positive anti-CCP2 were defined as $25.0 \leq \text{anti-CCP2} < 271.3$ U/ml, $271.3 \leq \text{anti-CCP2} < 941.2$ U/ml and $\text{anti-CCP2} \geq 941.2$ U/ml, respectively. Three equal groups of low, moderate and high positive RF were defined as $15.0 \leq \text{RF} < 114.0$ IU/ml, $114.0 \leq \text{RF} < 364.0$ IU/ml and $\text{RF} \geq 364.0$ IU/ml, respectively.
doi:10.1371/journal.pone.0092449.t004

We recognized that our study had limitations. We could not exclude the potential selection bias in our patients although individual patients were diagnosed as having ILD, according to the available diagnostic criteria [33]. Previous studies have estimated the prevalence of ILD in RA patients being about 3.7–33.0% in those without significant abnormal chest X-ray or respiratory symptoms [34–36]. It is possible that we may underestimate the prevalence of ILD in RA patients. In addition, we did exclude individual patients without complete medical records and those developed ILD prior to their RA diagnosis. Moreover, we had no complete medical records of all participants so that we were unable to evaluate whether a delay in treatment of patients with MTX and DMARDs can be a risk factor of ILD in RA patients. Finally, the small sample size in our study may interfere with some results, such as the lack of significant difference in the positivity of anti-CCP2 in different subtypes of ILD in this population. Furthermore,

small sample size can lead to a false positive result and overestimate the magnitude of an association, particularly in a situation with multiple confounding variants. Therefore, further study in a bigger population is warranted to validate the value of anti-CCP2 positivity in evaluating the prognosis of ILD in RA patients.

In summary, we found significantly higher positivity of anti-CCP2 in RA patients with ILD than that in the RA alone patients. Our findings indicated that the positivity of anti-CCP2 is associated with ILD in RA patients.

Author Contributions

Conceived and designed the experiments: YY Yongzhe Li XZ FZ FT XZ. Performed the experiments: YY DL LZ Yang Li WL YR. Analyzed the data: YY DL GS. Contributed reagents/materials/analysis tools: Yongzhe Li XZ FZ FT XZ. Wrote the paper: YY DL.

References

- Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, et al. (1988) The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 31: 315–324. PubMed: 3358796
- Turesson C, O'Fallon WM, Crowson CS, Gabriel SE, Matteson EL (2003) Extra-articular disease manifestations in rheumatoid arthritis: incidence trends and risk factors over 46 years. *Ann Rheum Dis* 62: 722–727. PubMed: 12860726
- Koduri G, Norton S, Young A, Cox N, Davies P, et al. (2010) Interstitial lung disease has a poor prognosis in rheumatoid arthritis: results from an inception cohort. *Rheumatology (Oxford)* 49: 1483–1489. PubMed: 20223814
- Olson AL, Swigris JJ, Sprunger DB, Fischer A, Fernandez-Perez ER, et al. (2011) Rheumatoid arthritis-interstitial lung disease-associated mortality. *Am J Respir Crit Care Med* 183: 372–378. PubMed: 20851924
- Turesson C, Matteson EL (2004) Management of extra-articular disease manifestations in rheumatoid arthritis. *Curr Opin Rheumatol* 16: 206–211. PubMed: 15103246
- Zeng X, Ai M, Tian X, Gan X, Shi Y, et al. (2003) Diagnostic value of anti-cyclic citrullinated Peptide antibody in patients with rheumatoid arthritis. *J Rheumatol* 30: 1451–1455. PubMed: 12858440
- Rodriguez-Mahou M, Lopez-Longo FJ, Sanchez-Ramon S, Esteche A, Garcia-Segovia A, et al. (2006) Association of anti-cyclic citrullinated peptide and anti-Sa/citrullinated vimentin autoantibodies in rheumatoid arthritis. *Arthritis Rheum* 55: 657–661. PubMed: 16874789
- Rantapää-Dahlqvist S, de Jong BAW, Berglin E, Hallmans G, Wadell G, et al. (2003) Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis & Rheumatism* 48: 2741–2749. PubMed: 14558078
- Reparon-Schuijt CC, van Esch WJ, van Kooten C, Schellekens GA, de Jong BA, et al. (2001) Secretion of anti-citrulline-containing peptide antibody by B lymphocytes in rheumatoid arthritis. *Arthritis Rheum* 44: 41–47. PubMed: 11212174
- Lopez-Longo FJ, Oliver-Minarro D, de la Torre I, Gonzalez-Diaz de Rabago E, Sanchez-Ramon S, et al. (2009) Association between anti-cyclic citrullinated peptide antibodies and ischemic heart disease in patients with rheumatoid arthritis. *Arthritis Rheum* 61: 419–424. PubMed: 19333979
- Liao KP, Gunnarsson M, Kallberg H, Ding B, Plenge RM, et al. (2009) Specific association of type 1 diabetes mellitus with anti-cyclic citrullinated peptide-positive rheumatoid arthritis. *Arthritis Rheum* 60: 653–660. PubMed: 19248096
- Alexiou I, Germeis A, Koutroumpas A, Kontogianni A, Theodoridou K, et al. (2008) Anti-cyclic citrullinated peptide-2 (CCP2) autoantibodies and extra-articular manifestations in Greek patients with rheumatoid arthritis. *Clin Rheumatol* 27: 511–513. PubMed: 18172572
- Gerli R, Bartoloni Bocci E, Sherer Y, Vaudo G, Moscatelli S, et al. (2008) Association of anti-cyclic citrullinated peptide antibodies with subclinical atherosclerosis in patients with rheumatoid arthritis. *Ann Rheum Dis* 67: 724–725. PubMed: 18408112
- Scott DL (2000) Prognostic factors in early rheumatoid arthritis. *Rheumatology (Oxford)* 39 Suppl 1: 24–29. PubMed: 11001376
- Jansen AL, van der Horst-Bruinsma I, van Schaardenburg D, van de Stadt RJ, de Koning MH, et al. (2002) Rheumatoid factor and antibodies to cyclic citrullinated Peptide differentiate rheumatoid arthritis from undifferentiated polyarthritis in patients with early arthritis. *J Rheumatol* 29: 2074–2076. PubMed: 12375314
- Inui N, Enomoto N, Suda T, Kageyama Y, Watanabe H, et al. (2008) Anti-cyclic citrullinated peptide antibodies in lung diseases associated with rheumatoid arthritis. *Clin Biochem* 41: 1074–1077. PubMed: 18638466
- Jearn LH, Kim TY (2012) Level of anticitrullinated peptide/protein antibody is not associated with lung diseases in rheumatoid arthritis. *J Rheumatol* 39: 1493–1494. PubMed: 22753807
- Aubart F, Crestani B, Nicaise-Roland P, Tubach F, Bollet C, et al. (2011) High levels of anti-cyclic citrullinated peptide autoantibodies are associated with co-occurrence of pulmonary diseases with rheumatoid arthritis. *J Rheumatol* 38: 979–982. PubMed: 21362759
- American Thoracic Society, European Respiratory Society (2002) American Thoracic Society/European Respiratory Society International Multidisciplinary Consensus Classification of the Idiopathic Interstitial Pneumonias. This joint statement of the American Thoracic Society (ATS), and the European Respiratory Society (ERS) was adopted by the ATS board of directors, June 2001 and by the ERS Executive Committee, June 2001. *Am J Respir Crit Care Med* 165: 277–304. PubMed: 11790668
- van der Heijde DM, van 't Hof MA, van Riel PL, Theunisse LA, Lubberts EW, et al. (1990) Judging disease activity in clinical practice in rheumatoid arthritis: first step in the development of a disease activity score. *Ann Rheum Dis* 49: 916–920. PubMed: 2256738
- Zhang W, Shi Q, Zhao LD, Li Y, Tang FL, et al. (2010) The safety and effectiveness of a chloroform/methanol extract of *Tripterygium wilfordii* Hook F (T2) plus methotrexate in treating rheumatoid arthritis. *J Clin Rheumatol* 16: 375–378. PubMed: 21085018
- Samouda H, Dutour A, Chaumoitre K, Panuel M, Dutour O, et al. (2013) VAT = TAAT-SAAT: innovative anthropometric model to predict visceral adipose tissue without resort to CT-Scan or DXA. *Obesity (Silver Spring)* 21: E41–50. PubMed: 23404678
- Ellman P, Ball RE (1948) Rheumatoid disease with joint and pulmonary manifestations. *Br Med J* 2: 816–820. PubMed: 18890308
- Brown KK (2007) Rheumatoid lung disease. *Proc Am Thorac Soc* 4: 443–448. PubMed: 17684286
- Castelino FV, Varga J (2010) Interstitial lung disease in connective tissue diseases: evolving concepts of pathogenesis and management. *Arthritis Res Ther* 12: 213. PubMed: 20735863
- van Noord C, Hooijkaas H, Dufour-van den Goorbergh BC, van Hagen PM, van Daele PL, et al. (2005) Diagnostic value of anti-cyclic citrullinated peptide antibodies to detect rheumatoid arthritis in patients with Sjogren's syndrome. *Ann Rheum Dis* 64: 160–162. PubMed: 15608321
- Morita Y, Muro Y, Sugiura K, Tomita Y (2008) Anti-cyclic citrullinated peptide antibody in systemic sclerosis. *Clin Exp Rheumatol* 26: 542–547. PubMed: 18799082
- Tobon GJ, Correa PA, Anaya JM (2005) Anti-cyclic citrullinated peptide antibodies in patients with primary Sjogren's syndrome. *Ann Rheum Dis* 64: 791–792. PubMed: 15834066
- Gottenberg JE, Mignot S, Nicaise-Rolland P, Cohen-Solal J, Aucouturier F, et al. (2005) Prevalence of anti-cyclic citrullinated peptide and anti-keratin antibodies in patients with primary Sjogren's syndrome. *Ann Rheum Dis* 64: 114–117. PubMed: 15231509
- Saravanan V, Kelly CA (2004) Reducing the risk of methotrexate pneumonitis in rheumatoid arthritis. *Rheumatology (Oxford)* 43: 143–147. PubMed: 12923285
- Ayhan-Ardic FF, Oken O, Yorgancioglu ZR, Ustun N, Gokharman FD (2006) Pulmonary involvement in lifelong non-smoking patients with rheumatoid arthritis and ankylosing spondylitis without respiratory symptoms. *Clin Rheumatol* 25: 213–218. PubMed: 16091838
- Mori S, Koga Y, Sugimoto M (2012) Different risk factors between interstitial lung disease and airway disease in rheumatoid arthritis. *Respir Med* 106: 1591–1599. PubMed: 22867979

33. Bongartz T, Nannini C, Medina-Velasquez YF, Achenbach SJ, Crowson CS, et al. (2010) Incidence and mortality of interstitial lung disease in rheumatoid arthritis: a population-based study. *Arthritis Rheum* 62: 1583–1591. PubMed: 20155830
34. Zrour SH, Touzi M, Bejjia I, Golli M, Rouatbi N, et al. (2005) Correlations between high-resolution computed tomography of the chest and clinical function in patients with rheumatoid arthritis. Prospective study in 75 patients. *Joint Bone Spine* 72: 41–47. PubMed: 15681247
35. Gochuico BR, Avila NA, Chow CK, Novero LJ, Wu HP, et al. (2008) Progressive preclinical interstitial lung disease in rheumatoid arthritis. *Arch Intern Med* 168: 159–166. PubMed: 18227362
36. Brusselle G (2010) Rheumatoid arthritis and interstitial lung disease. *Rheumatology (Oxford)* 49: 1425–1426. PubMed: 20223815