



Behavioral Plasticity in Ant Queens: Environmental Manipulation Induces Aggression among Normally Peaceful Queens in the Socially Polymorphic Ant *Leptothorax acervorum*

Jürgen Trettin*, Thomas Seyferth, Jürgen Heinze

Biologie I, Universität Regensburg, Regensburg, Germany

Abstract

The behavioral traits that shape the structure of animal societies vary considerably among species but appear to be less flexible within species or at least within populations. Populations of the ant *Leptothorax acervorum* differ in how queens interact with other queens. Nestmate queens from extended, homogeneous habitats tolerate each other and contribute quite equally to the offspring of the colony (polygyny: low reproductive skew). In contrast, nestmate queens from patchy habitats establish social hierarchies by biting and antennal boxing, and eventually only the top-ranking queen of the colony lays eggs (functional monogyny: high reproductive skew). Here we investigate whether queen-queen behavior is fixed within populations or whether aggression and high skew can be elicited by manipulation of socio-environmental factors in colonies from low skew populations. An increase of queen/worker ratio and to a lesser extent food limitation elicited queen-queen antagonism in polygynous colonies from Nürnberger Reichswald similar to that underlying social and reproductive hierarchies in high-skew populations from Spain, Japan, and Alaska. In manipulated colonies, queens differed more in ovarian status than in control colonies. This indicates that queens are in principle capable of adapting the magnitude of reproductive skew to environmental changes in behavioral rather than evolutionary time.

Citation: Trettin J, Seyferth T, Heinze J (2014) Behavioral Plasticity in Ant Queens: Environmental Manipulation Induces Aggression among Normally Peaceful Queens in the Socially Polymorphic Ant *Leptothorax acervorum*. PLoS ONE 9(4): e95153. doi:10.1371/journal.pone.0095153

Editor: Fabio S. Nascimento, Universidade de São Paulo, Faculdade de Filosofia Ciências e Letras de Ribeirão Preto, Brazil

Received: December 4, 2013; **Accepted:** March 24, 2014; **Published:** April 17, 2014

Copyright: © 2014 Trettin et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Deutsche Forschungsgemeinschaft (He1623/25) including the funding programme Open Access Publishing. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: juergen.trettin@ur.de

Introduction

The organization of animal societies and the underlying behavioral traits vary considerably among species but appear to be remarkably robust within species or at least within populations (e.g., [1–3]). For example, the share of individual group members in the offspring produced by the group (“reproductive skew”) ranges from equal partitioning among mutually tolerant group members (“low skew”) to reproduction being the privilege of only one individual, which dominates all other individuals (“high skew”); [4–6]. Reproductive skew may be affected by ecological constraints on solitary nesting, the relatedness and relative fighting strength of group members, and other factors (see e. g., [4], [7–9] and references therein). Consequently, reproductive skew and the causative behavioral processes typically vary less within than between species or populations [10–11].

In most social insects, queens in mature multi-queen societies show little antagonism and contribute equally to the offspring of the group (“polygyny”). High or even maximal skew has been described only from a small number of species with multi-queen colonies, where only one of several inseminated queens monopolizes reproduction ([12–15]). This “functional monogyny” [13] results from the formation of social rank orders among potential reproductives through overtly aggressive or ritualized dominance

behavior (e.g., [12], [14–17]). The magnitude of reproductive skew and/or the associated behavior of queens in mature or founding colonies appear to be largely fixed within species [13] or at least within populations [18–21]. Whether queens are capable of adaptively reacting to changed environmental conditions, as assumed by models of optimal skew [22], has rarely been investigated.

Here, we examine whether experimental manipulation of colonies from a population with mutually tolerant queens and low skew can elicit queen-queen fighting similar to that underlying reproductive hierarchies in high-skew populations, i.e., whether queens are in principle capable of adjusting their reproductive behavior to changed environmental conditions. Our study species, the Holarctic ant *Leptothorax acervorum* (Fabricius, 1793), is socially polymorphic and the magnitude of reproductive skew varies among populations [18–19]. Sexu-als of *L. acervorum* mate in late summer and thereafter young, mated queens may seek re-adoption into their natal nests. In low-skew populations, several nestmate queens continue to co-exist peacefully and lay eggs [18,23–26]. In contrast, queens in high-skew populations engage in aggressive interactions and form dominance hierarchies after queen re-adoption and again after hibernation. Subordinate queens may stay in the nest as hopeful reproductives, leave the nest alone or together with workers from the natal nest to start a new colony, or

are expelled through attacks by dominant queens and/or the workers [27–31]. Low-skew populations are abundant in the extended coniferous forests of boreal and alpine Central and Northern Eurasia with a high density of available nest sites for solitary foundresses [13,23–26]. In contrast, high skew populations with social and reproductive rank orders live in quickly saturated habitat patches with nest site limitation [19,27–31]. The distributional pattern of social organization parallels predictions from optimal skew models in that high skew appears to be associated with ecological constraints on solitary founding [32].

We aimed to elicit the formation of rank orders and high reproductive skew in colonies from a low-skew population by deteriorating the environmental conditions for queens. To do so, we experimentally increased queen-worker ratios in natural colonies and/or limited the availability of food. These treatments were motivated by studies according to which queen-worker ratios are considerably higher in functionally monogynous than polygynous species of *Leptothorax* [33] and food shortage may lead to increased skew in both ants [34–35] and social spiders [36]. Assuming that queen behavior and reproductive skew are plastic traits, we expected both manipulations to provoke queen-queen aggression and to lead to more pronounced differences in ovarian development. We show that in particular the reduction of worker number leads to queen fighting and more strongly skewed ovarian development.

Materials and Methods

Ant Collection and Rearing

Colonies of *L. acervorum* were collected from their nests in rotting branches in the well-studied low-skew population in Nürnberger Reichswald (June and August 2011), a pine-dominated forest near Nuremberg, Southern Germany (49°16'N, 11°10'E). Whole colonies were extracted from their nests in rotting twigs and transferred into standard three-chambered plastic boxes (10 cm×10 cm×3 cm) with plaster floor and reared under artificial spring/autumn conditions (12 h/12 h 20°C/10°C) in incubators as previously described [28], [37].

Experimental Set-up

From freshly collected colonies we set up experimental colonies with 40 workers, 30 brood items and three to a maximum of seven dealate (wingless) queens found in the respective field colony. Queens were marked individually with 30 µm thin wires (copper or red enameled) tied between alitrunk and petiole, petiole and postpetiole, or postpetiole and gaster. Within three days after the set up, we subjected the experimental colonies to the following four treatments (nine colonies per treatment): control colonies without stressor, food-stressed colonies, severely worker-reduced colonies from which 20 workers were removed at the start of the experiment, and colonies which received a combined treatment of food- and worker-reduction. Colonies were assigned to the four treatment groups so that the averages and the distributions of initial queen numbers did not differ significantly among treatment groups (total initial queen numbers per treatment, control = 44, food-stressed = 49, worker-reduced = 45, food- and worker-reduced = 45; Kruskal-Wallis test, $H_3 = 1.331$, $p = 0.727$; Kolmogorov-Smirnov two-sample tests, all $D < 0.444$, all $p > 0.336$).

Control and worker-reduced colonies were fed three-times during the observation period with chopped cockroaches and diluted honey ad libitum, food-stressed colonies only once. In the field, colonies often experience long periods during which workers cannot forage because of bad weather. Our food limitation experiment did not provide a similarly drastic reduction and

colonies can well survive under this condition, albeit without investing a lot of resources into new brood. An observation time of 10 days was chosen as a trade-off, to keep the stressful period as short as possible and, at the same time, to guarantee the minimal sample sizes required to yield statistically meaningful results.

To keep queen-worker ratios high, we removed dark pupae and callow workers from the worker-reduced colonies (with and without food stress) on the fifth day of the observation period. This additional manipulation did not affect the behavior of queens and the frequency of aggression was similar on day five and six in the worker-reduced treatments (Wilcoxon signed rank test, $W: V = 3$, $n_1 = n_2 = 8$, $p = 0.3711$; FW: $V = 3$, $n_1 = n_2 = 9$, $p = 1.0$).

The experiment was carried out during two different observation periods (first round: five colonies per treatment, 2011-07-03 to 2011-07-12; second round: four colonies per treatment, 2011-08-29 to 2011-09-18. During this latter period not all experimental colonies could be simultaneously subjected to the various treatments).

Observation and Ovary Dissection

Observations were started two days after the experimental manipulation. Each colony was observed under a binocular microscope in 20-min sessions twice per day over a period of ten consecutive days (total observation time per colony 400 min). Behavior was recorded by scan sampling every five minutes and in addition by ad libitum sampling [38]. The occurrence of all interactions involving queens (antennal boxing, mandible opening, biting, pulling, stinging/smearing, egg eating, egg laying, grooming, and trophallaxis, i.e., exchange of liquid food) was counted.

After the experiment, we killed all queens by freezing at -20°C and dissected their ovaries under a binocular microscope as described in [39]. We noted the presence of sperm in the spermatheca, corpora lutea, and mature oocytes and classified ovarian status as follows [16]: I undeveloped ovarioles, II slightly elongated ovarioles with a few immature oocytes, III elongated ovarioles, but no eggs in development (degenerated), IV fully elongated ovarioles with maturing oocytes, V fully elongated ovarioles with corpora lutea, but no eggs in development (degenerated). In addition, we recognized two intermediate stages (II–IV and IV–V).

Eighty-six percent of the observed queens ($n = 155$) were inseminated, the others were uninseminated and had shed their wings in the field without mating. Behavior of these virgin queens was excluded from the analysis because in *Leptothorax* they take over worker roles [40] and consequently do not represent an adequate substitute for mated queens. All colonies used in the analysis contained at least two mated queens except one worker-reduced colony, which was excluded from statistical analysis (for details on queen numbers per colony and treatment see Table S2). The exclusion of virgin queens or queens that died during the experiment led to slightly lowered average queen numbers relative to the beginning of the experiment, but as before neither average queen numbers nor the distribution of queen numbers were significantly different among the four treatments (total final queen number per treatment, control = 38, food-stressed = 45, worker-reduced = 34, food- and worker-reduced = 38; KW-test: $H_3 = 3.1086$, $p = 0.375$; Kolmogorov-Smirnov two sample tests, all $D < 0.333$, all $p > 0.699$).

As was the aim of our manipulation, queen-worker ratios were significantly lower in control/food-stressed colonies than in both types of worker-reduced colonies (Mann-Whitney U test: $U = 33.5$, $n_1 = 18$, $n_2 = 17$, $p < 0.001$). Experimental queen-worker ratios in control and food-stressed colonies were well within the range of queen-worker ratios from natural colonies in the low-skew




Figure 1. Frequency of aggression among queens of the ant *Leptothorax acervorum* from a low-skew population. Data shown as attacks per queen during the total observation period (median, quartiles, range). Individual colonies were subjected to different types of stress (food reduction F, worker reduction W, or both FW) or left unmanipulated (control C) in two different seasons (July, I, and September, II). Outliers are indicated as ○ and original data points as ×.
doi:10.1371/journal.pone.0095153.g001

population, while queen-worker ratios in worker-reduced colonies were similar to those previously reported from high skew populations (see Figure S1 and Table S1; but see [19]). The occurrence of queen mortality and the fact that the reproductive status of queens could only be determined after the observations ultimately resulted in a marginal overlap in queen-worker ratios among treatments (see boxplots A – D in Figure S1).

Data Analysis

Preliminary analyses in a subset of colonies showed that egg-laying rate, the frequency of grooming between queens and the rate of aggression and trophallaxis from workers towards queens were too rare to give meaningful results in statistical test. We therefore omitted these types of behavior from the final statistical analysis.

We analyzed the effect of the different treatments on the respective behavioral responses per colony by Scheirer-Ray-Hare tests (SRH), a non-parametric equivalent for a multi-way ANOVA [41], with worker reduction and food reduction as independent factors. To account for the two different observation periods we included “time” as a third factor. *P*-values were adjusted for multiple tests (queen-queen aggression, egg eating, trophallaxis, grooming) using sequential Bonferroni corrections [42]. In addition, we used Kruskal-Wallis tests to analyze data for an overall effect of different treatments on queen-queen aggression per colony for each observation period separately. To determine whether queens contribute equally to aggression we estimated the B-index [43]. For the statistical comparison of ovarian development among treatments we combined manipulated colonies due to the small sample size in each category.

All statistical analyses were carried out in R version 2.14.1 to 3.0.1 [44] or PAST v. 1.75b [45].

Ethics statement

As no protected species was sampled and colonies were collected only from a state-owned forest, no permits and approval for ant collection were required. All experiments comply with national and international law.

Results

Queen-Queen and Worker-Queen Behavior

Experimental manipulation of colonies from the low-skew Reichswald population resulted in queen-queen antagonism similar in quality and quantity to that previously observed in high-skew populations (antennal boxing, mandible opening, biting, pulling and stinging/smearing; [28], [30], [31]). In total, we observed 187 attacks among queens during 230 hours of direct observation (107 instances of antennal boxing, 21 threats with opened mandibles, 53 bites, 4 dragging on legs or antennae, 2 stings). The occurrence of aggressive behavior differed greatly among the four treatments and also between the observation periods (Figure 1). A Scheirer-Ray-Hare test gave evidence for a strong influence of the factors “observation period” (SRH: $H_1=12.68$, $n=26$, $p=0.0005$; corrected for multiple tests, $p'=0.0021$) and “worker reduction” (SRH: $H_1=10.17$, $p=0.0014$, $p'=0.0057$). Averaged over all treatments, more aggression occurred during the second observation period (median, quartiles, first observation period: 0, 0, 0.617 attacks per queen, $n=20$; second observation period: 2.167, 0.548, 2.928 attacks per queen, $n=15$, Mann-Whitney *U* test: $U=50$,

$p < 0.0008$). Statistical analysis also revealed significant differences between treatments for each observation period separately (Kruskal-Wallis tests, first observation period: $H_3 = 10.88$, $p = 0.01239$; second observation period: $H_3 = 8.79$, $p = 0.0323$).

In accordance with previous observations [18], queen-queen aggression almost never occurred in control colonies (median, quartiles 0, 0, 0.2 attacks per queen). The reduction of worker number alone and in combination with food reduction resulted in a considerable increase of aggressive behavior among queens (median, quartiles 2.1, 0.6, 2.9 attacks per queen). In principle, the drastically changed queen-worker ratio, the lower worker number and the experimental removal of workers and worker pupae might all have elicited queen-queen antagonism. However, the following observation suggests that queen aggression is a consequence of manipulated queen-worker ratio and not of disturbance or lower worker number alone: in one of the nine worker-reduced colonies, in which dissection at the end of the experiment revealed that several queens were not inseminated and queen-worker ratio thus was not greatly changed, no aggression was observed. Furthermore, over all worker-reduced colonies, the number of attacks appeared to increase with queen-worker ratio, albeit not significantly so (Gamma correlation: $\Gamma = 0.353$, $n = 17$, $p = 0.083$). Neither food-reduction nor worker-reduction had a significant effect on the level of aggression from workers towards queens (SRH, food-reduction: $H_1 = 1.30$, $p = 0.255$; worker-reduction: $H_1 = 0.13$, $p = 0.717$).

Food reduction alone did not have a significant effect on queen aggressiveness (SRH: $H_1 = 1.15$, $p = 0.283$, $p' = 0.850$). In none of the behaviors were interactions among the various factors significant (all $p > 0.07$). The short duration of the experimental manipulation did not allow deducing rank orders from the aggression. Nevertheless, the contribution of individual queens to the aggression was significantly different from random in three of those eight colonies with worker reduction (and worker reduction plus food reduction), in which more than 10 aggressive interactions among queens were observed (B-indices of 0.304, 0.978, and 0.214, with the confidence intervals not overlapping 0). Worker reduction in addition led to an increase of egg eating by queens (SRH: $H_1 = 6.19$, $n = 26$, $p = 0.0129$, $p' = 0.0350$, Figure S2), and food reduction significantly increased the frequency of trophallaxis between queens (SRH: $H_1 = 9.31$, $n = 27$, $p = 0.0023$, $p' = 0.0091$, Figure S3). Queens from worker-reduced colonies were more frequently groomed by workers than queens from food-reduction colonies (SRH: $H_1 = 6.36$, $n = 26$, $p = 0.0117$, $p' = 0.0350$, Figure S4).

Ovarian status

Ovarian status differed greatly between the observation periods. The ovaries of queens observed in September were elongated and contained corpora lutea, suggesting that the queens had been fully fertile, but they rarely contained maturing oocytes. Only six of 80 queens had still fully developed ovaries (stage IV), while the majority of queens had ovaries of stage V and probably had started to prepare for hibernation. We therefore did not examine differences in ovarian status among the different treatments in these colonies. In contrast, in July all five control colonies and nine of 14 stressed colonies (3 of 5 colonies with food reduction, 4 of 5 colonies with worker reduction and 2 of 4 colonies with both manipulations), contained one or several fully fertile queens with stage IV ovaries. Four of five control colonies but only two of 14 stressed colonies contained two or more fully fertile queens (Fisher's exact test, $p = 0.017$). Other queens had degenerated their ovaries.

Discussion

We investigated whether queens from a low-skew population of the ant *Leptothorax acervorum* adjust their behavior towards nestmate queens and the partitioning of reproduction in response to experimentally changed conditions. Our data show that experimental manipulation, in particular the reduction of worker numbers, provoked fighting and dominance interactions similar in quality and quantity to those previously observed in high-skew populations of this and other *Leptothorax* species. Though the absolute number of queen-queen attacks was low, few and infrequent interactions may suffice to establish clear social and reproductive rank orders among ants (e.g., [31]). Food reduction alone did not lead to an increase in aggression but resulted in a higher frequency of food begging and food exchange among queens.

In high skew populations of *L. acervorum* and other functionally monogynous ants queen-queen aggression establishes social rank orders among queens [14], [16], [28], [30], [31]. The incidence of aggression peaks directly after hibernation when queens begin to mature eggs and again in fall when queens prepare for hibernation and young, adopted queens integrate themselves into the hierarchy [46]. This temporal fluctuation of aggressiveness might explain the large difference in the frequency of queen antagonism between the two observation periods also in our study. The rank order of queens in functionally monogynous ants determines their reproductive status, and usually only the top-ranking queen lays eggs.

Dissections of queens in our experiment indicated that already after ten days of manipulation, queens from worker- and food-stressed colonies differed more strongly in their ovarian status than queens from control colonies. In most stressed colonies, only one queen had remained fully fertile, while other queens had begun to degenerate their ovaries. We cannot exclude that our experimental manipulation directly affected ovarian status in some queens more than in others. However, comparison with natural high-skew colonies suggests that the observed variation results from aggression and/or an increased rate of food exchange from subordinate to dominant queens. Begging food from subordinates is a subtle mechanism of domination through which queens in related species deplete workers of the resources needed to produce eggs [47], and food stress was reported to increase reproductive skew in the ant *Myrmica rubra* [34]. In our study, worker reduction also led to an increased frequency of egg-eating by queens. Egg eating by *L. acervorum* queens is well documented, and queens in low skew populations apparently do not discriminate between their own eggs and those laid by other queens [23], [48]. Such indiscriminate oophagy does not alter stable skew. In contrast, Ito [30] observed that dominant queens from a high-skew population on Hokkaido fed on eggs immediately after these were laid by subordinate queens and in this way increased reproductive skew. Unfortunately, we did not determine the origin of eggs in our study, but it is possible that egg eating would have increased skew in stressed colonies.

Ideally we should also have performed the opposite experiment, i.e., trying to induce queen tolerance in colonies from high skew populations of *L. acervorum* by adding workers or overfeeding the colonies. However, documenting the complete disappearance of a rare behavioral trait is more difficult than documenting its induction. Furthermore, queen aggressiveness and reproductive skew appear to vary to some extent among natural colonies of high skew populations of *L. acervorum* and related species, and not in all studied colonies were queens seen to engage in aggressive interactions [19], [28], [31], [49].

Our result that an increase of the queen-worker ratio increases queen-queen antagonism and reproductive skew matches the observation that queen-worker ratio is higher in high-skew species than in low-skew species [33]. Queen-worker ratio might co-vary with ecological constraints on solitary colony founding that have been suggested to affect the magnitude of reproductive skew. A high queen-worker ratio can result from a high re-adoption rate of young queens or high mortality of workers, e. g., during hibernation. Both might reflect adverse environmental conditions, which pose high costs on dispersal and colony founding by solitary queens [32]. In addition, the chance for individual queens to found new colonies accompanied by workers might be more limited at higher queen-worker ratios.

The easy induction of queen-queen aggression and increased reproductive skew in colonies in which queens normally do not fight is at odds with results from previous studies on social plasticity in ants. For example, the social structure of colonies of the red imported fire ant, *Solenopsis invicta*, depends on the genotypes of queens and workers at a particular locus [50], and whether co-founding queens of *Messor pergandei* and *Pogonomyrmex californicus* exhibited aggression or were mutually tolerant did not depend on group size but on region of origin [20], [21]. Similarly, queens of *Temnothorax rugatulus* were not capable of expanding their behavior to that shown during the workerless colony founding phase under experimental worker shortage [51]. Our finding also contrasts with previous studies in *L. acervorum*: dissections and population genetic analyses suggest that the level of skew is more or less fixed within populations, and that colonies maintain their social phenotype when kept under standardized laboratory conditions [18], [19], [25], [26]. Therefore, different levels of reproductive skew have been suggested to be an evolved rather than a behavioral response [19], [49]. This discrepancy might mean that ants from different populations have evolved different thresholds for fighting and dominance behavior, leading to more or less fixed local social organization. In addition, variation in skew within populations may have been underestimated by genetic analyses and dissections. Skew estimates from the genetic assignment of maternity of adults are often difficult because of the typically high relatedness among nestmate queens and rapid queen turnover. Genotyping of eggs and dissections give only a snap-shot account of the status quo of the partitioning of brood production. As it may be difficult to distinguish between formerly active, but now degenerate ovaries and not yet activated ovaries, the past or future contributions of queens that presently do not mature eggs may be underrated.

Conclusions

Our study demonstrates that queens of *L. acervorum* from low skew populations are able to react to changes in their social environment. In addition, it highlights the importance of queen-worker ratio for the adjustment of skew and the need for further studies to clarify its role and the role of other factors (e. g., egg cannibalism and habitat structure) in the formation and maintenance of reproductive skew in insect societies.

References

1. Wilson EO (1971) *The Insect Societies*. Cambridge, Mass: Harvard University Press. 548 p.
2. Ostfeld RS (1990) The ecology of territoriality in small mammals. *Trends Ecol Evol* 5: 411–415.
3. Linklater WL (2000) Adaptive explanation in socio-ecology: lessons from the Equidae. *Biol Rev* 75: 1–20.
4. Keller L, Reeve HK (1994) Partitioning of reproduction in animal societies. *Trends Ecol Evol* 9: 98–102.
5. Magrath RD, Heinsohn RG (2000) Reproductive skew in birds: models, problems and prospects. *J Avian Biol* 31: 247–258.
6. Reeve HK, Keller L (2001) Tests of reproductive-skew models in social insects. *Annu Rev Entomol* 46: 347–385.
7. Reeve HK, Ratnieks FLW (1993) Queen-queen conflicts in polygynous societies: mutual tolerance and reproductive skews. In: Keller L, editor. *Queen Number and Sociality in Insects*. Oxford: Oxford University Press, pp. 45–85.
8. Johnstone RA (2000) Models of reproductive skew: a review and synthesis. *Ethology* 106: 5–26.

Supporting Information

Figure S1 Queen-worker ratios of experimental treatments from this study (grey) as well as natural queen-worker ratios from several low skew (red) and high skew (blue) populations. (A: controls, B: food reduction, C: worker reduction, D: both treatments (FW), E–K: correspond to references in Table S1; E = 1, F = 2, G & H & I = 3, J = 4, K = 5 & 6). (TIF)

Figure S2 Frequency of egg eating (observations per queen during the total observation period, median, quartiles, range) in colonies of the ant *Leptothorax acervorum* from the low-skew population in Nürnberger Reichswald. Individual colonies were subjected to different types of stress (food reduction F, worker reduction W, or both FW) or left unmanipulated (control C) in two different seasons (July, I, and September, II). Outliers are indicated as circles. (TIF)

Figure S3 Frequency of trophallaxis, i.e., food exchange, among queens (observations per queen during the total observation period, median, quartiles, range) of the ant *Leptothorax acervorum* from the low-skew population in Nürnberger Reichswald. Individual colonies were subjected to different types of stress (food reduction F, worker reduction W, or both FW) or left unmanipulated (control C) in two different seasons (July, I, and September, II). Outliers are indicated as circles. (TIF)

Figure S4 Frequency of queens groomed by workers (observations per queen and worker during the total observation period, median, quartiles, range) of the ant *Leptothorax acervorum* from the low-skew population in Nürnberger Reichswald. Individual colonies were subjected to different types of stress (food reduction F, worker reduction W, or both FW) or left unmanipulated (control C) in two different seasons (July, I, and September, II). Outliers are indicated as circles. (TIF)

Table S1 Queen-worker ratios in high skew and low skew populations of the ant *Leptothorax acervorum*. (DOCX)

Table S2 Number of inseminated queens and initial queens (in brackets) per replicate colony and treatment. Individual colonies were subjected to different types of stress (food reduction F, worker reduction W, or both FW) or left unmanipulated (control C). (DOCX)

Acknowledgments

We thank Lisa Marie Braun and Martin Bircheneder for assistance during experimental preparation and observation.

Author Contributions

Conceived and designed the experiments: JH JT. Performed the experiments: TS JT. Analyzed the data: JT TS. Wrote the paper: JT JH.

9. Nonacs P, Hager R (2011) The past, present and future of reproductive skew theory and experiments. *Biol Rev* 86: 271–298.
10. Jamieson IG (1997) Testing reproductive skew models in a communally breeding bird, the pukeko, *Porphyrio porphyrio*. *Proc R Soc Lond B* 264: 335–340.
11. Kutsukake N, Nunn CL (2006) Comparative tests of reproductive skew in male primates: the roles of demographic factors and incomplete control. *Behav Ecol Sociobiol* 60: 695–706.
12. Pardi L (1946) Ricerche sui Polistini. VII. La “dominazione” e il ciclo ovarico annuale in *Polistes gallicus* L. *Boll Inst Entomol “Guido Grandi” Univ Stud Bologna* 15: 25–84.
13. Buschinger A (1968) Mono- und Polygynie bei Arten der Gattung *Leptothorax* Mayr (Hymenoptera Formicidae). *Insectes Soc* 15: 217–226.
14. Heinze J, Smith TA (1990) Dominance and fertility in a functionally monogynous ant. *Behav Ecol Sociobiol* 27: 1–10.
15. Yamauchi K, Ishida Y, Hashim R, Heinze J (2007) Queen-queen competition and reproductive skew in a *Cardiocondyla* ant. *Insectes Soc* 54: 268–274.
16. Heinze J, Lipski N, Hölldobler B (1992) Reproductive competition in colonies of the ant *Leptothorax gredleri*. *Ethology* 90: 265–278.
17. Reeve HK (2000) A transactional theory of within-group conflict. *Am Nat* 155: 365–382.
18. Heinze J, Lipski N, Hölldobler B, Bourke AFG (1995) Geographic variation in the social and genetic structure of the ant, *Leptothorax acervorum*. *Zoology* 98: 127–135.
19. Gill RJ, Arce A, Keller L, Hammond RL (2009) Polymorphic social organization in an ant. *Proc R Soc Lond B* 276: 4423–4431.
20. Helms KR, Newman NJ, Helms Cahan S (2013) Regional variation in queen and worker aggression in incipient colonies of the desert ant *Messor pergandei*. *Behav Ecol Sociobiol* 67: 1563–1573.
21. Overson R, Gadau J, Clark RM, Pratt SC, Fewell JH (in press) Behavioral transitions with the evolution of cooperative nest founding by harvester ant queens. *Behav Ecol Sociobiol*: DOI 10.1007/s00265-013-1618-2.
22. Kokko H (2003) Are reproductive skew models evolutionarily stable? *Proc R Soc Lond B* 270: 265–270.
23. Bourke AFG (1991) Queen behaviour, reproduction and egg cannibalism in multiple-queen colonies of the ant *Leptothorax acervorum*. *Anim Behav* 42: 295–310.
24. Stille M, Stille B, Douwes P (1991) Polygyny, relatedness and nest founding in the polygynous myrmicine ant *Leptothorax acervorum*. (Hymenoptera: Formicidae). *Behav Ecol Sociobiol* 28: 91–96.
25. Heinze J, Lipski N, Schlehmeier K, Hölldobler B (1995) Colony structure and reproduction in the ant *Leptothorax acervorum*. *Behav Ecol* 6: 359–367.
26. Hammond RL, Bruford MW, Bourke AFG (2006) A test of reproductive skew models in a field population of a multiple-queen ant. *Behav Ecol Sociobiol* 61: 265–275.
27. Ito F (1990) Functional monogyny of *Leptothorax acervorum* in northern Japan. *Psyche* 97: 203–211.
28. Heinze J, Ortius D (1991) Social organization of *Leptothorax acervorum* from Alaska (Hymenoptera: Formicidae). *Psyche* 98: 227–240.
29. Felke M, Buschinger A (1999) Social organization, reproductive behaviour and ecology of *Leptothorax acervorum* (Hymenoptera, Formicidae) from the Sierra de Albarracin in central Spain. *Insectes Soc* 46: 84–91.
30. Ito F (2005) Mechanisms regulating functional monogyny in a Japanese population of *Leptothorax acervorum* (Hymenoptera, Formicidae): dominance hierarchy and preferential egg cannibalism. *Belg J Zool* 135: 3–8.
31. Trettin J, Haubner M, Buschinger A, Heinze J (2011) Queen dominance and worker policing control reproduction in a threatened ant. *BMC Ecology* 11: 21.
32. Bourke AFG, Heinze J (1994) The ecology of communal breeding: the case of multiply-queened leptothoracine ants. *Phil Trans R Soc Lond B* 345: 359–372.
33. Heinze J (1992) Ecological correlates of functional monogyny and queen dominance in leptothoracine ants. In: Billen J, editor. *Biology and Evolution of Social Insects*. Leuven: Leuven University Press. pp. 25–33.
34. Sommeijer MJ, Van Veen JW (1990) The polygyny of *Myrmica rubra*: selective oophagy and trophallaxis as mechanisms of reproductive dominance. *Entomol Exp Appl* 56: 229–239.
35. Chen YP, Vinson SB (2000) Effects of Queen Attractiveness to Workers on the Queen Nutritional Status and Egg Production in the Polygynous *Solenopsis invicta* (Hymenoptera: Formicidae). *Ann Entomol Soc Am* 93: 295–302.
36. Salomon M, Mayntz D, Lubin Y (2008) Colony nutrition skews reproduction in a social spider. *Behav Ecol* 19: 605–611.
37. Buschinger A (1974) Experimente und Beobachtungen zur Gründung und Entwicklung neuer Sozietäten der sklavenhaltenden Ameise *Harpagoxenus sublaevis* (Nyl.). *Insectes Soc* 21: 381–406.
38. Martin P, Bateson P (2007) *Measuring behaviour: an introductory guide*. Cambridge: Cambridge University Press. 184 p.
39. Buschinger A, Alloway TM (1978) Caste polymorphism in *Harpagoxenus canadensis* M. R. Smith (Hym. Formicidae). *Insectes Soc* 25: 339–350.
40. Buschinger A (1974) Polymorphismus und Kastendetermination im Ameisenstribus Leptothoracini. In: Schmidt GH editor. *Sozialpolymorphismus bei Insekten*. Stuttgart: Wiss. Verlagsges. mbH. pp. 604–623.
41. Sokal RR, Rohlf FJ (1995) *Biometry: the Principles and Practices of Statistics in Biological Research*. New York: W.H. Freeman. 887 p.
42. Holm S (1979) A simple sequentially rejective multiple test procedure. *Scand J Stat* 6: 65–70.
43. Nonacs P (2000) Measuring and using skew in the study of social behavior and evolution. *Am Nat* 156: 577–589.
44. R Development Core Team (2011) R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
45. Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Palaeontological statistics software package for education and data analysis. *Palaeontol Electron* 4: 9.
46. Ortius D, Heinze J (1999) Fertility signaling in queens of a North American ant. *Behav Ecol Sociobiol* 45: 151–159.
47. Franks NR, Scovell E (1983) Dominance and reproductive success among slave-making worker ants. *Nature* 304: 724–725.
48. Bourke AFG (1994) Indiscriminate egg cannibalism and reproductive skew in a multiple-queen ant. *Proc R Soc Lond B* 255: 55–59.
49. Gill RJ, Hammond RL (2011) Workers influence royal reproduction. *Proc R Soc Lond B* 278: 1524–1531.
50. Gotzek D, Ross KG (2007) Genetic regulation of colony social organization in fire ants: an integrative overview. *Q Rev Biol* 82: 201–266.
51. Ruppell O, Schäffler L, Hölldobler B (2002) Lack of plasticity in the behaviour of queens of the ant *Leptothorax rugatulus* Emery (Formicidae: Hymenoptera). *J Insect Behav* 15: 447–454.