



Impact of Serum High Mobility Group Box 1 and Soluble Receptor for Advanced Glycation End-Products on Subclinical Atherosclerosis in Patients with Granulomatosis with Polyangiitis

Alexandre W. S. de Souza^{1,6*}, Karina de Leeuw¹, Mirjan M. van Timmeren², Pieter C. Limburg³, Coen A. Stegeman⁴, Marc Bijl⁵, Johanna Westra¹, Cees G. M. Kallenberg¹

1 Department of Rheumatology and Clinical Immunology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, **2** Department of Pathology and Medical Biology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, **3** Department of Laboratory Medicine, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, **4** Department of Internal Medicine, Division of Nephrology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, **5** Martini Hospital, Groningen, The Netherlands, **6** Rheumatology Division, Universidade Federal de São Paulo/Escola Paulista de Medicina (Unifesp/EPM), São Paulo-SP, Brazil

Abstract

The objective of this study was to evaluate whether levels of high mobility group box 1 (HMGB1) in granulomatosis with polyangiitis (GPA) patients are associated with carotid atherosclerosis, related to levels of soluble receptor for advanced glycation end-products (sRAGE) and influenced by immunosuppressive or lipid-lowering therapy. Twenty-three GPA patients and 20 controls were evaluated for HMGB1- and sRAGE levels and for carotid atherosclerosis using ultrasound to determine intima-media thickness (IMT). *In vitro* the effect of atorvastatin on the production of HMGB1 by lipopolysaccharide (LPS)-stimulated human umbilical vein endothelial cells (HUVEC) was assessed. Serum HMGB1 and sRAGE levels did not differ between patients and controls. A negative correlation was found between sRAGE and maximum IMT but HMGB1 and carotid IMT were not related. HMGB1 levels were reduced in GPA patients on statins and prednisolone. *In vitro*, atorvastatin reduced HMGB1 levels in supernatants of activated HUVEC. In conclusion, carotid IMT is inversely correlated with sRAGE levels but not with HMGB1 levels. Statins and prednisolone are associated with reduced serum HMGB1 levels and atorvastatin decreases HMGB1 release by activated HUVEC *in vitro*, indicating an additional anti-inflammatory effect of statins.

Citation: de Souza AWS, de Leeuw K, van Timmeren MM, Limburg PC, Stegeman CA, et al. (2014) Impact of Serum High Mobility Group Box 1 and Soluble Receptor for Advanced Glycation End-Products on Subclinical Atherosclerosis in Patients with Granulomatosis with Polyangiitis. PLoS ONE 9(4): e96067. doi:10.1371/journal.pone.0096067

Editor: Andreas Zirk, University Heart Center Freiburg, Germany

Received: December 6, 2013; **Accepted:** April 3, 2014; **Published:** April 28, 2014

Copyright: © 2014 de Souza et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was sponsored by the GUIDE Institute and by the Jan Kornelis de Cock Institute. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: alexandre.wagner@unifesp.br

Introduction

High mobility group box 1 (HMGB1) is a nuclear non-histone DNA binding protein that upon cellular death or activation is released into the extracellular milieu and acts as an alarmin. Extracellular HMGB1 stimulates cytokine production, cell proliferation, chemotaxis, angiogenesis, and cell differentiation through binding to its receptors that include the receptor for advanced glycation end-products (RAGE) and Toll-like receptors (TLR)-2, TLR4 and TLR9 [1]. Increased levels of HMGB1 have been found in patients with granulomatosis with polyangiitis (GPA) with active disease, especially in patients with predominantly granulomatous manifestations and in patients without renal involvement at disease onset [2–4]. Individuals with subclinical atherosclerosis and cardiovascular (CV) events present high HMGB1 levels whereas atorvastatin decreases serum HMGB1 in hyperlipidemia [5,6].

RAGE is a multi-ligand surface molecule involved in the pathogenesis of vascular diseases. Soluble RAGE (sRAGE) is a

decoy receptor for RAGE ligands and decreased sRAGE levels are associated with CV events [7]. Patients with GPA present increased advanced glycation end-products (AGEs) accumulation compared to controls and this accumulation is negatively correlated with sRAGE levels [8]. We hypothesized that serum HMGB1 levels contribute to subclinical atherosclerosis in GPA and that this process is influenced by sRAGE but also by treatment with prednisolone and statins.

Materials and Methods

Patients and controls

A cross-sectional study was performed on 23 GPA patients and 20 age- and gender-matched controls (Table 1) enrolled in a previous study to evaluate carotid atherosclerosis in GPA [9]. The study was approved by the Medical Ethical Committee (METC) of the University Medical Center Groningen (UMCG) and written informed consent was obtained from patients and controls. GPA

was classified according to the European Medicines Agency algorithm [10]. The mean disease duration was 131.2 ± 60.2 months. Birmingham Vasculitis Activity Score (BVAS) was recorded for all patients during admissions and out-patient visits allowing assessment of cumulative BVAS scores from the charts. All GPA patients were evaluated during remission (BVAS = 0) to exclude influence of disease activity. Prednisolone was used by 8 patients (34.8%) at a median of 5 mg/day (3.7–10.0 mg). Statins were prescribed for 5 (21.7%) GPA patients. Healthy controls did not use any medication at the time of the study. Carotid ultrasound was performed to assess intima-media thickness (IMT) and carotid plaques as described previously [9,11]. Overall mean IMT (mean of IMT measurements performed on carotid bulb, common and internal carotid arteries) and overall maximum IMT (average of highest IMT values found in above mentioned segments) were used for analysis. Traditional CV risk factors were evaluated according to established guidelines [12]. Serum HMGB1 (Shino Test, Kanagawa, Japan) and sRAGE levels (R&D Systems, Minneapolis, USA) were measured by enzyme-linked immunosorbent assay (ELISA).

Cell cultures

HUVEC (Lonza, Breda, The Netherlands) were cultured in EBM-2 medium supplemented with EGM-2 MV Single Quot Kit Supplements & Growth Factors (cat No. CC-3202, Lonza) and used when confluent. Three groups were evaluated: (1) HUVEC pre-incubated for 2 hours with 5 μ M atorvastatin (Sigma Aldrich, Saint Louis, USA) and treated with LPS (100 ng/ml) (Sigma Aldrich, Saint Louis, USA), (2) HUVEC treated only with LPS, and (3) unstimulated HUVEC. Supernatants were collected for measuring HMGB1 and interleukin (IL)-8 at baseline, 4 hours and 24 hours. All *in vitro* experiments were performed twice and in duplicate. Cell viability of HUVEC was checked with 0.2% trypan

blue dye (Invitrogen, Carlsbad, USA) on HUVEC treated with 0 μ M, 0.1 μ M, 1.0 μ M, 5.0 μ M and 10.0 μ M atorvastatin; percentages of living cells were 94%, 95%, 95%, 92% and 91% at 36 hours, respectively. Western blot was used to measure HMGB1 in supernatants as previously described [13] while IL-8 levels were measured by ELISA (R&D Systems, Minneapolis, USA).

Statistical analysis

Statistical analysis was performed with SPSS 18.0 software and graphs were built using GraphPad Prism 5. Continuous variables are presented as mean \pm SD or as median and interquartile range. Categorical variables are presented as total number and percentage. Comparisons between groups were performed using chi-square test or Fisher's exact test and Student's t-test or Mann-Whitney U test as appropriate. Analysis of longitudinal data from *in vitro* experiments was performed by two-way ANOVA and Bonferroni's test. Correlations were evaluated with Spearman's rank correlation coefficient. Differences were considered significant when $p < 0.05$.

Results

HMGB1, sRAGE and subclinical atherosclerosis

Figure 1 shows similar serum HMGB1 levels in patients and controls whereas Figures 2A and 2B depicts similar overall mean IMT and maximum IMT in carotid arteries in GPA and controls. Furthermore, sRAGE levels and the prevalence of carotid plaques did not differ between patients and controls (Table 1). In GPA patients, no correlation was found between HMGB1 and overall maximum IMT in carotid arteries ($\rho = 0.062$; $p = 0.820$). However, sRAGE levels were negatively correlated with overall maximum IMT in carotid arteries ($\rho = -0.565$; $p = 0.035$). No

Table 1. Risk factors for cardiovascular disease, carotid ultrasound, HMGB1 and sRAGE levels in GPA patients and controls.

| Variables | GPA patients (N = 23) | Controls (N = 20) | <i>p</i> |
|---|-----------------------|---------------------|----------|
| Median age at study, years | 55.2 (45.7–62.4) | 49.8 (43.0–57.4) | 0.173 |
| Females, n (%) | 9 (39.1) | 9 (45.0) | 0.697 |
| Age >45 years for men and 55 years for women, n (%) | 16 (69.6) | 12 (60.0) | 0.512 |
| Mean total cholesterol, mmol/L | 4.99 \pm 0.78 | 4.98 \pm 0.82 | 0.978 |
| Mean HDL, mmol/L | 1.41 \pm 0.37 | 1.51 \pm 0.33 | 0.359 |
| Mean LDL, mmol/L | 3.01 \pm 0.79 | 3.29 \pm 0.82 | 0.267 |
| Median TGL, mmol/L | 1.50 (1.00–2.00) | 1.00 (1.00–2.00) | 0.404 |
| Mean systolic BP, mmHg | 123.65 \pm 14.55 | 119.24 \pm 10.94 | 0.301 |
| Mean diastolic BP, mmHg | 70.61 \pm 9.38 | 75.12 \pm 8.07 | 0.120 |
| Smoking, n(%) | 2 (8.7) | 1 (5.0) | 0.635 |
| Family history of premature CVD, n(%) | 9 (39.1) | 8 (40.0) | 0.954 |
| Median BMI, kg/m ² | 26.0 (24.0–28.0) | 23.5 (22.0–26.5) | 0.256 |
| Previous CVD, n(%) | 3 (13.0) | 0 (0.0) | 0.236 |
| Carotid plaques, n (%) | 7 (30.4) | 3 (15.0) | 0.203 |
| Overall mean IMT, mm | 0.833 \pm 0.256 | 0.765 \pm 0.133 | 0.357 |
| Median overall maximum IMT, mm | 0.875 (0.810–1.215) | 0.880 (0.830–0.990) | 0.953 |
| HMGB1, ng/ml | 2.13 (1.53–4.15) | 2.42 (1.73–4.07) | 0.827 |
| sRAGE, pg/mL | 1256.1 \pm 559.6 | 1483.3 \pm 399.8 | 0.155 |

Numerical data are presented as mean \pm standard deviation or as median and interquartile range; BMI: body mass index; BP: blood pressure; CVD: cardiovascular disease; HDL: high-density lipoprotein; HMGB1: high mobility group box-1; IMT: intima-media thickness; LDL: low-density lipoprotein; n: number of individuals; sRAGE: soluble receptor for advanced glycation end-products; TGL: triglycerides.

doi:10.1371/journal.pone.0096067.t001




Figure 1. Serum HMGB1 levels in GPA patients and controls. GPA patients in remission present similar median HMGB1 levels compared to controls. doi:10.1371/journal.pone.0096067.g001

correlation was present between HMGB1 and sRAGE ($\rho = 0.068$; $p = 0.777$), between overall mean carotid IMT and cumulative BVAS ($\rho = -0.070$; $p = 0.805$) or between overall mean carotid IMT and time in remission since last relapse/

presentation ($\rho = 0.337$; $p = 0.201$). Moreover, no correlation was found between overall maximum carotid IMT and cumulative BVAS ($\rho = -0.050$; $p = 0.859$) and between overall maximum IMT and time in remission since last relapse/presentation ($\rho = 0.338$; $p = 0.200$). Also, no significant difference was found between patients with and without carotid plaques regarding HMGB1 [$1.52 (1.20-2.89)$ ng/ml vs. $2.48 (1.74-4.22)$ ng/ml; $p = 0.300$] and sRAGE levels (1144.66 ± 817.33 pg/ml vs. 1303.99 ± 438.86 pg/ml; $p = 0.574$).

Impact of treatment on HMGB1 and sRAGE

Since HMGB1 and sRAGE levels did not differ between GPA patients in remission and controls, we evaluated whether therapy would have any influence on both biomarkers. Serum HMGB1 levels were lower in patients on statins [$1.26 (1.16-1.68)$ ng/ml vs. $2.70 (1.75-4.76)$ ng/ml; $p = 0.014$] and on prednisolone [$1.49 (1.21-2.61)$ ng/ml vs. $2.51 (1.79-5.61)$ ng/ml; $p = 0.017$] compared to patients without these drugs, respectively (Figure 3A and 3B). Although not significant, we observed higher serum HMGB1 levels in 12 GPA patients without statins or prednisolone compared with 20 controls (4.03 ± 2.00 ng/ml vs. 2.84 ± 1.58 ng/ml, $p = 0.073$). sRAGE levels did not differ between patients with and without statins (976.20 ± 345.64 pg/ml vs. 1326.19 ± 588.79 pg/ml; $p = 0.275$) or prednisolone (1366.96 ± 450.80 pg/ml vs. 1196.55 ± 619.08 pg/ml; $p = 0.531$).




Figure 2. Intima media thickness in common carotid arteries in GPA and controls. GPA patients present mean IMT (2A) and maximum IMT (2B) in common carotid arteries similar to controls. doi:10.1371/journal.pone.0096067.g002




Figure 3. HMGB1 levels in GPA patients treated with statins or prednisolone. median serum levels of HMGB1 were significantly lower in GPA patients treated with statins (3A) and prednisolone (3B) when compared to patients without these drugs. doi:10.1371/journal.pone.0096067.g003




Figure 4. Effect of atorvastatin on HMGB1 and IL-8 levels in HUVEC supernatants. 4A – A time curve was built to evaluate HMGB1 release from HUVEC upon LPS stimulation. HMGB1 intensity in supernatants increases 8 hours after stimulation and a peak is observed at 24 hours with a decrease at 48 hours. 2B and C – HMGB1 and IL-8 levels are significantly higher at 24 hours in supernatants from HUVEC treated with LPS in comparison to unstimulated HUVEC while pre-incubation with 5 µM atorvastatin followed by LPS stimulation lowered HMGB1 and IL-8 levels significantly at 24 hours in HUVEC's supernatants. Experiments were performed twice and in duplicate. Data are presented as median and range, * $p < 0.01$ and ** $p < 0.001$. doi:10.1371/journal.pone.0096067.g004

HMGB1 and atorvastatin in HUVEC

To investigate the relation between statins and HMGB1, we tested the effect of atorvastatin on HMGB1-release from LPS-stimulated HUVEC. HMGB1 in supernatants gradually increased in time with a peak after 24 hours of LPS stimulation and a decrease at 48 hours (Figure 4A). Thus, we compared HMGB1

and IL-8 levels in supernatants of HUVEC pre-incubated with atorvastatin followed by 24 hours of LPS stimulation. HUVEC stimulated with LPS showed a significantly higher production of HMGB1 and IL-8 than unstimulated cells. Pre-incubation with 5 µM atorvastatin prior to LPS stimulation led to significantly lower levels of both HMGB1 and IL-8 in supernatants after

24 hours in comparison to HUVEC stimulated by LPS only (Figures 4B and 4C).

Discussion

In this study, in GPA patients HMGB1 levels were not correlated with overall carotid maximum IMT, whereas sRAGE levels were negatively correlated with IMT. Furthermore, statins or prednisolone use was associated with lower HMGB1 levels.

In previous studies, GPA patients presented higher IMT levels in carotid arteries and an increased number of CV events compared to controls [11,14]. Accelerated atherosclerosis in GPA was not associated with traditional CV risk factors, rather enhanced levels of markers of vascular inflammation and remodeling were associated with atherosclerotic disease [11]. In this study, we evaluated HMGB1 and sRAGE levels in GPA patients who had been enrolled in a follow-up study to evaluate progression of atherosclerosis [9]. Patients were in remission and a reduction of traditional CV risk factors was achieved during follow-up of these patients that possibly resulted in similar IMT and similar prevalence of carotid plaques compared to controls [9]. Also, cumulative BVAS scores and time in remission were not related to IMT values. Moreover, proteinase 3 antineutrophil cytoplasmic antibody (PR3-ANCA) status has been shown to have a protective role on the risk of CV events in patients with ANCA-associated vasculitis and most of our patients were PR3-ANCA positive [15].

HMGB1 and sRAGE have been implicated in the pathogenesis of atherosclerotic disease. Higher serum HMGB1 levels are a potential marker of subclinical atherosclerosis and CV events while lower serum sRAGE levels are associated with risk factors for CV disease and CV events [5,7]. Thus, we evaluated the association between both biomarkers and atherosclerotic disease in GPA. sRAGE was associated with subclinical carotid atherosclerosis whereas HMGB1 was not. We hypothesized that due to its multifaceted nature, HMGB1 is possibly more influenced by other factors such as therapy while sRAGE levels are more stable. Indeed, prednisolone or statins use were associated with lower serum HMGB1 levels in our study.

Previous studies had also shown reduction of HMGB1 levels upon statins in an experimental model of atherosclerosis and in humans with hyperlipidemia [6,16]. In this study, we demonstrated that statins are associated with lower serum HMGB1 levels in GPA patients in remission while a tendency for higher serum HMGB1 levels was observed in GPA patients in remission without

statins compared with controls. Moreover, the addition of 5 μ M atorvastatin which is within the range of atorvastatin concentration used in other *in vitro* studies [17–21] led to a decrease in extracellular HMGB1 levels in activated HUVEC. These findings indicate that endothelial cells are a possible source of extracellular HMGB1 that could be inhibited by atorvastatin. The decrease in IL-8 levels in supernatants of activated HUVEC by atorvastatin was used as a comparator since it is already known that statins inhibit mRNA expression of IL-8 in activated HUVEC [22]. Inhibition of HMGB1 release by activated HUVEC points to another potential anti-inflammatory effect of statins on vascular endothelium. The actual mechanism of inhibition of HMGB1 release by HUVEC (i.e. inhibition of mRNA expression or cytoplasmic translocation of HMGB1) still needs further elucidation.

HMGB1 levels were also lower in GPA patients on prednisolone therapy. Previous studies have demonstrated that corticosteroids can reduce extracellular release of HMGB1 by monocytes *in vitro* and they can also reduce HMGB1 expression and circulating levels *in vivo* [23,24]. However, these findings were not confirmed in patients with rheumatoid arthritis [25].

In conclusion, no association was observed between subclinical carotid atherosclerosis and HMGB1 while sRAGE was negatively associated with carotid IMT in GPA. Statin use was associated with lower HMGB1 levels suggesting an additional anti-inflammatory property of statins. As subclinical atherosclerosis was similar between patients and controls, we suggest that development of premature atherosclerosis in GPA patients might be postponed by sRAGE and use of statins or prednisolone. Since our study had a relatively low number of patients and had a cross-sectional design, further longitudinal studies are needed to evaluate if reduction of serum HMGB1 levels might be important in CV risk management in GPA.

Acknowledgments

The authors thank, Hans L. A. Nienhuis, Andries J Smit, Paulina Sosicka for their contribution to this study. The authors also thank Henk Moorlag from the Endothelial Cell Facility (UMCG, Groningen) for the HUVEC.

Author Contributions

Conceived and designed the experiments: JW PCL MB CGMK. Performed the experiments: AWSS MMvT JW. Analyzed the data: AWSS JW PCL MB CAS CGMK. Contributed reagents/materials/analysis tools: KL JW MMvT. Wrote the paper: AWSS.

References

- Harris HE, Andersson U, Pisetsky DS (2012) HMGB1: a multifunctional alarmin driving autoimmune and inflammatory disease. *Nat Rev Rheumatol* 8: 195–202.
- Wibisono D, Csernok E, Lamprecht P, Holle JU, Gross WL, et al. (2010) Serum HMGB1 levels are increased in active Wegener's granulomatosis and differentiate between active forms of ANCA-associated vasculitis. *Ann Rheum Dis* 69: 1888–1889.
- Henes FO, Chen Y, Bley TA, Fabel M, Both M, et al. (2011) Correlation of serum level of high mobility group box 1 with the burden of granulomatous inflammation in granulomatosis with polyangiitis (Wegener's). *Ann Rheum Dis* 70: 1926–1929.
- De Souza AW, Westra J, Bijzet J, Limburg PC, Stegeman CA, et al. (2013) Is serum HMGB1 a biomarker in ANCA-associated vasculitis? *Arthritis Res Ther* 15: R104.
- De Souza AW, Westra J, Limburg PC, Bijl M, Kallenberg CG (2012) HMGB1 in vascular diseases: its role in vascular inflammation and atherosclerosis. *Autoimmun Rev* 11: 909–917.
- Jin D, Wu Y, Zhao L, Guo J, Zhang K, et al. (2012) Atorvastatin reduces serum HMGB1 levels in patients with hyperlipidemia. *Exp Ther Med* 4: 1124–1126.
- Kalea AZ, Schmidt AM, Hudson BI (2009) RAGE: a novel biological and genetic marker for vascular disease. *Clin Sci (Lond)* 116: 621–637.
- de Leeuw K, Nienhuis H, Smit A, Stegeman C, Kallenberg C, et al. (2010) Increased accumulation of advanced glycation endproducts in patients with Wegener's granulomatosis. *Ann Rheum Dis* 6: 625–627.
- de Leeuw K, van der Graaf AM, Bijzet J, Stegeman CA, Smit AJ, et al. (2010) Patients with Wegener's granulomatosis: a long-term follow-up study. *Clin Exp Rheumatol (Suppl. 57)*:S18–23.
- Watts R, Lane S, Hanslik T, Hauser T, Hellmich B, et al. (2007) Development and validation of a consensus methodology for the classification of the ANCA-associated vasculitides and polyarteritis nodosa for epidemiological studies. *Ann Rheum Dis* 66: 222–227.
- de Leeuw K, Sanders JS, Stegeman C, Smit A, Kallenberg CG, et al. (2005) Accelerated atherosclerosis in patients with Wegener's granulomatosis. *Ann Rheum Dis* 64: 753–759.
- Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol in Adults. Executive summary of the third report of the national cholesterol education program (adult treatment panel III) (2001) *JAMA* 285: 2486–2497.
- Abdulhadi DA, Westra J, Bijzet J, Limburg PC, Kallenberg CG, et al. (2011) High mobility group box 1 (HMGB1) and anti-HMGB1 antibodies and their relation to disease characteristics in systemic lupus erythematosus. *Arthritis Res Ther* 13: R71.

14. Faurschou M, Mellemkjaer L, Sorensen IJ, Svalgaard Thomsen B, Dreyer L, et al. (2009) Increased morbidity from ischemic heart disease in patients with Wegener's granulomatosis. *Arthritis Rheum* 60: 1187–1192.
15. Suppiah R, Judge A, Batra R, Flossmann O, Harper L, et al. (2011) A model to predict cardiovascular events in patients with newly diagnosed Wegener's granulomatosis and microscopic polyangiitis. *Arthritis Care Res* 63: 588–596.
16. Haraba R, Suica VI, Uyy E, Ivan L, Antohe F (2011) Hyperlipidemia stimulates the extracellular release of the nuclear high mobility group box 1 protein. *Cell Tissue Res* 346: 361–368.
17. Hol J, Otterdal K, Breland UM, Stang E, Pedersen TM, et al. (2012) Statins affect the presentation of endothelial chemokines by targeting to multivesicular bodies. *PLoS One* 7:e40673.
18. Jia F, Wu C, Chen Z, Lu G (2012) Atorvastatin inhibits homocysteine-induced endoplasmic reticulum stress through activation of AMP-activated protein kinase. *Cardiovasc Ther* 30:317–25.
19. Izidoro-Toledo TC, Guimaraes DA, Belo VA, Gerlach RF, Tanus-Santos JE (2011) Effects of statins on matrix metalloproteinases and their endogenous inhibitors in human endothelial cells. *Naunyn Schmiedeberg Arch Pharmacol* 383:547–54.
20. Dulak J, Loboda A, Jazwa A, Zagorska A, Dörler J, et al. (2005) Atorvastatin affects several angiogenic mediators in human endothelial cells. *Endothelium* 12:233–41.
21. Bao XM, Wu CF, Lu GP (2009) Atorvastatin attenuates homocysteine-induced apoptosis in human umbilical vein endothelial cells via inhibiting NADPH oxidase-related oxidative stress-triggered p38MAPK signaling. *Acta Pharmacol Sin* 30:1392–8.
22. Hot A, Lavocat F, Lenief V, Miossec P (2013) Simvastatin inhibits the pro-inflammatory and pro-thrombotic effects of IL-17 and TNF- α on endothelial cells. *Ann Rheum Dis* 72: 754–760.
23. Schierbeck H, Wahämaa H, Andersson U, Harris HE (2010) Immunomodulatory drugs regulate HMGB1 release from activated human monocytes. *Mol Med* 16: 343–351.
24. Huang YH, Wang PW, Tiao MM, Chou MH, Du YY, et al. (2011) Glucocorticoid modulates high-mobility group box 1 expression and Toll-like receptor activation in obstructive jaundice. *J Surg Res* 170: e47–55.
25. Pullerits R, Urbonaviciute V, Voll RE, Forsblad-D'Elia H, Carlsten H (2011) Serum levels of HMGB1 in postmenopausal patients with rheumatoid arthritis: associations with proinflammatory cytokines, acute-phase reactants, and clinical disease characteristics. *J Rheumatol* 38: 1523–1525.