



The Relation of Hepcidin to Iron Disorders, Inflammation and Hemoglobin in Chronic Kidney Disease

Lucile Mercadel^{1,2*}, Marie Metzger^{1,3}, Jean Philippe Haymann⁴, Eric Thervet⁵, Jean-Jacques Boffa⁶, Martin Flamant^{7,8}, François Vrtovsnik^{8,9}, Pascal Houillier¹⁰, Marc Froissart¹ and Bénédicte Stengel^{1,3} on behalf of the NephroTest Study Group[†]

1 Inserm, Centre for research in Epidemiology and Population Health, U1018, Epidemiology of Diabetes, Obesity, and Kidney Diseases Team, Villejuif, France, **2** Department of Nephrology, Hôpital Pitié-Salpêtrière, Assistance Publique-Hôpitaux de Paris, Paris, France, **3** Université Paris Sud 11, U1018, Villejuif, France, **4** Department of Physiology and Nephrology, Hôpital Tenon, Assistance Publique-Hôpitaux de Paris, Université Pierre et Marie Curie, Inserm U702, Paris, France, **5** Department of Nephrology, Hôpital Européen G Pompidou, Assistance Publique-hôpitaux de Paris, Paris, France, **6** Department of Nephrology, Hôpital Tenon, Assistance Publique-hôpitaux de Paris, université Pierre et Marie Curie, Paris, France, **7** Department of Physiology, Hôpital Bichat, Assistance Publique-hôpitaux de Paris, Paris, France, **8** Inserm U699, Paris, France, **9** Department of Nephrology, Hôpital Bichat, Assistance Publique-hôpitaux de Paris, Université Pierre et Marie Curie, Paris, France, **10** Department of Physiology, Hôpital Européen G Pompidou, Assistance Publique-hôpitaux de Paris, Université Paris Descartes, Paris, France

Abstract

The metabolism of hepcidin is profoundly modified in chronic kidney disease (CKD). We investigated its relation to iron disorders, inflammation and hemoglobin (Hb) level in 199 non-dialyzed, non-transplanted patients with CKD stages 1–5. All had their glomerular filtration rate measured by ⁵¹Cr-EDTA renal clearance (mGFR), as well as measurements of iron markers including hepcidin and of erythropoietin (EPO). Hepcidin varied from 0.2 to 193 ng/mL. The median increased from 23.3 ng/mL [8.8–28.7] to 36.1 ng/mL [14.1–92.3] when mGFR decreased from ≥ 60 to < 15 mL/min/1.73 m² ($p = 0.02$). Patients with absolute iron deficiency (transferrin saturation (TSAT) $< 20\%$ and ferritin < 40 ng/mL) had the lowest hepcidin levels (5.0 ng/mL [0.7–11.7]), and those with a normal iron profile (TSAT $\geq 20\%$ and ferritin ≥ 40), the highest (34.5 ng/mL [23.7–51.6]). In multivariate analysis, absolute iron deficiency was associated with lower hepcidin values, and inflammation combined with a normal or functional iron profile with higher values, independent of other determinants of hepcidin concentration, including EPO, mGFR, and albuminemia. The hepcidin level, although it rose overall when mGFR declined, collapsed in patients with absolute iron deficiency. There was a significant interaction with iron status in the association between Hb and hepcidin. Except in absolute iron deficiency, hepcidin's negative association with Hb level indicates that it is not down-regulated in CKD anemia.

Citation: Mercadel L, Metzger M, Haymann JP, Thervet E, Boffa J-J, et al. (2014) The Relation of Hepcidin to Iron Disorders, Inflammation and Hemoglobin in Chronic Kidney Disease. PLoS ONE 9(6): e99781. doi:10.1371/journal.pone.0099781

Editor: Jaap A. Joles, University Medical Center Utrecht, Netherlands

Received: February 5, 2014; **Accepted:** May 16, 2014; **Published:** June 30, 2014

Copyright: © 2014 Mercadel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The NephroTest CKD cohort study is supported by grants from: Inserm GIS-IReSP AO 8113LS TGIR (BS); French Ministry of Health AOM 09114 (MFr); Inserm AO 8022LS (BS); Agence de la Biomédecine R0 8156LL (BS), AURA (MFr), and Roche 2009-152-447G (MFr). The Nephrotest initiative was also sponsored by unrestricted grants from F. Hoffman-La Roche Ltd. (LM). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: This study was partly funded by La Roche Ltd. M. Froissart has received consulting or lecture fees or research funds from Affymax, Genzyme, Hoffmann-La Roche, Novartis, Sandoz, Shire, Takeda, and Vifor International. M. Froissart has been employed by Amgen since January 1, 2011, but was a full-time academic associate professor during the time of study conception and data collection. B.S. has received research funds from Amgen, Baxter, Genzyme (Sanofi), Fresenius, MSD, and GSK. Dr Barbara Sasu and Vimal Patel provided advised and performed hepcidin measurements at Amgen laboratories (Amgen Inc., Thousand Oaks, CA). In order to enable hepcidin measurements, this study was labelled as an Investigator Sponsored Study (BS) by Amgen. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* Email: lucile.mercadel@psl.aphp.fr

† Membership of the NephroTest Study Group is provided in the Acknowledgments.

Introduction

Hepcidin is a new iron marker, discovered in 2001 and studied especially in hemochromatosis. The first investigations concerned prohepcidin, an inactive precursor. Prohepcidin has no impact on iron metabolism in either healthy individuals [1] or patients with chronic kidney disease (CKD) [2], [3]. The involvement of hepcidin in iron disorders was clarified with the development of methods to quantify hepcidin-25 [2], [4]. Hepcidin has been evaluated in absolute iron deficiency, where its concentration is low, and in patients with chronic disease-related anemia [5]. The subsequent development of more specific assays for hepcidin-25 has led to a redefinition of normal values from around 50 ng/mL

for the first assays to less than 10 ng/mL for those recently validated by the FDA [6]. Except in hemochromatosis, however, hepcidin has not yet proved its usefulness in clinical practice.

Hepcidin involvement in CKD anemia has been explored in both dialysis and non-dialysis CKD patients: both CKD-related inflammation and lower hepcidin clearance tend to increase hepcidin concentration in these patients [7], [8], [9]. The specificity of hepcidin for the diagnosis of absolute iron deficiency may thus be limited in this group. The hepcidin level does not seem to improve the prediction of erythropoietin response [10], [11]. Furthermore, its close relation with ferritin raises questions about its performance in the diagnosis of iron disturbances [12]. In

hemodialysis patients, its performance is no better than that of traditional iron markers [2], [12]. Nonetheless, a better understanding of the determinants of hepcidin levels is essential to assess its usefulness in CKD anemia. We therefore studied the determinants of hepcidin in non-dialysis CKD patients, particularly its relation to iron disorders, inflammation, and glomerular filtration rate measured (mGFR) by a reference method. We also tested the hypothesis that hepcidin and hemoglobin (Hb) levels are related, independently of other Hb determinants.

Methods

Population

The NephroTest study is a prospective hospital-based cohort that has enrolled patients with any diagnosis of CKD stages 1 through 5, recruited from three nephrology departments. Patients younger than 18 years, dialysis patient or with a kidney transplant, and pregnant women were excluded. Between January 2000 and January 2012, NephroTest included 1095 patients after they had provided written informed consent. The NephroTest study design was approved by the relevant ethics committee (Direction Générale pour la Recherche et l'Information, Comité Consultatif sur le Traitement de l'Information en matière de Recherche dans le domaine de la Santé MG/CP09.503) and adheres to the Declaration of Helsinki. Patients not receiving EPO and intravenous iron were included in this analysis if a plasma sample was available for them. More precisely, 114 men were drawn from the NEPHROTEST cohort samples from 2000 to 2004 and all women of the same period were included ($n = 48$). Additionally, 37 women with plasma samples available from the 2006 to 2008 period were added to obtain a study population that was balanced with respect to sex. Hepcidin was measured for these 199 patients. They were similar to the overall study population with respect to age and mGFR distribution (Appendix 1). Biopsy-proven nephropathy was identified in 21% of all patients, but fewer than 10% of those with diabetes. Clinical criteria based on a history of urinary albumin >300 mg/g creatinine and of other microangiopathy damage (retinopathy or neuropathy) were used to classify diabetes patients who had not had renal biopsies with diabetic glomerular nephropathy. Other diabetes patients were classified with another nephropathy type, most probably vascular.

Laboratory measurements

Patients undergo an extensive annual check-up, including a measurement of their GFR (mGFR) by $^{51}\text{Cr-EDTA}$, in the hospitals' physiology departments [13]. Serum iron (DxC800 Beckman-Coulter, ferrozine, emitted light 560 nm), ferritin (BN-Siemens, N-latex ferritin immunonephelometry), and transferrin (BN-Siemens, N Antiserum antitransferrin immunonephelometry) were measured. TIBC (total iron binding capacity, $\mu\text{mol/L}$) was calculated as $25 \times \text{transferrin (g/L)}$ and TSAT (%) as $\text{serum iron} \times 100 / \text{TIBC}$. Firstly, we used Lipschitz's iron index [13] to define iron status based on TSAT and ferritin together as follows: normal, $\text{TSAT} \geq 20\%$; absolute iron deficiency, $\text{TSAT} < 20\%$ and ferritin < 40 ng/mL; functional iron deficiency, $\text{TSAT} < 20\%$ and ferritin ≥ 40 ng/mL. The 40 ng/mL threshold value was chosen for ferritin because this threshold is usually recommended to diagnose absolute iron deficiency, in contrast with the KDIGO threshold of 100 ng/mL used to define the need for iron supply in patients with non-dialysis chronic kidney disease. Secondly, we used a combined iron marker containing 4 classes, built like the Lipschitz iron index but separating $\text{TSAT} \geq 20\%$ and ferritin ≥ 40 ng/mL from $\text{TSAT} \geq 20\%$ and ferritin < 40 ng/mL. Endogenous EPO levels were determined in serum (100 μL) with the

Quantitine IVD Epo double-antibody sandwich ELISA method from R&D Systems (Minneapolis, MN), as reported elsewhere [14]. EPO measurements were missing for 38 patients (37 women and 1 men).

The assay for the quantification of hepcidin in human serum was developed at Amgen, Thousand Oaks, CA, and is described in appendix 2. Briefly, this immunoassay is based on hepcidin capture by an anti-human hepcidin monoclonal antibody, followed by electrochemiluminescent detection of the complex. The normal reference level of hepcidin in healthy volunteers was < 10 ng/mL. The lower limit of detection was 0.1 ng/mL.

Statistical analyses

We studied crude and mGFR-adjusted relations of hepcidin with age, gender, ethnicity, diabetes and diabetic nephropathy, body mass index (BMI), mGFR, the urinary protein to creatinine ratio (PCR), albuminemia, C-reactive protein (CRP), oral iron treatment, erythropoietin, ferritin, TSAT, TIBC, and the combined iron marker; we used ANOVA to compare categorical variables and Pearson's correlations for quantitative variables. The hepcidin determinants were then analysed by multivariate regression analysis that included age, gender, center, albumin, BMI, CRP, mGFR, EPO, oral iron therapy, and the combined iron marker. Finally, we tested the association of Hb levels with the hepcidin concentration treated continuously after adjustment for other Hb determinants. These included the combined iron marker (or ferritin), gender, diabetes, BMI, mGFR, oral iron treatment, albuminemia, CRP, and angiotensin converting enzyme inhibitors/angiotensin receptor blockers. Potential interactions between hepcidin and the combined iron marker (or ferritin) in the relation with Hb were also tested. In all analyses, hepcidin was transformed by its square root to meet the criterion of a normal parameter.

For covariates with less than 3% missing observations (albumin and CRP), the median value was imputed in the multivariate analysis. A missing data category was created for the combined iron marker and EPO. Statistical analyses were performed with SAS 9.2 (SAS Institute Inc., Cary, NC, USA) and R 2.15 (R Foundation for Statistical Computing, Vienna, Austria, 2012).

Results

Patients' characteristics

Patients' characteristics are shown in Table 1. Hepcidin ranged from 0.2 to 193 ng/mL with a median value of 27.9 ng/mL [IQR 16.5–45.4]. According to Lipschitz's iron index [15], 72.4% had normal iron profiles, 6.3% absolute iron deficiency and 21.4% functional iron deficiency. This distribution was similar to that observed in the entire cohort (71.3%, 6.2%, and 18.6%). The median mGFR for each class of the combined iron marker was 34.7 mL/min/1.73 m² [IQR 24.0–48.3], 37.2 mL/min/1.73 m² [24.8–56.0], and 36.0 mL/min/1.73 m² [26.3–49.5].

Factors associated with hepcidin levels

Women younger than 55 years had lower hepcidin values than either men or older women (Table 2). Hepcidin levels increased from 23.3 ng/mL IQR [8.8–28.7] to 36.1 ng/mL IQR [14.1–92.3] when mGFR declined from >60 mL/min/1.73 m² to < 15 mL/min/1.73 m². Independent of mGFR, higher levels of body mass index, albuminemia, C-reactive protein (CRP), oral iron therapy and lower levels of proteinuria and EPO were significantly associated with higher hepcidin concentrations (Tables 2 and 3). Hepcidin levels were strongly related to all iron markers and particularly to ferritin (Table 3 and Figure 1). The hepcidin level varied in the different combined classes of TSAT

Table 1. Patient characteristics.

	Mean ± SD, median (IQR) or %(N)
Men	57.3 (114)
Age, years	58.4±14.8
African origin	5.8 (11)
Body mass index, kg/m ²	25.5±5.0
Systolic/Diastolic Blood Pressure, mmHg	138±20/76±11
Diabetes	
No	76.4 (152)
Yes, with diabetic nephropathy	15.1 (30)
Yes, with other nephropathy type	8.5 (17)
History of cardiovascular disease	12.6 (25)
mGFR, ml/min/1.73 m ²	35.3 (24.2–49.3)
<15	7.0 (14)
15–30	28.1 (56)
30–45	33.7 (67)
45–60	18.1 (36)
>60	13.1 (26)
eGFR MDRD, ml/min/1.73 m ²	33.3 (22.6–45.9)
eGFR CKD-EPI, ml/min/1.73 m ²	34.2 (23.2–48.4)
PCR, mg/mmol	28.9 (15.3–115.1)
Serum albumin, g/L	39.53±5.18
C-reactive protein >8 mg/L	11.0 (21)
Hb, g/dL	12.32±1.47
WHO Anemia	54.8 (109)
Serum Iron, µmol/L	14.06±5.16
Total iron binding capacity (TIBC), µmol/L	56.5±10.9
Transferrin saturation (TSAT), %	25.47±9.83
Ferritin, µg/L	110 (63–201)
Folates, ng/L	6.8 (5.0–9.5)
Vitamin B12, pg/mL	418 (291–558)
ACEi or ARB %	68.3 (136)
Oral iron therapy	11.1 (22)
Lipschitz's iron index	
TSAT ≥20%	72.4 (139)
TSAT <20% and ferritin <40	6.3 (12)
TSAT <20% and ferritin ≥40	21.4 (41)

PCR: urinary protein to creatinine ratio; ACEi/ARBs: angiotensin converting enzyme inhibitors/angiotensin receptor blockers. WHO anemia: Hb <13 g/dL for men and < 12 g/dL for women.
doi:10.1371/journal.pone.0099781.t001

and ferritin (Table 4). It was six times lower in patients with absolute iron deficiency and three times in those with low ferritin alone than in those with normal iron status. The multivariate analysis showed that oral iron therapy, albuminemia, EPO levels, BMI ≥ 30 kg/m² and mGFR all remained significantly associated with hepcidin levels. The four-class iron index produced the model with the best fit (Table 5). There was an interaction between iron status profile and CRP in their relation with hepcidin (p = 0.01). Patients with inflammation and either a functional or normal iron profile had higher hepcidin levels than their counterparts without inflammation (Table 5). Patients with ferritin <40 ng/mL had an hepcidin collapse.

Relation between Hb and hepcidin levels

There was a significant interaction with the four-class iron index in the relation between hepcidin and Hb (p = 0.003). Hb levels tended to decrease as hepcidin levels increased when TSAT < 20% and ferritin >40 µg/L (Hb = -0.14±0.08, p = 0.06), but the relation was reversed in patients with absolute iron deficiency, defined by TSAT <20% and ferritin <40 µg/L (Hb = 0.30±0.14, p = 0.04). These relations were not modified when the four-class iron index was combined with inflammation (data not shown). Using ferritin instead of the combined iron marker produced similar results for the relation between hepcidin and Hb, with significant interaction between tertiles of ferritin and hepcidin (p < 0.001). Hb levels decreased significantly with increasing hepcidin

Table 2. Hepcidin level according to categorial factors.

		N	Hepcidin (ng/mL) Median (IQR)	p-value	mGFR adjusted p-value
Age×Gender	Men	114	27.7 (19.0–43.1)	0.009	0.003
	Women <55 yrs	35	15.6 (8.8–40.2)		
	Women ≥55 yrs	50	34.6 (21.3–57.4)		
Origin	African	11	34.5 (8.0–63.9)	0.6	0.5
	Caucasian or other	180	27.7 (16.6–43.6)		
	missing	8	47.6 (16.2–66.0)		
Diabetes	No	152	28.8 (17.7–46.1)	0.8	0.8
	Yes with diabetic nephropathy	30	26.2 (12.5–45.2)		
	Yes with other nephropathy type	17	21.4 (16.3–42.4)		
Oral Iron Therapy	No	177	27.1 (16.0–42.3)	0.005	0.01
	Yes	22	46.5 (19.0–75.9)		
ACEI/ARBs	No	63	26.5 (16.0–48.3)	0.8	0.8
	Yes	136	29.3 (17.2–44.7)		
CRP	≤8 mg/L	170	26.7 (16.4–40.4)	0.0002	0.0005
	>8	21	56.4 (26.1–73.2)		
	missing	8	23.8 (7.7–87.7)		

CRP: C reactive protein; ACE/ARBs: angiotensin converting enzyme inhibitors/angiotensin receptor blockers.
doi:10.1371/journal.pone.0099781.t002

at intermediate (74–167 ng/mL, Hb = -0.22 ± 0.09 , $p = 0.02$) and high ferritin values (>167 ng/mL, Hb = -0.20 ± 0.08 , $p = 0.01$), and increased with hepcidin at low ferritin values (<74 ng/mL, Hb = 0.15 ± 0.07 , $p = 0.04$). Further adjustment for the EPO level did not change the association of Hb and hepcidin (data not shown).

Discussion

This study identified several factors other than mGFR that may determine hepcidin levels in CKD patients. The most original finding is our showing that even though hepcidin levels generally increase as mGFR declines, CKD patients with absolute iron deficiency still experienced a profound hepcidin collapse, in

contrast to those with other iron profiles. We also confirmed the observation of an interaction between iron status and hepcidin in their relation to Hb levels. These findings are especially noteworthy because they are based on a patient population carefully phenotyped for mGFR and a specific and accurate hepcidin-25 assay with a low limit of detection.

Hepcidin is influenced by four regulation processes: hypoxia [16], erythropoiesis, inflammation, and iron status. We confirmed the main determinant factors of hepcidin seen in the general population [17] and in the hemodialysis population [18], [19], [20]. EPO deficiency and the chronic inflammation observed in CKD patients play a role in the hepcidin increase. BMI was a determinant of hepcidin, independently of inflammation. Some hepcidin secretion is located in and directly related to the adipose

Table 3. Crude and mGFR-adjusted Pearson’s correlations of square root-transformed hepcidin with quantitative factors.

	Crude correlation		mGFR-adjusted partial correlation	
	r	p-value	r	p-value
mGFR	-0.31	<.0001	-	-
BMI	0.21	0.003	0.16	0.03
LOG PCR	0.01	0.9	-0.18	0.01
Albuminemia	0.04	0.6	0.17	0.02
Free iron	0.05	0.5	0.18	0.01
TIBC	-0.32	<.0001	-0.22	0.002
TSAT	0.15	0.04	0.24	0.0008
LOG Ferritin	0.62	<.0001	0.71	<.0001
LOG EPO	-0.20	0.01	-0.18	0.03

mGFR: measured glomerular filtration rate; BMI: body mass index; PCR: urinary protein to creatinine ratio; CRP: C-reactive protein; EPO: erythropoietin; TIBC: total iron binding capacity; TSAT: transferrin saturation.
doi:10.1371/journal.pone.0099781.t003




Figure 1. Hepcidin concentration according to quintiles of ferritin (A), transferrin saturation (B) and total iron-binding capacity (C).
doi:10.1371/journal.pone.0099781.g001

tissue [21]. Finally, iron status influences hepcidin levels. A low iron store down-regulates hepcidin, in a relation modulated by transferrin [22], [23]. Our 3-marker iron index with five classes produced similar results in this analysis (data not shown) [24]. In our cohort, the combined iron marker we used to classify the different iron profiles sheds light on the persistent process of hepcidin down-regulation in absolute iron deficiency at the same time that the patients with all other iron profiles had increasing hepcidin levels associated with mGFR decline. This persistent down-regulation has also been shown in animal models of chronic disease anemia [5]. The down-regulation process remains effective in and specific to absolute iron deficiency in CKD patients. The possibility of defining a hepcidin threshold value for this diagnosis requires further validation in CKD cohorts.

The influence of mGFR on hepcidin has been interpreted in different ways in recent articles reporting the role of the hepcidin assay. Hepcidin measurement is based on either mass spectrometry or antibody-based hepcidin detection. Competitive radioimmunoassays and ELISA tests have been developed for the antibody-based methods. Several studies have reported that hepcidin increases as mGFR declines [4], [17], [25]. The hepcidin measurements in these articles, however, were performed with a radioimmunoassay that might have cross-reacted with the isoform hepcidin-20 for about 10% of the total measured value of hepcidin-25. Further contestation of the relation between mGFR

and hepcidin has been based on suspicion that this relation is restricted to hepcidin-20, which is inactive [26]. A large cohort study of 505 CKD patients that used a highly specific liquid chromatography tandem mass-spectrometry method found that estimated GFR was associated with hepcidin levels in the univariate but not the multivariate analysis [27]. Finally, from a physiological point of view, the isoform hepcidin-25, the active form, is a small peptide, freely filtered by the glomerulus and highly reabsorbed by the proximal tubules with an excretion fraction around 8%; it shares the renal clearance characteristics of $\beta 2$ microglobulin [28], [29]. A major strength of our study is the use of $^{51}\text{Cr-EDTA}$ renal clearance to determine GFR, which reduced measurement errors and improved our ability to understand the relation between renal function and hepcidin level. It is likely that levels of the serum isoform hepcidin-25 increase with declining renal function, but the relation between mGFR and hepcidin may have been disrupted by patients with absolute iron deficiency because of their down-regulated hepcidin level. Besides the difference in methods, differences in study populations could have influenced the relation between hepcidin and mGFR in some studies.

Anemia in CKD cohorts has been associated with elevated hepcidin levels. This association reflects the ratios of the prevalence rates of the different iron disturbances: absolute ID, which depresses hepcidin, is relatively rare in CKD cohorts such as

Table 4. Hepcidin levels (ng/mL) and percentage of patients with low hepcidin values according to ferritin and transferrin saturation (TSAT) levels.

		Ferritin		Total
		≥ 40 ng/mL	< 40 ng/mL	
TSAT	$\geq 20\%$	34.5 (23.7–51.6) ² 10% (12) N = 123	10.3 (6.9–14.1) ² 80% (12) N = 15	31.2 (20.9–45.7) ¹ 17% (24) N = 138
	$< 20\%$	22.6 (16.3–43.1) ^{1,2} 29% (12) N = 41	5.0 (0.7–11.7) ^{1,2} 92% (11) N = 12	19.2 (12.2–40.3) 43% (23) N = 53
Total		32.1 (21.35–51.3) 15% (24) N = 164	9.6 (3.8–12.7) 85%(23) N = 27	

Median (Interquartile range).

% (N) of patients with hepcidin values below a threshold value of hepcidin defined as the 10th percentile of patients with normal iron profile (16.7 ng/mL).

¹Three classes of Lipschitz's iron index.

²Four classes iron index.

doi:10.1371/journal.pone.0099781.t004

Table 5. Multivariate analyses of hepcidin levels according to different definitions of iron status.

	N	BIC/AIC	hepcidin $\beta \pm \text{sd}$	p-value
Lipschitz's iron index^{1,2}		877/821		
TSAT $\geq 20\%$	139		ref	
TSAT $< 20\%$ and Ferritin $< 40 \mu\text{g/L}$	12		-2.71 ± 0.55	$< .0001$
TSAT $< 20\%$ and Ferritin $\geq 40 \mu\text{g/L}$	41		-0.32 ± 0.32	0.3
Missing	7		-0.04 ± 0.71	0.9
Four classes iron index^{1,2}		848/788		
TSAT $\geq 20\%$ and Ferritin $\geq 40 \mu\text{g/L}$	123		ref	
TSAT $\geq 20\%$ and Ferritin $< 40 \mu\text{g/L}$	15		-2.84 ± 0.46	$< .0001$
TSAT $< 20\%$ and Ferritin $< 40 \mu\text{g/L}$	12		-3.25 ± 0.51	$< .0001$
TSAT $< 20\%$ and Ferritin $\geq 40 \mu\text{g/L}$	41		-0.65 ± 0.30	0.03
Missing	8		-0.16 ± 0.61	0.8
Four classes iron index according to CRP levels¹		848/785		
TSAT $\geq 20\%$, Ferritin $\geq 40 \mu\text{g/L}$				
CRP $\leq 8 \text{ ng/mL}$	115		ref	
CRP $> 8 \text{ ng/mL}$	8		1.87 ± 0.61	0.002
TSAT $\geq 20\%$ and Ferritin $< 40 \mu\text{g/L}$	15		-2.54 ± 0.46	$< .0001$
TSAT $< 20\%$ and Ferritin $< 40 \mu\text{g/L}$	12		-3.09 ± 0.51	$< .0001$
TSAT $< 20\%$, Ferritin $\geq 40 \mu\text{g/L}$				
CRP $\leq 8 \text{ ng/mL}$	32		-0.90 ± 0.32	0.005
CRP $> 8 \text{ ng/mL}$	9		2.33 ± 0.57	$< .0001$
Missing	8		0.06 ± 0.60	0.9

Abbreviations: TSAT, Transferrin saturation; CRP, C-reactive protein; BIC, Bayesian Information Criterion; AIC, Akaike information Criterion; Ref, reference class.

$\beta \pm \text{sd}$ Regression coefficients for the different iron indexes in the linear regression models of hepcidin values (square-root transformed).

¹Models were adjusted for measured glomerular filtration rate, gender, age, body mass index, albuminemia, erythropoietin, oral iron and centre.

²Models for Lipschitz's iron index and the four-class iron index were also adjusted for C-reactive protein (CRP) in two classes (≤ 8 vs $> 8 \text{ ng/mL}$).

doi:10.1371/journal.pone.0099781.t005

ours (6.3%). Functional iron deficiency is more frequent and is associated with an elevated hepcidin level. Our functional iron profile group had elevated hepcidin levels, similar to those of the normal iron profile group. Combining a functional iron deficiency and overt inflammation further increased hepcidin in our study. The short half-life of hepcidin (22 hours), its high intra-patient variability, and the cross-sectional design of our study might have weakened the link between functional iron deficiency and a hepcidin increase. Recently in a mouse model, the hepcidin increase has been proved to be limited to the early phase of the inflammation process [30].

We confirmed that the relation between Hb and hepcidin varies according to iron status. In the study by Uehata et al [27], hepcidin was not related to Hb in the low ferritin group ($< 91 \text{ ng/mL}$) and was negatively related to it in the high ferritin group. The low ferritin group probably had absolute iron deficiency, with a very low hepcidin level of 7 ng/mL (IQR 2.3–17.3 ng/mL , with liquid chromatography mass spectrometry). We showed that in the absolute iron deficiency group, Hb decreased with hepcidin. On the contrary, in patients with functional iron deficiency, anemia was related to an increased hepcidin level. The hepcidin level thus provides information about Hb above and beyond that contained in the iron profiles. Elevated hepcidin can predict poor gut absorption of iron and ineffective oral iron treatment. In vitro studies suggest that hepcidin has a direct effect on erythropoiesis by inhibiting erythroid colony formation when erythropoietin is low and by impairing red blood cell survival. To date, however, no

study has found that hepcidin has any predictive value for response to either IV iron or ESA.

The limitations of our study are linked mainly to its cross-sectional design, its sample size, and the measurement method. The cross-sectional design makes it difficult to infer causality between the hepcidin level and the risk of anemia. In terms of sample size among non-dialysis CKD cohorts, our study is the second largest examining hepcidin levels, after that of Uehata et al [27]. The immunochemical method that we used measured hepcidin levels in patients with absolute iron deficiency similar to those in the study of Uehata et al, which used mass spectrometry. The antibody-based hepcidin assays may diagnose and discriminate absolute iron deficiency quite well. On the contrary, when hepcidin values are elevated, the lower specificity for hepcidin-25 of the immunochemical test we used might have impaired its power to discriminate patients with normal iron profile from those with functional iron deficiency profile.

In conclusion, this study primarily shows the importance of considering iron profiles in improving our understanding of the changes of the hepcidin concentration as renal function declines as well as its association with CKD anemia. While absolute iron deficiency is associated with a profound hepcidin collapse independently of the GFR decline, hepcidin is elevated in all other CKD patients and thus impairs the oral iron absorption in these patients. These findings have prompted research on drug development focusing on a hepcidin antagonist in CKD and possibly other chronic disease anemia with functional iron

deficiency, although the possible side effects of these drugs may limit their potential.

Acknowledgments

We are grateful to and would like to thank Pr Nicole Casadevall for her advice and the EPO measurements performed in her laboratory (Laboratoire d'Hématologie, Hôpital Saint-Antoine, Paris, France).

We are indebted and thankful to Dr Barbara Sasu and Vimal Patel who advised us and performed hepcidin measurements at Amgen laboratories (Amgen Inc., Thousand Oaks, CA). In order to enable hepcidin measurements, this study was labelled as an Investigator Sponsored Study (BS) by Amgen.

NephroTest Steering Committee

Martin Flamant, P. Houillier, Jean Philippe Haymann, Jean-Jacques Boffa, Eric Thervet, François Vrtovsniak, Bénédicte Stengel (benedicte.stengel@inserm.fr).

References

- Ulukol B, Orhon FS, Hanoluk A, Akar N (2007) Serum pro-hepcidin levels and relationship with ferritin in healthy individuals non-anaemic infants. *Acta Haematol* 118(2): 70–2.
- Kato A, Tsuji T, Luo J, Sakao Y, Yasuda H, et al. (2008) Association of prohepcidin and hepcidin-25 with erythropoietin response and ferritin in hemodialysis patients. *Am J Nephrol* 28(1): 115–121.
- Malyszko J, Malyszko JS, Mysliwiec M (2009) A possible role of hepcidin in the pathogenesis of anemia among kidney allograft recipients. *Transplant Proc* 41(8): 3056–9.
- Tomosugi N, Kawabata H, Wakatabe R, Higuchi M, Yamaya H, et al. (2006) Detection of serum hepcidin in renal failure and inflammation by using ProteinChip System. *Blood* 108(4): 1381–7.
- Theurl I, Aigner E, Theurl M, Nairz M, Seifert M, et al. (2009) Regulation of iron homeostasis in anemia of chronic disease and iron deficiency anemia: diagnostic and therapeutic implications. *Blood* 113(21): 5277–86.
- Grebentchikov N, Geurts-Moespot AJ, Kroot JJ, den Heijer M, Tjalsma H, et al. (2009) High-sensitive radioimmunoassay for human serum hepcidin. *Br J Haematol* 146(3): 317–25.
- Li H, Rose MJ, Tran L, Zhang J, Miranda LP, et al. (2009) Development of a method for the sensitive and quantitative determination of hepcidin in human serum using LC-MS/MS. *J Pharmacol Toxicol Methods* 59(3): 171–80.
- Ashby DR, Gale DP, Busbridge M, Murphy KG, Duncan ND, et al. (2009) Plasma hepcidin levels are elevated but responsive to erythropoietin therapy in renal disease. *Kidney Int* 75(9): 976–81.
- Zaritsky J, Young B, Gales B, Wang HJ, Rastogi A, et al. (2010) Reduction of serum hepcidin by hemodialysis in pediatric and adult patients. *Clin J Am Soc Nephrol* 5(6): 1010–4.
- Malyszko J, Malyszko JS, Mysliwiec M (2009) Hyporesponsiveness to erythropoietin therapy in hemodialyzed patients: potential role of prohepcidin, hepcidin, and inflammation. *Ren Fail* 31(7): 544–8.
- Costa E, Swinkels DW, Laarakkers CM, Rocha-Pereira P, Rocha S, et al. (2009) Hepcidin serum levels and resistance to recombinant human erythropoietin therapy in haemodialysis patients. *Acta Haematol* 122(4): 226–9.
- Ford BA, Eby CS, Scott MG, Coyne DW (2010) Intra-individual variability in serum hepcidin precludes its use as a marker of iron status in hemodialysis patients. *Kidney Int* 78(8): 769–73.
- Moranne O, Froissart M, Rossert J, Gauci C, Boffa JJ, et al. (2009) Timing of onset of CKD related metabolic complications. *J Am Soc Nephrol* 20(1): 164–71.
- Mercadal L, Metzger M, Casadevall N, Haymann JP, Karras A, et al. (2012) Timing and determinants of erythropoietin deficiency in chronic kidney disease. *Clin J Am Soc Nephrol* 7(1): 35–42.
- Lipschitz DA, Cook JD, Finch CA (1992) A clinical evaluation of serum ferritin as an index of iron stores. *Nutrition* 8(6): 443–8.
- Hintze KJ, McClung JP (2011) Hepcidin A Critical Regulator of Iron Metabolism during Hypoxia. *Adv Hematol* 2011; 510304.
- Galesloot TE, Vermeulen SH, Geurts-Moespot AJ, Klaver SM, Kroot JJ, et al. (2011) Serum hepcidin: reference ranges and biochemical correlates in the general population. *Blood* 117(25): e218–25.
- van der Weerd NC, Grooteman MP, Bots ML, van den Dorpel MA, den Hoedt CH, et al. (2012) Hepcidin-25 in chronic hemodialysis patients is related to residual kidney function and not to treatment with erythropoiesis stimulating agents; CONTRAST Investigators. *PLoS One* 7(7): e39783.
- Valenti L, Messa P, Pelusi S, Camprostrini N, Girelli D (2013) Hepcidin levels in chronic hemodialysis patients: a critical evaluation. *Clin Chem Lab Med* 14: 1–7.
- Pelusi S, Girelli D, Rametta R, Camprostrini N, Alfieri C, et al. (2013) The A736V TMPRSS6 polymorphism influences hepcidin and iron metabolism in chronic hemodialysis patients: TMPRSS6 and hepcidin in hemodialysis. *BMC Nephrol* 14: 4.
- Bekri S, Gual P, Anty R, Luciani N, Dahman M, et al. (2006) Increased adipose tissue expression of hepcidin in severe obesity is independent from diabetes and NASH. *Gastroenterology* 131(3): 788–96.
- Bartnikas TB, Andrews NC, Fleming MD (2011) Transferrin is a major determinant of hepcidin expression in hypotransferrinemic mice. *Blood* 117(2): 630–7.
- Gkouvatso K, Papanikolaou G, Pantopoulos K (2012) Regulation of iron transport and the role of transferrin. *Biochim Biophys Acta* 1820(3): 188–202.
- Mercadal L, Metzger M, Haymann JP, Thervet E, Boffa JJ, et al. (2014) NephroTest Study Group. A 3-Marker Index Improves the Identification of Iron Disorders in CKD Anaemia. *PLoS One* 9(2): e84144.
- Zaritsky J, Young B, Wang HJ, Westerman M, Olbina G, et al. (2009) Hepcidin—a potential novel biomarker for iron status in chronic kidney disease. *Clin J Am Soc Nephrol* 4(6): 1051–6.
- Peters HP, Laarakkers CM, Swinkels DW, Wetzels JF (2010) Serum hepcidin-25 levels in patients with chronic kidney disease are independent of glomerular filtration rate. *Nephrol Dial Transplant* 25(3): 848–53.
- Uchata T, Tomosugi N, Shoji T, Sakaguchi Y, Suzuki A, et al. (2012) Serum hepcidin-25 levels and anemia in non-dialysis chronic kidney disease patients: a cross-sectional study. *Nephrol Dial Transplant* 27(3): 1076–83.
- Canavesi E, Alfieri C, Pelusi S, Valenti L (2012) Hepcidin and HFE protein: Iron metabolism as a target for the anemia of chronic kidney disease. *World J Nephrol* 1(6): 166–176.
- Peters HP, Laarakkers CM, Pickkers P, Masereeuw R, Boerman OC, et al. (2013) Tubular reabsorption and local production of urine hepcidin-25. *BMC Nephrol* 14: 70.
- Kim A, Fung E, Parikh SG, Valore EV, Gabayan V, et al. (2014) A mouse model of anemia of inflammation: complex pathogenesis with partial dependence on hepcidin. *Blood* 123(8): 1129–36.

NephroTest Study Group

Bichat Hospital: François Vrtovsniak, Eric Daugas (Nephrology) and Martin Flamant, Emmanuelle Vidal-Petiot (Physiology); European Georges Pompidou Hospital: Christian Jacquot, Alexandre Karras, Eric Thervet, Christian d'Auzac (Nephrology) and P. Houillier, M. Courbebaisse, D. Eladari et G. Maruani (Physiology); Tenon Hospital: Jean-Jacques Boffa, Pierre Ronco, H. Fessi, Eric Rondeau (Nephrology) and Emmanuel Letavernier, Jean Philippe Haymann (Physiology); clinique du Landy, Saint-Ouen: P. Urena-Torres.

Author Contributions

Conceived and designed the experiments: M. Froissart BS LM MM. Analyzed the data: MM LM. Wrote the paper: LM MM BS. Carried out the laboratory reference measurements: M. Flamant JPH PH M. Froissart. Managed the biobank: M. Froissart BS. Managed the data collection: MM BS M. Froissart. Discussed analyses, interpretation and presentation: ET JJB M. Flamant FV PH.