



Complement Factor H, Vitronectin, and Opticin Are Tyrosine-Sulfated Proteins of the Retinal Pigment Epithelium

Yogita Kanan^{1*}, Joseph C. Siefert¹, Michael Kinter^{2,3}, Muayyad R. Al-Ubaidi^{1*}

1 Department of Cell Biology, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, United States of America, **2** Department of Geriatric Medicine, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, United States of America, **3** Free Radical Biology and Aging Research Program, Oklahoma Medical Research Foundation, Oklahoma City, Oklahoma, United States of America

Abstract

Lack of tyrosine sulfation of ocular proteins results in disorganized photoreceptor structure and drastically reduced visual function, demonstrating the importance of this post-translational modification to vision. To understand the role that tyrosine sulfation plays in the function of ocular proteins, we identified some tyrosine-sulfated proteins in the retinal pigment epithelium using two independent methods, immuno-affinity column purification with an anti-sulfotyrosine specific antibody and computer-based sequence analysis of retinal pigment epithelium secretome by means of the prediction program Sulfinator. Radioactive labeling followed by thin layer electrophoresis revealed that three proteins, vitronectin, opticin, and complement factor H (CFH), were post-translationally modified by tyrosine sulfation. The identification of vitronectin and CFH as tyrosine-sulfated proteins is significant, since both are deposited in drusen in the eyes of patients with age-related macular degeneration (AMD). Furthermore, mutations in CFH have been determined to be a major risk factor in the development of AMD. Future studies that seek to understand the role of CFH in the development of AMD should take into account the role that tyrosine sulfation plays in the interaction of this protein with its partners, and examine whether modulating sulfation provides a potential therapeutic target.

Citation: Kanan Y, Siefert JC, Kinter M, Al-Ubaidi MR (2014) Complement Factor H, Vitronectin, and Opticin Are Tyrosine-Sulfated Proteins of the Retinal Pigment Epithelium. PLoS ONE 9(8): e105409. doi:10.1371/journal.pone.0105409

Editor: Alfred S. Lewin, University of Florida, United States of America

Received: May 30, 2014; **Accepted:** July 23, 2014; **Published:** August 19, 2014

Copyright: © 2014 Kanan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. Relevant data are included within the paper.

Funding: This study was supported by the National Eye Institute to MRA, Oklahoma Center for the Advancement of Science and Technology to YK, and Foundation Fighting Blindness to MRA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of NIH or any of its institutes.

Competing Interests: The authors have declared that no competing interests exist.

* Email: ykanan@ouhsc.edu (YK); muayyad-al-ubaidi@ouhsc.edu (MRA)

Introduction

Tyrosine *O* sulfation, a post-translational modification employed in higher eukaryotes [1], is catalyzed by two Type II transmembrane enzymes, tyrosylprotein sulfotransferases 1 & 2 (TPST 1 & 2). It was initially described by Bettelheim in 1954, but was later found to be a common post-translational modification [2,3]. Tyrosine sulfation occurs in the trans-Golgi compartment and requires 3'-phosphoadenosine 5'-phosphosulfate (PAPS) as a sulfate donor for the reaction [4]. It is only observed on secreted and transmembrane proteins: nuclear and cytoplasmic proteins have not been reported to have this modification [1,5]. However, the role of tyrosine sulfation in protein function has only recently been investigated [6–8].

Initial analyses of the amino acid sequences surrounding the identified sulfated tyrosines showed a predominance of acidic amino acids within 5 residues surrounding the sulfated tyrosines [9]. However, later studies showed that some tyrosine-sulfated proteins do not follow these criteria, and it is the secondary structure that may expose the tyrosine residue to a TPST to be sulfated [10].

Mouse models that lack either or both TPST enzymes exhibit distinctly different phenotypes [11–13]. Mice lacking both TPSTs

show the most drastic phenotype of cardio-pulmonary insufficiency and subsequent death within 2 months after birth [11]. Previous studies have demonstrated that these animals also display ocular defects [14,15]. The *Tpsti*^{-/-} mice have reduced rod electroretinographic (ERG) responses during early development, but these normalize by postnatal day 90 [14]. However, the *Tpst2*^{-/-} mice exhibit a non-progressive reduction in rod and cone ERG functions that persists throughout the life of the animal [14]. Most importantly, the rod and cone ERG responses of the double knockout mice were reduced to 25% and 15% of normal littermates' levels [15].

The significant reduction in rod and cone light-evoked responses in the absence of both TPSTs underscores the role that tyrosine *O*-sulfated proteins play in vision. Previous work using the anti-sulfotyrosine antibody (PSG2) showed that tyrosine-sulfated proteins are present in different ocular tissues, including the neurosensory retina and the retinal pigment epithelium (RPE) [16,17]. Although most of the tyrosine-sulfated proteins present in the interphotoreceptor matrix are produced locally, others originate in the RPE [17]. As a first step towards the systematic identification of tyrosine-sulfated proteins involved in sensory retina/RPE function and homeostasis, we used two approaches to

isolate and characterize tyrosine *O*-sulfated proteins in the RPE. The first method involved immunoaffinity purification from cow RPE extracts using the anti-sulfotyrosine antibody PSG2, as has been established previously [18]. Several proteins were identified by mass spectrometry analysis following the affinity purification. Further analyses by ectopic expression and barium hydroxide hydrolysis confirmed that vitronectin and opticin were tyrosine-sulfated.

Affinity purification using PSG2 [19] has its own limitations, such as the co-purification of partners that interact with tyrosine-sulfated proteins that may not be tyrosine sulfated, and the lack of recognition by PSG2 due to variations in the sequences surrounding the sulfation site. Therefore, a second approach was adopted in which the secretome profile of human RPE [20] was examined for proteins associated with/causing retinal diseases listed on RetNet (<https://sph.uth.edu/retnet/>). While three proteins in the RPE secretome, collagen type 2, alpha 1, and complement factor H (CFH), were identified to be involved in retinal disease, CFH was the only candidate predicted to be tyrosine-sulfated by the program Sulfinator [21]. Sulfinator predicted that CFH contains 5 putative sulfated tyrosines. Subjecting it to barium hydroxide hydrolysis confirmed that, similar to vitronectin and opticin, CFH is also tyrosine sulfated. The finding that both CFH and vitronectin are sulfated, combined with the identified role of tyrosine sulfation in protein-protein interactions [6,22–24], is of significance to studies of AMD since drusen, which are characteristic extracellular deposits in AMD, contain both CFH and vitronectin [25,26]. Furthermore, some AMD cases are caused by mutations in CFH [27–30].

Opticin, as an extracellular matrix protein, has been shown to have anti-angiogenic properties [31], and can bind collagen [32] and retinal growth hormone (GH) in chick embryonic vitreous humor [33]. Similar to CFH and vitronectin, it would be interesting to determine in future experiments whether tyrosine sulfation modulates the interaction of opticin with collagen or GH and its anti-angiogenic properties.

Materials and Methods

Animal studies and ethics statement

Mouse work was performed after approval from the Institutional Animal Care and Use Committee at the University of Oklahoma Health Sciences Center (IACUC 13-001) and strictly adhered to rules and regulations set forth by the National Institute of Health Guide for the Care and Use of Laboratory Animals and the Association for Research in Vision and Ophthalmology Resolution on the Use of Animals in Research. Adult mice were euthanized using CO₂ asphyxiation. Then, the sensory retina and RPE were harvested.

Preparation of lysates from human donor eyes

Three human donor eyes were from 57-to-67-year-old Caucasian males that had no visual problems. Eyes were obtained from either Lions Eye Institute (Tampa, FL) or from the Illinois Eye Bank (Chicago, IL). We dissected the eyes and separated the neurosensory retina and RPE, flash froze each in liquid nitrogen, and then stored at -80°C until use. Small samples of each were homogenized in buffer A (25 mM MOPS, 100 mM NaCl, pH 7.5).

Preparation of lysates from cow and pig eyes

Adult cow and pig eyes were obtained from Country Home Meat Slaughter House (Edmond, OK). Eyes were dissected and RPE was immediately frozen in liquid nitrogen until lysate

preparation as described above. Cows were not tested for bovine spongiform encephalopathy (BSE) since they were under 30 months of age, and none of the cows in the herd in that region showed any signs of the disease.

Processing of cow eyes for affinity purification

The RPE from four fresh independent adult cow eyes were homogenized in buffer A (above) in a dounce homogenizer. The homogenate was centrifuged at $50,000\times g$ for 30 min at 4°C and the supernatant was collected. Protein concentrations were determined in the supernatant fraction by Bradford assay, adjusted to 4 mg/ml in wash buffer 1 (W1, 25 mM MOPS, 100 mM NaCl), and loaded on the PSG2 column.

PSG2 affinity purification of tyrosine O sulfated proteins

About 10 mg of cow RPE supernatant was filtered using a 0.45 μm syringe filter (Millipore, Billerica, MA) and loaded onto the PSG2-Affi-Gel-10 HPLC column at a flow rate of 0.1 ml/min [18]. The column was washed successively with wash buffer 1 (W1, 25 mM MOPS, 100 mM NaCl), wash buffer 2 (W2, 25 mM MOPS, 200 mM NaCl), and wash buffer 3 (W3, 25 mM MOPS, 400 mM NaCl) at a flow rate of 0.2 ml/min, then eluted with elution buffer (EB, 25 mM MOPS, 400 mM NaCl, 4 mM sulfated pentapeptide). The entire run was monitored by recording absorbance at 280 nm. The eluted samples were concentrated with acetone precipitation and fractionated by SDS-PAGE. The tyrosine-sulfated pentapeptide LDY^SDF was synthesized (Bio-Synthesis Inc., Lewisville, TX).

Mass spectrometry

Column fractions were separated by SDS-PAGE and the gel lane was cut into 1 mm slices. Each slice was reduced, alkylated, and subjected to in-gel trypsin digestion. The samples were analyzed using a Thermo Scientific LTQ-XL linear ion trap system with an Eksigent splitless nanoflow HPLC. Ten μL volumes of each digest were injected. A data-dependent analysis acquired one mass spectrum and 9 collision induced dissociation (CID) spectra per cycle. The CID spectra were used to search the cow RefSeq database using the program Mascot. All identified proteins exceeded a minimal identification criteria of at least 2 CID spectra matching unique peptide sequences with ion scores greater than 50. Since at this time only secreted or transmembrane proteins have been identified as tyrosine-sulfated [1,5], only these proteins were included in Tables 1 & 2. Nuclear and cytoplasmic proteins that co-purified in the column were therefore eliminated.

RPE secretome analysis

Members of the RPE secretome [20] were screened for disease-causing proteins listed in the Retinal Information Network (RetNet). The candidate proteins that caused disease were then plugged into the tyrosine sulfation prediction program Sulfinator [21] to predict their tyrosine sulfation status.

Immunoprecipitation and immunoblotting

Immunoprecipitation and immunoblotting were performed according to previously published methods [17,34]. The following antibodies were used: anti-vitronectin antibody (Santa Cruz Biotechnology Inc., Dallas, TX), anti-opticin antibody (Santa Cruz Biotechnology Inc.), and anti-CFH antibody (AbD Serotec, Raleigh, NC). The purified human plasma CFH was obtained from AbD Serotec.

Table 1. List of potential tyrosine-sulfated proteins in cow RPE.

	Protein name	Mascot Score	RefSeq#	Coverage %
*	Opticin	605	45429965	26
*	Lumican	630	27806853	31
*	Vitronectin	313	78045497	15
*	Heparin cofactor II	722	76639676	26
*	Similar to Amyloid beta A4 protein precursor	851	76607645	22
1.	Fibrinogen, gamma	1093	27806893	43
2.	Complement C4	978	76650940	10
3.	Dickkopf related protein 3	863	76635678	28
4.	Fibulin 2	774	76649536	13
5.	Fibrinogen beta	657	76638241	49
6.	Secreted frizzled related protein	662	27806625	39
7.	Tubby like protein 1	557	76672237	54
8.	Spondin 1	480	27807443	14
9.	Retinol Binding protein 3 (IRBP)	467	27806445	9
10.	Fibroblast growth factor	381	27806627	31
11.	Fibrinogen alpha	319	75812954	10

Sixteen proteins were identified in cow RPE by MALDI-MS analysis of SDS-PAGE gel slices of PSG2 immunoaffinity column eluent. The asterisks mark those proteins at 75 and 50 kD, shown in Figure 2C. NCBI reference sequence database ID numbers (RefSeq#) and percent sequence coverage (coverage %) are also indicated for each protein. Inclusion in this table does not confirm that the protein is tyrosine-sulfated. Some of the proteins may be isolated by the affinity column as a result of their interaction with sulfated proteins.

doi:10.1371/journal.pone.0105409.t001

Table 2. List of potential tyrosines that may be sulfated on human and cow vitronectin, opticin, and CFH, as identified by Sulfinator.

Name	Site	Sequence	Reference for Tyrosine sulfation
	75*	MPDEYTVYDD	
	78*	DEYTVYDDGEE	
Human Vitronectin	282	KQYWEYQFQHR	Yu Y, Hoffhines AJ, Moore KL, Leary JA (2007) Determination of the sites of tyrosine O-sulfation in peptides and proteins. Nat Methods 4: 583–588.
	417	LGANNYDDYRM	
	420	NNYDDYRMDWL	
Cow Vitronectin	75	LPEDEYGFHDY	Novel
Human Opticin	65	IDLSNYEELTD	Novel
	71	EELTDYGDQLP	
Cow Opticin	61	DELIDYGDQLP	Novel
Human CFH	243	NMGYEYSERGD	
	534	NDTLDYECHDY	
	709	LSSPPYYGDS	Novel
	710	SSPPYYGDSV	
	711	SPPYYGDSVE	
Cow CFH	168	EPDQEYTYGQV	Novel
	170	DQEYTYGQVVQ	
	465	ESTFTYPLNKQ	
	473	NKQTEYKCKPG	
	575	PEMDPYLNAYP	
	579	PYLNAYPRKET	
	585	PRKETYKVGDV	

Those sites that have been experimentally proven are marked by an asterisks and citation is provided.

doi:10.1371/journal.pone.0105409.t002

Metabolic labeling, barium hydroxide hydrolysis, and thin layer electrophoresis analysis

HEK 293T cells were independently transiently transfected with recombinant human vitronectin (VTN, Genecopoeia, Rockville, MD), recombinant myc-tagged human Opticin (OPTC, Genecopoeia), or human recombinant complement factor H (CFH, Genecopoeia). Eight hours after transfection, media was replaced with sulfate-free Joklik-modified Eagle's media (Sigma, St. Louis, MO) containing 2% dialyzed fetal bovine serum. To this media, 0.15 mCi/ml of $\text{Na}_2^{35}\text{SO}_4$ (Perkin Elmer, Waltham, MA) was added. About 48 hours later, the media was harvested, fractionated by SDS-PAGE, and transferred onto polyvinylidene difluoride (PVDF) membranes (Millipore, Billerica, MA). The radioactive bands were cut out and subjected to barium hydroxide hydrolysis and thin layer electrophoresis (TLE) according to published methods [11,18].

Deglycosylation assays

CFH was treated for 2 hours at 37°C with PNGase F, according to the manufacturer's instruction (NEB BioLabs, Ipswich, MA).

Results

RPE extracts are a rich reservoir of tyrosine-sulfated proteins

Tyrosine-sulfated proteins have been previously shown to be expressed in the neurosensory retina and RPE [16,17]. Furthermore, some of the sulfated proteins that were present in the neurosensory retina originated in the RPE [17]. To determine the differential distribution of tyrosine-sulfated proteins in the neurosensory retina and RPE, immunoblot analyses of mouse, pig, cow, and human neurosensory retinal and RPE lysates were performed and probed with PSG2. The analysis revealed that the RPE harbors a relatively higher number of tyrosine-sulfated proteins than the neurosensory retina in each of the four mammalian species tested (Figure 1). The sizes of sulfated proteins in the RPE ranged from 20 kD to >250 kD. While some sulfated protein bands appeared to be conserved across species tested, other tyrosine-sulfated protein bands seemed to be species-specific. An across-species conserved sulfated protein may appear at different sizes on immunoblots because tyrosine-sulfated proteins are secreted proteins and, therefore, may be differentially glycosylated [35,36] due to species-specific glycosylation patterns. The tyrosine-sulfated proteins in human lysates appear more prominent when compared to those in other species. This may be partly due to the fact that PSG2 was raised against the tyrosine-sulfated N terminus of human PSGL-1 [19], and hence may better recognize human tyrosine-sulfated proteins. Alternatively, since recognition by PSG2 was shown to be context-dependent [19], better recognition of human proteins may also have to do with the environment surrounding the sulfated tyrosines.

Immunoaffinity purification of tyrosine-sulfated proteins from the RPE

We chose to use cow RPE extracts in the affinity purification with PSG2 because RPE lysates contain more tyrosine-sulfated proteins than the neurosensory retina, most of the sulfated proteins that appear in human RPE are also observed in cow RPE, and it was easy to obtain a sufficient number of independent samples. Monitoring the column flow-through by measuring absorbance at 280 nm showed that most of the unbound proteins passed through the column in the flow-through (FT) fraction (Figure 2A). The remaining nonspecifically bound proteins were washed off the




Figure 1. Immunoblot analysis of neurosensory (NS) retinal and RPE extracts to identify tyrosine-sulfated proteins. Immunoblot analysis of 50 μg of RPE (lanes 1–4) and neurosensory retinal extracts (lanes 5–8) from human, cow, pig, and mouse RPE probed with the anti-sulfotyrosine antibody PSG2. Blots were repeated 3 independent times using biologically different samples. doi:10.1371/journal.pone.0105409.g001

column with increasing salt washes (W1, W2 & W3, Figure 2A), while the potential tyrosine-sulfated proteins were eluted with EB containing 4 mM sulfated pentapeptide (EB, Figure 2A).

Proteins in the FT, W1, W2, W3, and EB fractions were separated by SDS-PAGE, followed by Coomassie blue staining of the gel. Comparing equal volume aliquots from each fraction confirmed the column absorbance results and showed that the majority of the non-specific proteins were already cleared away by W2 (compare W2 and W3, Figure 2B&C). The EB fraction contained two enriched bands (asterisks, Figure 2C, CB) visible after Coomassie blue staining of the gel. PSG2-immunoblot analysis of EB showed that the two enriched bands were tyrosine-sulfated proteins: it is important to note that there were other tyrosine-sulfated proteins in this fraction that were not obvious by Coomassie blue staining (Figure 2C, PSG2). Interestingly, the upper asterisk-marked band in Figure 2C was also observed in W3, and was also sulfated, as shown after probing with PSG2 (W3&EB, Figure 2C). While only 20% of the peptide eluted fraction was run on the SDS-PAGE, the remaining 80% was acetone-precipitated and fractionated by SDS-PAGE for 20 minutes to remove the pentapeptide. The entire lane was sliced into multiple 1 mm portions and subjected to in-gel trypsin digestion, followed by LC-tandem MS analysis. Table 1 presents the identity of 16 proteins that were selectively eluted from the column and identified by MS/MS. Among the identified proteins, lumican, heparin cofactor II, complement C4, amyloid beta protein, fibulin 2, and fibrinogen (α , β , and γ) have been previously identified to be tyrosine-sulfated and their sites of sulfation identified [37–43]. Retinol binding protein 3, Spondin 1,




Figure 2. PSG2-immunoaffinity column purification of tyrosine-sulfated proteins from cow RPE. (A) The elution profile was monitored by following absorbance at 280 nm. Following loading, the column was washed with buffers W1, W2, and W3. Elution was performed in buffer W3 containing 4 mM sulfated pentapeptide (EB). (B) Twenty-six microliter aliquots from input (IN), flow-through (FT), wash 1 (W1), and wash 2 (W2) were fractionated by SDS-PAGE, and proteins were visualized by staining with Coomassie blue dye. (C) Left, SDS-PAGE of 26 μ L of wash 3 (W3) and eluted samples (EB) from the immunoaffinity column stained with Coomassie blue (CB) and right, immunoblotted with PSG2. Asterisks indicate the bands that were prominent on Coomassie blue-stained gel (CB) and were also recognized by PSG2 as tyrosine-sulfated. doi:10.1371/journal.pone.0105409.g002

basic fibroblast growth factor, secreted frizzled related protein, and Tubby-like protein 1 have not previously been shown to be tyrosine-sulfated. It is important to mention that there is no obvious functional relationship between the identified proteins, since they belong to different classes of extracellular proteins. For example, retinol binding protein is involved in the retinoid cycle [44], dickkopf-related protein is involved in a signaling pathway [45], and fibrinogen is involved in the blood clotting pathway [46].

Due to the myriad functions performed by vitronectin, and since opticin has recently been shown to exert anti-angiogenic properties, we chose to focus on vitronectin and opticin. However, the unavailability of anti-bovine specific antibodies capable of immunoprecipitating vitronectin and opticin from bovine RPE limited our studies to human proteins instead. While vitronectin has previously been shown to be tyrosine-sulfated in human hepatoma-derived cell line Hep G2 and human plasma [47,48], its tyrosine sulfation status in the human RPE is unknown. Furthermore, the tyrosine-sulfation of opticin has not been demonstrated.

Vitronectin is tyrosine-sulfated in human RPE

To confirm that human vitronectin is tyrosine-sulfated *in vivo*, we immunoprecipitated it from human RPE lysates using anti-vitronectin antibody. Vitronectin in human RPE was observed to migrate as two bands of sizes 65 kD and 75 kD (lane 1, Figure 3A). These two isoforms have been detected in multiple tissues [49–51]. It has been shown that the 65 kD band results from the cleavage of the 75 kD vitronectin band by the protease furin [51]. Both the 65 kD and 75 kD isoforms, detected in the RPE input, were immunoprecipitated by the anti-vitronectin antibody (asterisks, Figure 3A, upper panel), but not by mouse IgG. Additionally, using PSG2, we detected that both isoforms were tyrosine-sulfated (Figure 3A, lower panel).

While the PSG2 antibody identifies VTN as a tyrosine-sulfated protein, the gold standard for classifying a protein as tyrosine-sulfated is the actual detection of sulfated tyrosines in the protein by barium hydroxide hydrolysis. To accomplish that, recombinant vitronectin was expressed in a heterologous system in the presence of radioactive sulfate, followed by barium hydroxide hydrolysis and thin layer electrophoresis (TLE) in the presence of a non-radioactive sulfated tyrosine standard. The co-localization of the radiolabelled sulfated tyrosine from the recombinant protein with the non-radioactive sulfated tyrosine standard on the TLE plate confirms that the protein of interest is subjected to tyrosine sulfation. To this purpose, recombinant human vitronectin and a control vector (pcDNA 3.1) were transfected into HEK293T cells. Since vitronectin is a secreted protein, media was collected and immunoblotted with anti-vitronectin antibody. Interestingly, only the 75 kD isoform was expressed in cells transfected with recombinant vitronectin (Figure 3B). The sole presence of the 75 kD isoform may be due to the absence of furin in HEK293T cells. The transfected 75 kD vitronectin isoform was immunoprecipitated from the media and shown by PSG2 to be tyrosine-sulfated (Figure 3B). The recombinant vitronectin was then metabolically labeled with 35 S followed by immunoprecipitation and autoradiography to confirm the radiolabeling of the protein (Figure 3C, AR). The radioactive 75 kD band was confirmed to be vitronectin by the anti-vitronectin antibody (Figure 3C). The radiolabeled vitronectin band and a similar-sized membrane from an equivalent region of the IgG control lane were cut from the membrane and subjected to alkaline hydrolysis using the barium hydroxide method described by Huttner et al [2]. The co-localization of the radioactive tyrosine-sulfate from the radiolabelled vitronectin hydrolysates (Figure 3D) with the non-radioac-




Figure 3. Native human RPE vitronectin is tyrosine-sulfated. Vitronectin was immunoprecipitated from 500 μ g human RPE lysates using anti-VTN antibody (lane 2) or mouse IgG (lane 3). Immunoprecipitants were fractionated by SDS-PAGE, transferred, and immunoblotted using anti-VTN antibody or anti-sulfotyrosine PSG2 antibody. Immunoprecipitation and western blots were repeated 3 independent times using biologically different human RPE samples. (B). Ectopically-expressed vitronectin is tyrosine-sulfated. Recombinant VTN or empty vector (pcDNA3.1) were transfected into HEK 293T cells and immunoprecipitated from conditioned media using anti-VTN antibody (lane 4) or mouse IgG (lane 3). Immunoprecipitants were electrophoresed and immunoblotted using anti-VTN antibody or anti-sulfotyrosine PSG2 antibody. Immunoprecipitation and western blots were repeated 3 independent times after independent VTN transfections. (C). 35 S-metabolic labeling of recombinant vitronectin *in vitro*. Vitronectin-transfectants were radiolabelled with 35 Sulfate. Following radiolabeling, vitronectin was immunoprecipitated and blots were either subjected to autoradiography (AR) or immunoblotted with anti-VTN antibody. (D). Radiolabeled vitronectin bands were excised from the membrane along with equivalent areas from mouse IgG immunoprecipitants, and alkaline hydrolysis was performed. The samples were then spiked with sulfo-amino standards tyrosine sulfate, threonine sulfate, and serine sulfate, and subjected to thin layer electrophoresis (TLE) on cellulose plates. Following TLE analysis, sulfo-amino standards (NHD) were visualized either by spraying with Ninhydrin or autoradiography (AR). TLE experiments were repeated at least three independent times. doi:10.1371/journal.pone.0105409.g003

tive tyrosine-sulfate standard detected after ninhydrin staining of the TLE plate (Figure 3D) confirmed that human vitronectin is tyrosine-sulfated.

Opticin is tyrosine-sulfated in vivo

Opticin was identified in human RPE as a 55 kD protein (lane 1, Figure 4A). Since this size is similar to the heavy chain of the antibody, which is also tyrosine-sulfated [52], opticin was immunoprecipitated using anti-opticin antibody and eluted under non-reducing conditions, so that the tyrosine-sulfated heavy chain of the antibody would not mask the opticin isoform. The anti-opticin antibody was successful in bringing down the 55 kD

monomeric opticin band as well as multiple aggregated opticin bands of sizes 55–75 kD, with the 75 kD band being the most abundant (asterisks, Figure 4A). Opticin was not pulled down with the mouse IgG antibody (Figure 4A, upper panel). A companion blot probed with PSG2 showed tyrosine sulfation recognition by the antibody of the 75 kD aggregated opticin band (asterisk, Figure 4A, lower panel).

We then expressed recombinant opticin ectopically in HEK293T cells and metabolically labelled the protein with 35 S. This method provides an advantage over cold experiments since it allow us to differentiate between 35 S-opticin and the non-radiolabeled IgG heavy chain, even under reduced conditions. After radiolabeling, two isoforms (55 kD and 65 kD) were found to have incorporated the radiolabel only in the conditioned media of transfected cells (Figure 4B). It is worth mentioning that the two isoforms of opticin observed here have been reported by other groups [53–55]. Opticin has also been shown to be O-glycosylated and, therefore, the two isoforms may be differentially O-glycosylated isoforms [56]. The two isoforms were immunoprecipitated (lane 4, Figure 4B) and visualized after autoradiography. These bands are identified in Figure 4B as ‘lower band (L)’ for the 55 kD protein and ‘higher band (H)’ for the 65 kD protein.

To verify that the radiolabelled bands were opticin, after autoradiography we blotted the bands with anti-opticin antibody, which recognized both the L and the H isoforms of opticin in the media of transfected cells (Figure 4B). However, in the anti-opticin immunoprecipitated lane (lane 4, Figure 4B), only the 65 kD (H) opticin band is distinct, because the heavy chain IgG band from the antibody masks the L opticin isoform under reducing conditions (Figure 4B). Finally, we performed barium hydroxide hydrolysis to confirm that the radiolabelled isoforms were tyrosine-sulfated. Again, the co-migration of the radioactive tyrosine-sulfate from the radioactive protein with the non-radioactive tyrosine-sulfate standard on TLE plate confirms that opticin contains sulfated tyrosines (Figure 4C).

Analysis of RPE secretome identifies CFH as a potentially sulfated protein

Parallel to the immunoaffinity purification of tyrosine-sulfated proteins from the RPE, we performed an independent analysis on the human RPE secretome [20]. Two criteria were chosen for this analysis. The first was to select members of the RPE secretome in which mutations had been identified to be associated with human visual disorders according to the Retinal Information Network (RetNet). The second criterion was to determine which of these selected proteins is predicted to be tyrosine-sulfated by Sulfinator [21].

Our analysis of the human RPE secretome yielded two tyrosine-sulfated proteins, complement factor H (CFH) and collagen type 2 (alpha 1), which are implicated in AMD and Stickler syndrome, respectively [27,57–59]. Analyses by Sulfinator predicted only CFH as a tyrosine-sulfated protein, with 5 possible sites at tyrosines 243, 534, 709, 710, and 711 (Table 2). In addition, human vitronectin and opticin were similarly analyzed by Sulfinator and were found to be tyrosine-sulfated (Table 2). Human vitronectin was predicted to contain five sulfated tyrosines at 75, 78, 282, 417, and 420. Two of these residues, 75 and 78, have been confirmed to be sulfated [48]. Finally, human opticin was predicted to contain two tyrosine-sulfated residues at positions 65 and 71 (Table 2). It is interesting to note that while vitronectin and opticin were originally identified using immunoaffinity purification from bovine extracts, Sulfinator also predicts bovine opticin and vitronectin to be tyrosine-sulfated (Table 2). In addition, bovine CFH was predicted to be tyrosine-sulfated by




Figure 4. Tyrosine sulfation of ectopically expressed opticin. (A). Opticin was immunoprecipitated from 500 μ g human RPE lysates using anti-OPT antibody (lane 3) or mouse IgG (lane 2). Immunoprecipitants were electrophoresed and immunoblotted using anti-OPT antibody (upper panel) or PSG2 (lower panel). Immunoprecipitation and Western blots were repeated 3 independent times using biologically different human RPE samples. (B). Metabolic labeling of recombinant opticin. Recombinant opticin or empty vectors were transfected into HEK293T cells and cells grown in presence of radioactive sulfate. Following radiolabeling, opticin isoforms were immunoprecipitated using anti-OPT antibody (lane 4) or mouse IgG (lane 3). The blots were then subjected to autoradiography (AR) and immunoblotted for opticin. The 65 kD and 55 kD isoforms are depicted as 'H' and 'L', respectively. Immunoprecipitation and western blots were repeated 3 independent times after independent OPT transfections. (C). Both the 65 kD 'H' and 55 kD 'L' isoforms were excised from the blot, along with equivalent areas around 55 kD from mouse IgG immunoprecipitants, and analyzed by barium hydroxide hydrolysis. The samples were then spiked with a mixture of tyrosine sulfate, threonine sulfate, and serine sulfate as standards, and subjected to thin layer electrophoresis (TLE) on cellulose plates. Following TLE analysis, the plate was sprayed with ninhydrin to visualize the sulfo-amino standards (NHD) and was autoradiographed (AR). TLE experiments were repeated at least three independent times. doi:10.1371/journal.pone.0105409.g004

sequence analysis (Table 2). This suggests the exciting possibility that protein tyrosine sulfation is likely conserved across species.

Complement factor H is tyrosine-sulfated in its native form

CFH has been shown to be mainly produced in the liver and secreted into the plasma [60,61], and secreted by the RPE in ocular tissue [28,62]. To demonstrate that CFH is tyrosine-sulfated *in vivo*, purified human plasma CFH was fractionated by SDS-PAGE and immunoblotted with anti-CFH and PSG2 antibodies. Immunoblot analysis with PSG2 identified human CFH as tyrosine-sulfated (Figure 5A). Since CFH has eight N-glycosylated residues [63], deglycosylation of the protein with PNGase F eliminated the glycosidic residues and decreased the size of the protein by \sim 18 kD [63]. This glycosidic elimination enhanced detection by PSG2 (Figure 5A). The same strategy was adopted to human RPE immunoprecipitated CFH. Human RPE CFH showed the presence of 2 CFH bands, a glycosylated and non-(or partially)-glycosylated isoform (Figure 5A, IB:CFH), even in the absence of deglycosylation with PNGase F. However, PNGase F treatment of the immunoprecipitated CFH collapsed the two bands into one non-glycosylated CFH band (Figure 5A, IB: CFH). Both of the non-glycosylated bands were recognized by PSG2 in the presence and absence of PNGase F (Figure 5A, IB: PSG2). Interestingly, PSG2 showed reactivity only to the native non-glycosylated isoform of CFH, not the glycosylated endogenous CFH (lower right panel of Figure 5A, IB: PSG2).

The same deglycosylation strategy was used in *in vitro* studies. Recombinant CFH was immunoprecipitated from the media of human CFH-transfected HEK293T cells. One portion was subjected to PNGase F treatment while another was left untreated. Immunoblot analysis showed a size decrease with PNGase F treatment (Figure 5B). Both the PNGase F-treated and untreated immunoprecipitants were recognized by PSG2, with the PSG2

showing better reactivity to the PNGase F- treated immunoprecipitant (compare lane 2 to 1, Figure 5B).

In vitro metabolic labeling with 35 S showed that CFH incorporated the label (Figure 5C). PNGase F treatment eliminated \sim 75% of 35 S radioactivity as determined by densitometry (Figure 5C). This suggests that the N-glycosylated residues on CFH are also sulfated, which is in agreement with a previous report [64]. Barium hydroxide hydrolysis and TLE showed that the remaining \sim 25% of label on CFH is on tyrosine(s), as indicated by the co-localization of the radioactive tyrosine sulfate from radioactive CFH with the non-radioactive tyrosine sulfate standard (Figure 5D).

Discussion

Previous studies using the *Tpst1*^{-/-}, *Tpst2*^{-/-}, and double knockout mouse models demonstrated the importance of tyrosine sulfation for vision. Since proper visual transduction results from a hemostatic relationship between two closely associated tissues, the neurosensory retina and the RPE, we compared the abundance of tyrosine-sulfated proteins in these two tissues. Immunoblot analysis with PSG2 showed a greater abundance of tyrosine-sulfated proteins in the RPE compared to the neurosensory retina, across species. To identify tyrosine-sulfated proteins in the RPE, two approaches were used. In the first approach, we used immunoaffinity purification with PSG2 antibody to isolate the tyrosine-sulfated proteins. The second method involved applying the tyrosine sulfate prediction program Sulfinator to analyze proteins in the human RPE secretome to select candidate protein(s) that are involved in human disease, as well as potentially tyrosine-sulfated.

Vitronectin and opticin were identified using the first approach, and complement factor H (CFH) was identified as a possible sulfated protein using the second approach. To confirm that these three proteins are tyrosine-sulfated, multiple methods were




Figure 5. Native human plasma and RPE CFH is sulfated. (A) About 4 μ g of purified human plasma CFH was either untreated (lane 1) or treated with PNGase F (lane 2), and then immunoblotted with anti-CFH antibody or with PSG2. CFH was also immunoprecipitated from human RPE, and was either untreated (lane 1) or treated with PNGase F (lane 2), and then immunoblotted with anti-CFH antibody or with PSG2. Western blots were repeated 3 independent times from human plasma and RPE samples. (B) A recombinant human CFH clone was transfected into HEK293T cells, immunoprecipitated with anti-CFH antibody, and was either directly electrophoresed (lane 1) or first treated with PNGase F (lane 2). The immunoprecipitants were immunoblotted with anti-CFH antibody or with PSG2. Immunoprecipitation and Western blots were repeated 3 independent times after CFH transfections. (C) Recombinant CFH was transfected into HEK293T cells and radiolabelled with radioactive sulfate, then immunoprecipitated and subjected to PNGase treatment (lane 3) or left untreated (lane 2). The blots were autoradiographed (AR), then immunoblotted with anti-CFH antibody. (D) Radiolabelled CFH was analyzed by barium hydroxide hydrolysis and autoradiographed (AR). TLE experiments were repeated at least three independent times.
doi:10.1371/journal.pone.0105409.g005

adapted. A non-radioactive scheme involved immunoprecipitating the said protein with antigen-specific antibodies, followed by immunoblotting with anti-sulfotyrosine antibody PSG2. A radioactive method included metabolically labeling the protein, and performing immunoprecipitation followed by barium hydroxide analysis and thin layer electrophoresis. The barium hydroxide method is the gold standard for confirming tyrosine sulfation because it hydrolyzes the carbohydrate moieties on a protein and has the ability to determine whether the sulfate is on a tyrosine or on a carbohydrate moiety [2].

Opticin was isolated following immunoaffinity purification with PSG2. Further analysis by immunoblotting with PSG2 and barium hydroxide hydrolysis confirmed that it was tyrosine-sulfated. Opticin is an extracellular matrix protein that has recently been shown to have anti-angiogenic properties *in vivo* and *in vitro*. *In vivo* comparisons of wild-type and opticin knockout animals in the oxygen-induced retinopathy model of neovascularization showed more neovascularization in the knockout animal [31]. These results were further examined *in vitro*, in

which it was shown that opticin binds collagen and thereby inhibits endothelial cell integrins $\alpha(1)\beta(1)$ and $\alpha(2)\beta(1)$ from binding collagen, a necessity for pro-angiogenic signaling [32]. It has also been shown that opticin can bind retinal growth hormone (GH) in chick embryonic vitreous humor [33]. It would be interesting to see if removing sulfation on opticin modulates its interaction with collagen or GH and if it affects anti-angiogenic functions.

Vitronectin has been previously shown to be tyrosine-sulfated in human plasma [48]. Its tyrosine-sulfated residues were identified as Tyr-75 and Tyr-78 [48]. These two residues are close to the 'RGD' cell attachment site on the protein, which resides between residues 64–66. The RGD sites on vitronectin have been previously shown to bind integrin receptor $\alpha v\beta 3$ and $\alpha v\beta 5$ [52,65]. However, for this to occur, the 'RGD' site must be exposed to the surface, which can be influenced by type of surrounding residues. For example, the presence of a proline residue that follows the 'RGD' site silences the motif by preventing surface accessibility [66]. Therefore, the function of tyrosine sulfation, due to its hydrophilic nature and close proximity to the 'RGD' site, may be to expose the RGD domain at the surface of the protein, facilitating its interaction with integrin receptors on cells. Since tyrosine sulfation has been shown to be necessary for protein-protein interactions [6,22–24], the presence of a highly charged sulfate group may facilitate the interaction of vitronectin with a positively charged domain(s) on an interacting partner. As the RGD-Integrin attachment influences many functions such as cell migration, adhesion, growth, and differentiation [67,68], the presence of tyrosine sulfation may potentially modulate all of these functions.

Another function of vitronectin is complement regulation. Vitronectin inhibits the complement cascade by binding the membrane attack complex (MAC), the final product in the complement cascade [69], and therefore protects cells against complement attack. Complement damage in the eye is a leading cause of vision loss and causes AMD [70–72], which results in the increased presence of drusen in Bruch's membrane [73]. A major component of drusen is vitronectin and complement proteins [25]. It has previously been shown that vitronectin is upregulated during complement attack on RPE cells [74,75]. Since vitronectin is a tyrosine-sulfated protein, it would be interesting to study if eliminating tyrosine sulfation decreases its binding to the MAC complex and results in its inability to inhibit the complement cascade, therefore exacerbating complement damage to cells. Future studies in which tyrosine-sulfated sites on vitronectin are mutated to phenylalanines will address the role of tyrosine sulfation in MAC binding.

Sulfinator analysis of the RPE secretome predicted CFH as a tyrosine-sulfated protein. This protein is produced by the RPE [20,62] and is also involved in complement regulation. CFH inhibits complement activation by acting as a cofactor in Factor I-mediated decay of C3-covertase [76]. Mutations in CFH have been implicated in AMD [27,29,30]. In addition to binding C3 convertase, CFH self-interacts [81,82] and has also been shown to bind multiple proteins in the complement system, such as C-reactive protein (CRP) [77,78], C3b [79], and C3d [80,80]. Since the main function of tyrosine sulfation is modulation of protein-protein interaction [6,22–24], the function of tyrosine sulfation in CFH may be to influence the interactions between some of these complement proteins and to inhibit complement activation. To test this possibility, future studies will focus on the identification of the sulfated tyrosine(s) on CFH using site-directed mutagenesis to phenylalanines, followed by functional studies to examine if the interaction of the protein with its binding partner is affected.

Ultimately, we would generate knock-in mice that have the endogenous wild-type sulfated tyrosine(s) replaced with phenylalanine(s). This mouse will be the ideal model with which to better understand the role that tyrosine sulfation plays in the function of CFH in complement regulation.

Due to the involvement of tyrosine sulfation in protein-protein interactions [6,22–24], using the PSG2 immunoaffinity column to isolate tyrosine *O*-sulfated proteins would not only pull down tyrosine-sulfated proteins, but also nonsulfated proteins which may be co-purifying due to their direct or indirect interaction with tyrosine-sulfated proteins. Therefore, each protein isolated from the column needs to be independently verified as either a tyrosine-sulfated protein or an interacting protein.

In conclusion, this report identifies the tyrosine-sulfated proteins present in the RPE. Since tyrosine sulfation can influence protein-

protein interactions, any future studies of vitronectin, opticin, or CFH should involve identifying the tyrosine-sulfation sites on these proteins. This will be followed by functional studies to examine if the interaction of the protein with its binding partner is affected.

Acknowledgments

We are grateful for the technical assistance provided by Mr. Robert Hamilton.

Author Contributions

Conceived and designed the experiments: YK MRA. Performed the experiments: YK JCS MK. Analyzed the data: YK MRA MK. Contributed reagents/materials/analysis tools: MK. Contributed to the writing of the manuscript: YK MRA.

References

- Moore KL (2003) The biology and enzymology of protein tyrosine *O*-sulfation. *J Biol Chem* 278: 24243–24246.
- Huttner WB (1984) Determination and occurrence of tyrosine *O*-sulfate in proteins. *Methods Enzymol* 107: 200–223.
- Stone MJ, Chuang S, Hou X, Shoham M, Zhu JZ (2009) Tyrosine sulfation: an increasingly recognised post-translational modification of secreted proteins. *N Biotechnol* 25: 299–317.
- Baeuerle PA, Huttner WB (1987) Tyrosine sulfation is a trans-Golgi-specific protein modification. *J Cell Biol* 105: 2655–2664.
- Huttner WB (1988) Tyrosine sulfation and the secretory pathway. *Annu Rev Physiol* 50: 363–376.
- Zhu JZ, Millard CJ, Ludeman JP, Simpson LS, Clayton DJ, et al (2011) Tyrosine sulfation influences the chemokine binding selectivity of peptides derived from chemokine receptor CCR3. *Biochemistry* 50: 1524–1534.
- Liu ZJ, Yang YJ, Jiang L, Xu YC, Wang AX, et al (2011) Tyrosine sulfation in N-terminal domain of human C5a receptor is necessary for binding of chemotaxis inhibitory protein of *Staphylococcus aureus*. *Acta Pharmacol Sin* 32: 1038–1044.
- Choe H, Farzan M (2009) Chapter 7. Tyrosine sulfation of HIV-1 coreceptors and other chemokine receptors. *Methods Enzymol* 461: 147–170.
- Baeuerle PA, Lottspeich F, Huttner WB (1988) Purification of yolk protein 2 of *Drosophila melanogaster* and identification of its site of tyrosine sulfation. *J Biol Chem* 263: 14925–14929.
- Nicholas HB Jr, Chan SS, Rosenquist GL (1999) Reevaluation of the determinants of tyrosine sulfation. *Endocrine* 11: 285–292.
- Westmuckett AD, Hoffhines AJ, Borghei A, Moore KL (2008) Early postnatal pulmonary failure and primary hypothyroidism in mice with combined TPST-1 and TPST-2 deficiency. *Gen Comp Endocrinol* 156: 145–153.
- Ouyang YB, Crawley JT, Aston CE, Moore KL (2002) Reduced body weight and increased postimplantation fetal death in tyrosylprotein sulfotransferase-1-deficient mice. *J Biol Chem* 277: 23781–23787.
- Borghei A, Ouyang YB, Westmuckett AD, Marcello MR, Landel CP, et al (2006) Targeted disruption of tyrosylprotein sulfotransferase-2, an enzyme that catalyzes post-translational protein tyrosine *O*-sulfation, causes male infertility. *J Biol Chem* 281: 9423–9431.
- Sherry DM, Kanan Y, Hamilton R, Hoffhines A, Arbogast KL, et al (2012) Differential developmental deficits in retinal function in the absence of either protein tyrosine sulfotransferase-1 or -2. *PLoS One* 7: e39702.
- Sherry DM, Murray AR, Kanan Y, Arbogast KL, Hamilton RA, et al (2010) Lack of protein-tyrosine sulfation disrupts photoreceptor outer segment morphogenesis, retinal function and retinal anatomy. *Eur J Neurosci* 32: 1461–1472.
- Kanan Y, Hamilton RA, Moore KL, Al-Ubaidi MR (2012) Protein tyrosine-*O*-sulfation in bovine ocular tissues. *Adv Exp Med Biol* 723: 835–841.
- Kanan Y, Hoffhines A, Rauhauser A, Murray A, Al-Ubaidi MR (2009) Protein tyrosine-*O*-sulfation in the retina. *Exp Eye Res* 89: 559–567.
- Hoffhines AJ, Jen CH, Leary JA, Moore KL (2009) Tyrosylprotein sulfotransferase-2 expression is required for sulfation of RNase 9 and Mfge8 *in vivo*. *J Biol Chem* 284: 3096–3105.
- Hoffhines AJ, Damoc E, Bridges KG, Leary JA, Moore KL (2006) Detection and purification of tyrosine-sulfated proteins using a novel anti-sulfotyrosine monoclonal antibody. *J Biol Chem* 281: 37877–37887.
- An E, Lu X, Flippin J, Devaney JM, Halligan B, et al (2006) Secreted proteome profiling in human RPE cell cultures derived from donors with age related macular degeneration and age matched healthy donors. *J Proteome Res* 5: 2599–2610.
- Monigatti F, Gasteiger E, Bairoch A, Jung E (2002) The Sulfinator: predicting tyrosine sulfation sites in protein sequences. *Bioinformatics* 18: 769–770.
- Rodgers SD, Camphausen RT, Hammer DA (2001) Tyrosine sulfation enhances but is not required for PSGL-1 rolling adhesion on P-selectin. *Biophys J* 81: 2001–2009.
- Nishimura Y, Wakita T, Shimizu H (2010) Tyrosine sulfation of the amino terminus of PSGL-1 is critical for enterovirus 71 infection. *PLoS Pathog* 6: e1001174.
- Costagliola S, Panneels V, Bonomi M, Koch J, Many MC, et al (2002) Tyrosine sulfation is required for agonist recognition by glycoprotein hormone receptors. *EMBO J* 21: 504–513.
- Hageman GS, Mullins RF, Russell SR, Johnson LV, Anderson DH (1999) Vitronectin is a constituent of ocular drusen and the vitronectin gene is expressed in human retinal pigmented epithelial cells. *FASEB J* 13: 477–484.
- Wyatt MK, Tsai JY, Mishra S, Campos M, Jaworski C, et al (2013) Interaction of complement factor h and fibulin3 in age-related macular degeneration. *PLoS One* 8: e68088.
- Edwards AO, Ritter R III, Abel KJ, Manning A, Panhuysen C, et al (2005) Complement factor H polymorphism and age-related macular degeneration. *Science* 308: 421–424.
- Hageman GS, Anderson DH, Johnson LV, Hancox LS, Taiber AJ, et al (2005) A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A* 102: 7227–7232.
- Haines JL, Hauser MA, Schmidt S, Scott WK, Olson LM, et al (2005) Complement factor H variant increases the risk of age-related macular degeneration. *Science* 308: 419–421.
- Klein RJ, Zeiss C, Chew EY, Tsai JY, Sackler RS, et al (2005) Complement factor H polymorphism in age-related macular degeneration. *Science* 308: 385–389.
- Le Goff MM, Lu H, Ugarte M, Henry S, Takanosu M, et al (2012) The vitreous glycoprotein opticin inhibits preretinal neovascularization. *Invest Ophthalmol Vis Sci* 53: 228–234.
- Le Goff MM, Sutton MJ, Slevin M, Latif A, Humphries MJ, et al (2012) Opticin exerts its anti-angiogenic activity by regulating extracellular matrix adhesiveness. *J Biol Chem* 287: 28027–28036.
- Sanders EJ, Walter MA, Parker E, Aramburo C, Harvey S (2003) Opticin binds retinal growth hormone in the embryonic vitreous. *Invest Ophthalmol Vis Sci* 44: 5404–5409.
- Kanan Y, Matsumoto H, Song H, Sokolov M, Anderson RE, et al (2010) Serine/threonine kinase akt activation regulates the activity of retinal serine/threonine phosphatases, PHLPP and PHLPL. *J Neurochem* 113: 477–488.
- Rose JK, Doms RW (1988) Regulation of protein export from the endoplasmic reticulum. *Annu Rev Cell Biol* 4: 257–288.
- Budarf ML, Herbert E (1982) Effect of tunicamycin on the synthesis, processing, and secretion of pro-opiomelanocortin peptides in mouse pituitary cells. *J Biol Chem* 257: 10128–10135.
- Onnerfjord P, Heathfield TF, Heinegard D (2004) Identification of tyrosine sulfation in extracellular leucine-rich repeat proteins using mass spectrometry. *J Biol Chem* 279: 26–33.
- Farrell DH, Mulvihill ER, Huang SM, Chung DW, Davie EW (1991) Recombinant human fibrinogen and sulfation of the gamma' chain. *Biochemistry* 30: 9414–9420.
- Hirose S, Oda K, Ikehara Y (1988) Tyrosine *O*-sulfation of the fibrinogen gamma B chain in primary cultures of rat hepatocytes. *J Biol Chem* 263: 7426–7430.
- Krajewski T, Blomback B (1968) The location of tyrosine-*O*-sulphate in fibrinopeptides. *Acta Chem Scand* 22: 1339–1346.
- Hortin G, Tollefsen DM, Strauss AW (1986) Identification of two sites of sulfation of human heparin cofactor II. *J Biol Chem* 261: 15827–15830.
- Hortin G, Sims H, Strauss AW (1986) Identification of the site of sulfation of the fourth component of human complement. *J Biol Chem* 261: 1786–1793.
- Kanan Y, Brobst D, Han Z, Naash MI, Al-Ubaidi MR (2014) Fibulin 2, a tyrosine *O*-sulfated protein, is up-regulated following retinal detachment. *J Biol Chem*. 'In press'.

44. Pepperberg DR, Okajima TL, Wiggert B, Ripps H, Crouch RK, et al (1993) Interphotoreceptor retinoid-binding protein (IRBP). Molecular biology and physiological role in the visual cycle of rhodopsin. *Mol Neurobiol* 7: 61–85.
45. Caraci F, Busceti C, Biagioni F, Aronica E, Mastroiacovo F, et al (2008) The Wnt antagonist, Dickkopf-1, as a target for the treatment of neurodegenerative disorders. *Neurochem Res* 33: 2401–2406.
46. Morse EE (1971) Fibrinogen and the fibrinogenopathies. *Ann Clin Lab Sci* 1: 155–161.
47. Jenne D, Hille A, Stanley KK, Huttner WB (1989) Sulfation of two tyrosine-residues in human complement S-protein (vitronectin). *Eur J Biochem* 185: 391–395.
48. Yu Y, Hoffhines AJ, Moore KL, Leary JA (2007) Determination of the sites of tyrosine O-sulfation in peptides and proteins. *Nat Methods* 4: 583–588.
49. Hintner H, Dahlback K, Dahlback B, Pepys MB, Breathnach SM (1991) Tissue vitronectin in normal adult human dermis is non-covalently bound to elastic tissue. *J Invest Dermatol* 96: 747–753.
50. Pohl WR, Conlan MG, Thompson AB, Ertl RF, Romberger DJ, et al (1991) Vitronectin in bronchoalveolar lavage fluid is increased in patients with interstitial lung disease. *Am Rev Respir Dis* 143: 1369–1375.
51. Seger D, Shaltiel S (2000) Evidence showing that the two-chain form of vitronectin is produced in the liver by a selective furin cleavage. *FEBS Lett* 480: 169–174.
52. Choi Y, Kim E, Lee Y, Han MH, Kang IC (2010) Site-specific inhibition of integrin alpha v beta 3-vitronectin association by a ser-asp-val sequence through an Arg-Gly-Asp-binding site of the integrin. *Proteomics* 10: 72–80.
53. Monfort J, Tardif G, Roughley P, Reboul P, Boileau C, et al (2008) Identification of opticin, a member of the small leucine-rich repeat proteoglycan family, in human articular tissues: a novel target for MMP-13 in osteoarthritis. *Osteoarthritis Cartilage* 16: 749–755.
54. Ramesh S, Bonshek RE, Bishop PN (2004) Immunolocalisation of opticin in the human eye. *Br J Ophthalmol* 88: 697–702.
55. Ma J, Zhu TP, Moe MC, Ye P, Yao K (2012) Opticin production is reduced by hypoxia and VEGF in human retinal pigment epithelium via MMP-2 activation. *Cytokine* 59: 100–107.
56. Le Goff MM, Hindson VJ, Jowitt TA, Scott PG, Bishop PN (2003) Characterization of opticin and evidence of stable dimerization in solution. *J Biol Chem* 278: 45280–45287.
57. Annunen S, Korkko J, Czarny M, Warman ML, Brunner HG, et al (1999) Splicing mutations of 54-bp exons in the COL11A1 gene cause Marshall syndrome, but other mutations cause overlapping Marshall/Stickler phenotypes. *Am J Hum Genet* 65: 974–983.
58. Richards AJ, Yates JR, Williams R, Payne SJ, Pope FM, et al (1996) A family with Stickler syndrome type 2 has a mutation in the COL11A1 gene resulting in the substitution of glycine 97 by valine in alpha 1 (XI) collagen. *Hum Mol Genet* 5: 1339–1343.
59. Boon CJ, Klevering BJ, Hoyng CB, Zonneveld-Vrieling MN, Nabuurs SB, et al (2008) Basal laminar drusen caused by compound heterozygous variants in the CFH gene. *Am J Hum Genet* 82: 516–523.
60. Schwaeble W, Zwirner J, Schulz TF, Linke RP, Dierich MP, et al (1987) Human complement factor H: expression of an additional truncated gene product of 43 kDa in human liver. *Eur J Immunol* 17: 1485–1489.
61. Morris KM, Aden DP, Knowles BB, Colten HR (1982) Complement biosynthesis by the human hepatoma-derived cell line HepG2. *J Clin Invest* 70: 906–913.
62. Chen M, Forrester JV, Xu H (2007) Synthesis of complement factor H by retinal pigment epithelial cells is down-regulated by oxidized photoreceptor outer segments. *Exp Eye Res* 84: 635–645.
63. Fenaile F, Le MM, Groseil C, Ramon C, Riande S, et al (2007) Site-specific N-glycan characterization of human complement factor H. *Glycobiology* 17: 932–944.
64. Muthana SM, Campbell CT, Gildersleeve JC (2012) Modifications of glycans: biological significance and therapeutic opportunities. *ACS Chem Biol* 7: 31–43.
65. Vogel BE, Lee SJ, Hildebrand A, Craig W, Pierschbacher MD, et al (1993) A novel integrin specificity exemplified by binding of the alpha v beta 5 integrin to the basic domain of the HIV Tat protein and vitronectin. *J Cell Biol* 121: 461–468.
66. Pierschbacher MD, Ruoslahti E (1987) Influence of stereochemistry of the sequence Arg-Gly-Asp-Xaa on binding specificity in cell adhesion. *J Biol Chem* 262: 17294–17298.
67. Ramachandran V, Nollert MU, Qiu H, Liu WJ, Cummings RD, et al (1999) Tyrosine replacement in P-selectin glycoprotein ligand-1 affects distinct kinetic and mechanical properties of bonds with P- and L-selectin. *Proc Natl Acad Sci U S A* 96: 13771–13776.
68. Ruoslahti E (1996) RGD and other recognition sequences for integrins. *Annu Rev Cell Dev Biol* 12: 697–715.
69. Podack ER, Preissner KT, Muller-Eberhard HJ (1984) Inhibition of C9 polymerization within the SC5b-9 complex of complement by S-protein. *Acta Pathol Microbiol Immunol Scand Suppl* 284: 89–96.
70. Sparrow JR, Ueda K, Zhou J (2012) Complement dysregulation in AMD: RPE-Bruch's membrane-choroid. *Mol Aspects Med* 33: 436–445.
71. Bradley DT, Zipfel PF, Hughes AE (2011) Complement in age-related macular degeneration: a focus on function. *Eye (Lond)* 25: 683–693.
72. Karagianni N, Adamis AP (2010) The case for complement and inflammation in AMD: open questions. *Adv Exp Med Biol* 703: 1–7.
73. Pauleikhoff D, Chen J, Bird AC, Wessing A (1992) [The Bruch membrane and choroid. Angiography and functional characteristics in age-related changes]. *Ophthalmologie* 89: 39–44.
74. Lueck K, Wasmuth S, Williams J, Hughes TR, Morgan BP, et al (2011) Sub-lytic C5b-9 induces functional changes in retinal pigment epithelial cells consistent with age-related macular degeneration. *Eye (Lond)* 25: 1074–1082.
75. Wasmuth S, Lueck K, Bachler H, Lommatzsch A, Pauleikhoff D (2009) Increased vitronectin production by complement-stimulated human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci* 50: 5304–5309.
76. Pechtl IC, Kavanagh D, McIntosh N, Harris CL, Barlow PN (2011) Disease-associated N-terminal complement factor H mutations perturb cofactor and decay-accelerating activities. *J Biol Chem* 286: 11082–11090.
77. Perkins SJ, Okemefuna AI, Nan R (2010) Unravelling protein-protein interactions between complement factor H and C-reactive protein using a multidisciplinary strategy. *Biochem Soc Trans* 38: 894–900.
78. Okemefuna AI, Nan R, Miller A, Gor J, Perkins SJ (2010) Complement factor H binds at two independent sites to C-reactive protein in acute phase concentrations. *J Biol Chem* 285: 1053–1065.
79. Schmidt CQ, Herbert AP, Kavanagh D, Gandy C, Fenton CJ, et al (2008) A new map of glycosaminoglycan and C3b binding sites on factor H. *J Immunol* 181: 2610–2619.
80. Bhattacharjee A, Lehtinen MJ, Kajander T, Goldman A, Jokiranta TS (2010) Both domain 19 and domain 20 of factor H are involved in binding to complement C3b and C3d. *Mol Immunol* 47: 1686–1691.
81. Fernando AN, Furtado PB, Clark SJ, Gilbert HE, Day AJ, et al (2007) Associative and structural properties of the region of complement factor H encompassing the Tyr402His disease-related polymorphism and its interactions with heparin. *J Mol Biol* 368: 564–581.
82. Okemefuna AI, Gilbert HE, Griggs KM, Ormsby RJ, Gordon DL, et al (2008) The regulatory SCR-1/5 and cell surface-binding SCR-16/20 fragments of factor H reveal partially folded-back solution structures and different self-associative properties. *J Mol Biol* 375: 80–101.