



## Correction

# Correction: Fenofibrate Improves Renal Lipotoxicity through Activation of AMPK-PGC-1 $\alpha$ in *db/db* Mice

## The PLOS ONE Staff

There is an error in the fourth sentence of the “Experimental methods” section of the Materials and Methods. The corrected sentence should read: In total, 20 mg/kg/day of fenofibrate was administered to the treated *db/db* and *db/m* groups.

There are errors in the sixth sentence of the second paragraph of the Discussion. The corrected sentence should read: Recently, we demonstrated that the AMPK activator resveratrol improves lipotoxicity in the kidney by enhancing the PPAR $\alpha$ -ERR-1 $\alpha$ -SREBP-1-mediated removal of lipid accumulation and decreasing apoptotic renal cell death and oxidative stress related to FoxO3a dephosphorylation in diabetic nephropathy.

## Reference

1. Hong YA, Lim JH, Kim MY, Kim TW, Kim Y, et al. (2014) Fenofibrate Improves Renal Lipotoxicity through Activation of AMPK-PGC-1 $\alpha$  in *db/db* Mice. PLoS ONE 9(5): e96147. doi:10.1371/journal.pone.0096147

**Citation:** The PLOS ONE Staff (2014) Correction: Fenofibrate Improves Renal Lipotoxicity through Activation of AMPK-PGC-1 $\alpha$  in *db/db* Mice. PLoS ONE 9(8): e105921. doi:10.1371/journal.pone.0105921

**Published:** August 11, 2014

**Copyright:** © 2014 The PLOS ONE Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.