

CORRECTION

# Correction: Caco-2 Cell Acquisition of Dietary Iron(III) Invokes a Nanoparticulate Endocytic Pathway

The PLOS ONE Staff

There is an error in the first sentence of the “Cellular uptake studies” subsection of the Materials and Methods. The correct sentence is: To avoid aggregation/agglomeration of the nanoparticulate iron, the medium for cellular uptake consisted of a balanced salt solution (BSS) containing 130 mM NaCl, 10 mM KCl, 1 mM MgSO<sub>4</sub>, 5 mM Glucose and 1.8 mM CaCl<sub>2</sub> in 10 mM PIPES buffer (pH 7.4).

## Reference

1. Pereira DIA, Mergler BI, Faria N, Bruggraber SFA, Aslam MF, Poots LK, et al. (2013) Caco-2 Cell Acquisition of Dietary Iron(III) Invokes a Nanoparticulate Endocytic Pathway. PLoS ONE 8(11): e81250. doi:[10.1371/journal.pone.0081250](https://doi.org/10.1371/journal.pone.0081250) PMID: [24278403](https://pubmed.ncbi.nlm.nih.gov/24278403/)



## OPEN ACCESS

**Citation:** The PLOS ONE Staff (2015) Correction: Caco-2 Cell Acquisition of Dietary Iron(III) Invokes a Nanoparticulate Endocytic Pathway. PLoS ONE 10(3): e0119747. doi:[10.1371/journal.pone.0119747](https://doi.org/10.1371/journal.pone.0119747)

**Published:** March 20, 2015

**Copyright:** © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.