

RESEARCH ARTICLE

Social Stress Increases Cortisol and Hampers Attention in Adolescents with Excess Weight

Antonio Verdejo-Garcia^{1,2*}, Maria Moreno-Padilla¹, M. Carmen Garcia-Rios³, Francisca Lopez-Torrecillas¹, Elena Delgado-Rico¹, Jacqueline Schmidt-Rio-Valle³, Maria J. Fernandez-Serrano⁴

1 Department of Clinical Psychology and Institute of Neuroscience F. Oloriz, Universidad de Granada, Granada, Spain, **2** School of Psychological Sciences, Monash University, Melbourne, Australia, **3** School of Health Sciences, Universidad de Granada, Granada, Spain, **4** Department of Psychology, Universidad de Jaen, Jaen, Spain

* antonio.verdejo@monash.edu



Abstract

Objective

To experimentally examine if adolescents with excess weight are more sensitive to social stress and hence more sensitive to harmful effects of stress in cognition.

Design and Methods

We conducted an experimental study in 84 adolescents aged 12 to 18 years old classified in two groups based on age adjusted Body Mass Index percentile: Normal weight (n=42) and Excess weight (n=42). Both groups were exposed to social stress as induced by the virtual reality version of the Trier Social Stress Task –participants were requested to give a public speech about positive and negative aspects of their personalities in front of a virtual audience. The outcome measures were salivary cortisol levels and performance in cognitive tests before and after the social stressor. Cognitive tests included the CANTAB Rapid Visual Processing Test (measuring attention response latency and discriminability) and the Iowa Gambling Task (measuring decision-making).

Results

Adolescents with excess weight compared to healthy weight controls displayed increased cortisol response and less improvement of attentional performance after the social stressor. Decision-making performance decreased after the social stressor in both groups.

Conclusion

Adolescents who are overweight or obese have increased sensitivity to social stress, which detrimentally impacts attentional skills.

OPEN ACCESS

Citation: Verdejo-Garcia A, Moreno-Padilla M, Garcia-Rios MC, Lopez-Torrecillas F, Delgado-Rico E, Schmidt-Rio-Valle J, et al. (2015) Social Stress Increases Cortisol and Hampers Attention in Adolescents with Excess Weight. PLoS ONE 10(4): e0123565. doi:10.1371/journal.pone.0123565

Academic Editor: Suliann Ben Hamed, Centre de Neuroscience Cognitive, FRANCE

Received: October 23, 2014

Accepted: March 4, 2015

Published: April 21, 2015

Copyright: © 2015 Verdejo-Garcia et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study has been funded by grants PSI2010-17290 (INTEROBE) from the Ministry of Innovation and Science (MICINN), and P-10-HUM-6635 (NEUROCOBE).

Competing Interests: The authors have declared that no competing interests exist.

Introduction

Adolescents with excess weight suffer substantial social stress including frequent peer bullying and social marginalization and exclusion [1,2]. Crucially, the degree of exposure to these social stressors is the most important predictor of poor psychological adjustment and poor academic achievement in adolescents with obesity [3]. Moreover, neuroendocrine studies have shown that non-fasting levels of the “hunger hormone” ghrelin increase in response to social stressors (i.e., the Trier Social Stress Task, involving a public speak) [4] and that the awakening response of the “stress hormone” cortisol positively associates with subsequent lipid intake [5]. Therefore, social stress is a potent determinant of poor cognition and poor food choices in adolescents with excess weight. This phenomenon could be explained by the harmful impact of social stress on cognitive skills such as attention, cognitive control and decision-making, which contribute to obesity-related behaviours in adolescents [6]. The harmful impact of persistent social stressors on cognition in adolescents with obesity is likely to be enduring as stress induces neuroadaptations in prefrontal and limbic regions particularly during adolescence [7,8]. Therefore, examining whether social stress hampers cognition in adolescents with excess weight is essential for prevention of cognitive decline and hence progression of obesity. However, to date no studies have experimentally assessed this notion. In this study we examined if a social stressor—the Trier public speaking stress task—specifically increases cortisol levels and hampers cognitive performance in adolescents with excess weight compared to adolescents with normal weight. We specifically assessed the impact of social stress on outcome measures of attention, cognitive inhibition and decision-making. We selected these outcomes because they reflect the function of frontal-limbic systems [9,10] and are longitudinally associated with weight gain in pediatric populations [11,12]. We hypothesized that adolescents with excess weight would show greater cortisol response to the social stressor, and greater detrimental impact of social stress on attention and decision-making performance.

Methods

Participants

Eighty-four adolescents aged between 12 and 18 years old participated in the study. They were classified in two groups (Normal weight [$n = 42$] and Excess weight [$n = 42$]) based on their age adjusted Body Mass Index (BMI) percentile [13]. Sample size was estimated through power analysis. The existing evidence about the impact of the Trier Social Stress Task (TSST) on selected outcome variables was correlational (i.e., the association between TSST-induced cortisol changes and decision-making performance is between 0.3 and 0.4) [14,15]. Therefore, we estimated that in order to achieve adequate power (80%) to detect a $\rho_{H1} = 0.3$ association between the independent variable (stress) and the cognitive outcomes (attention and decision-making) 84 participants would be required (S1 Fig). This sample size was deemed acceptable for the mixed repeated-measures design. The classification of the two groups was conducted in alignment with the guidelines of the International Obesity Task Force and the Centers for Disease Control and Prevention: Normal weight participants had age adjusted BMI percentiles in the range between the 5th and the 84th percentile, and Excess weight participants had age adjusted BMI percentiles ≥ 85 (Table 1). Three participants from the Excess weight group provided invalid cortisol samples, and therefore the final study sample comprised 42 Normal weight and 39 Excess weight participants. Participants' socio-demographic characteristics, BMIs, percentage fat and blood count obtained biochemical parameters are as well displayed in Table 1. Participants also completed The Dutch Eating Behavior Questionnaire [16] which was used to

Table 1. Descriptive scores for the demographic, biometric and blood count characteristics of adolescents with excess and normal weight.

Variables	Excess weight		Normal weight		t ^a /chi square ^b	p
	Mean	SD	Mean	SD		
Age	15.59	1.91	15.62	1.83	-.07 ^a	.944
Gender (% Men/Women)	52.4/47.6		43.2/56.8		.72 ^b	.262
BMI	29.87	3.57	20.87	2.06	13.73 ^a	.000
Fat (%)	31.97	9.15	17.99	6.94	7.69 ^a	.000
DEBQ						
Emotional	23.68	9.33	24.03	9.02	-0.17 ^a	.858
External	28.64	7.11	31.28	7.68	-1.71 ^a	.091
Restraint	25.55	7.22	19.30	7.81	3.98 ^a	.000
Glucose	92.57	6.03	92.14	6.64	.276 ^a	.783
Cholesterol	158.07	27.35	148.47	20.89	1.64 ^a	.104
Triglycerides	70.64	28.76	63.13	27.32	1.09 ^a	.279
HDL	56.80	12.49	58.73	12.82	-.61 ^a	.541
LDL	90.85	21.08	80.51	14.96	2.38 ^a	.020
Insulin	47.28	57.78	53.34	114.18	-.25 ^a	.802
Uric Acid	5.08	0.87	4.39	0.97	2.98 ^a	.004
Thyroxine	1.33	0.43	1.44	0.68	-0.79 ^a	.451

^avalue of Student's *t*;

^bvalue of Chi-square χ^2

doi:10.1371/journal.pone.0123565.t001

characterise psychological traits relevant to maladaptive eating behaviours (i.e., external eating, emotional eating and restraint) (Table 1).

Participants were recruited from the paediatrics and endocrinology services of the Hospital “Virgen de las Nieves” in Granada (Spain), and from schools located in the same geographical area. The inclusion criteria for participants were defined as follows: (i) age range between 12 and 18 years old; (ii) BMI percentiles falling within the intervals categorized as overweight or obesity (≥ 85 —Excess weight group), or normal weight (5–85—Normal weight group); and (iii) absence of history or current evidence of neurological or psychiatric disorders, assessed by participants and parents interviews and the Eating Disorder Inventory [17]. All participants had normal or corrected-to-normal vision.

Experimental procedures

Fig 1 displays a schematic representation of the experiment. In order to induce social stress in the laboratory we utilised a previously validated Virtual Reality version of the Trier Social Stress Task (TSST) [18]. Participants had to perform a stressing task which consisted of delivering a speech about personal characteristics including both positive and negative aspects of themselves in front of a simulated audience. Participants were told that this audience would attend the speech and subsequently evaluate its quality. However, the virtual audience was programmed to look progressively bored and disappointed with the speech. The speech was followed by a mental calculation test (serially subtracting 17, starting from 2013). Cortisol levels were measured via saliva samples collected before onset of the TSST (T1), after completion of the TSST and the calculation test (10 minutes after TSST onset—T2) and after performance on each of the attention and decision making cognitive probes (20 and 30 minutes after TSST onset-T3 and T4- respectively). Cognitive measures were conducted in a fixed order before TSST onset (pre-TSST, overlapping with T1) and after completion of the TSST and the



Fig 1. Schematic representation of the experiment.

doi:10.1371/journal.pone.0123565.g001

calculation test (post-TSST, overlapping with T2). To minimize practice effects, we utilised parallel versions of all tasks in the post-TSST administration. The original validation study showed that this virtual reality TSST is able to induce modest but sizeable increases in cortisol and subjective stress responses [18]. Moreover the virtual audience tamed the ethical concerns associated with the negative impact of the social stressor on adolescents' participants. The Ethics Committee for Human Research of the Universidad de Granada approved the study. Both participants and parents signed informed consent.

Cortisol measurement. Participants were told not to smoke, eat or drink coffee for at least 30 minutes before the experiment. All the experimental sessions were conducted at the same time of the day (4–5 pm) based on pilot data obtained in this cohort prior to study onset indicating that diurnal cortisol levels were stable during these hours. Saliva was collected via a commercially available device: Salivette Cortisol (Sarstedt, Numbrecht, Germany). This device is composed of a cotton tube (similar to dental cotton), and two plastic tubes that fit one inside the other. Subjects were told to place cotton salivettes inside their mouth and gently chew and/or suck on them for 1–3 min until they became soaked in saliva. The cotton tube was inserted inside the plastic tube, which was then capped. Saliva samples were stored at -20°C until required for assay. Samples were analyzed at the University Hospital, using the electrochemiluminescence immunoassay “ECLIA” method. This method is designed for use in Roche Elecsys 1010/2010 automated analyzers and in the Elecsys MODULAR NALYTICS E170 module. We computed two different metrics from each cortisol sample (microgram/deciliter and nanomol/liter). The correlation between both metrics at the different time points ranged from 0.8 and 0.9.

Cognitive measures. We utilized three computerized tests: two subtests from the Cambridge Neuropsychological Test Automated Battery (CANTAB) [19], *Motor Screening (MOT)* and *Rapid Visual Information Processing (RVP)*, and the Iowa Gambling Task (IGT) [20]. Alternate versions of each test were used in pre-stress and post-stress administrations.

MOT. The main objective of this test is to provide a baseline measure of the subjects' basic motor skills in terms of reaction times and accuracy. After a demonstration of the correct way to point on the computer screen using the forefinger of the dominant hand, the subjects must point to a series of stimuli (crosses) popping up in turn. The outcome measure of this test was response latency.

RVP. This is a test of visual sustained attention with an impulse control component. A white box is displayed in the centre of the computer screen, inside which digits, from 2 to 9, are displayed in a pseudo-random order, at the rate of 100 digits per minute. The subject must detect consecutive odd or even sequences of digits (for example, 2-4-6) and respond by pressing the touch pad. The outcome measures of this test were response latency and response

discriminability (B') scores, which are sensitive to attention and impulse control domains respectively. The B' score is the signal detection measure of the strength of trace required to elicit a response (range -1.00 to +1.00). Thus, it is the tendency to respond regardless of whether the target sequence is present and uses the $p(\text{hit})$ and $p(\text{fa})$ results. A score close to +1.00 indicates that the subject gave few false alarms.

IGT. This is a computer task measuring reward/punishment based decision-making. It involves four decks of cards (A, B, C and D). Each time a participant selects a card, a specified amount of play money is awarded. However, interspersed among these rewards, there are probabilistic punishments (monetary losses). Two of the decks of cards (A and B) produce high immediate gains; however, in the long run, they will take more money than they give, and are thus considered disadvantageous. The other two decks (C and D) are considered advantageous, as they result in small, immediate gains, but will yield more money than they take in the long run. The performance measure was the net score calculated by subtracting the number of disadvantageous choices (decks A and B) from the number of advantageous choices (decks C and D). An equivalent parallel version of the ABCD task in which decks are labelled K, L, M and N was utilised in the post-TSST administration. These versions have shown adequate test-retest reliability and ecological validity in relation to decision-making [21].

Visual Analogue Scales (VAS). We used two Visual Analogue Scales (VAS) designed to rate arousal and stress. For arousal scale the individual must indicate the extent to which they perceived as active and alert (from nothing active to very active). For stress scale they must indicate how much stress they feel (from no stress to very much stress). We used the mean scores of each dimension.

Statistical analyses

The main hypotheses were examined utilizing mixed repeated measures analyses of variance including Time as the repeated-measures factor, Group as the between-groups factor, and cortisol levels (as measured in $\mu\text{g/dl}$) and RVP's mean response latency and B' scores and IGT's net scores as dependent measures. Cortisol and RVP performance measures were log-transformed (base 10) to meet the normal distribution, but for the sake of clarity the Figures report non-transformed measures. IGT scores fitted to the normal distribution as assessed by Kolmogorov-Smirnov tests. We also performed correlation analyses between change scores of cortisol levels ($T2-T1$) and change scores of cognitive performance ($T2-T1$) and between both change scores and biological and psychological measures. These change measures were non-normally distributed and therefore we applied Spearman's rank correlation analyses. Two participants from the Excess weight group ($n = 37$) and one participant from the Normal weight group had missing cortisol data at T1 and T2 ($n = 41$). With regard to cognitive tests, there was no missing data in the Excess weight group ($n = 39$), whereas in the Normal weight group three participants had invalid data for RVP response latency and IGT ($n = 39$) at T1 or T2, and three participants had invalid data for RVP B' ($n = 38$) at T1 or T2.

Results

Cortisol response

We found a significant Time x Group interaction on cortisol levels, $F(3,74) = 4.36$, $p = 0.008$. Cortisol mildly increased in Excess weight participants after the TSST. Independent-sample t -tests showed that Excess weight and Normal weight participants did not significantly differ on cortisol levels before TSST (T1). However, Excess weight adolescents showed significantly increased cortisol levels after TSST (T2), $t = 1.94$, $p = 0.05$, Cohen's $d = 0.5$ (Fig 2). Moreover, cortisol increase between T2 and T1 correlated with amount of fat, Spearman's $Rho = 0.30$,




Fig 2. Cortisol levels (µg/dl units) in adolescents with excess weight and adolescents with normal weight before and after exposure to the Trier Social Stress Task (TSST). T1 represents cortisol levels before TSST; T2 represents cortisol levels immediately after TSST termination; T3 and T4 represents cortisol levels 10 and 20 minutes after TSST termination.

doi:10.1371/journal.pone.0123565.g002

$p = 0.01$. Between-group differences were also statistically significant at T3, $t = 2.44$, $p = 0.02$, and T4, $t = 2.63$, $p = 0.01$. However, this effect seems to be driven by decreased cortisol levels in the Normal weight group (Fig 2).

Cognitive performance

MOT. Pre-TSST scores showed that both groups had similar baseline response latencies. Further, both groups showed mild reductions of response latencies between the pre-TSST measure and the post-TSST measure (Fig 3).

RVP—Response latency. We found a significant Time x Group interaction, $F(1,76) = 6.35$, $p = 0.01$ (Fig 3). Independent-sample t-tests showed that Excess weight and Normal weight participants did not significantly differ in the pre-TSST measure. However, they showed marginally significant differences in the post-TSST measure, $t(78) = 1.75$, $p = 0.08$, Cohen’s $d = 0.4$, with Excess weight participants performing significantly poorer than Normal weight controls. There was no significant correlation between T2—T1 cortisol levels and T2—T1 RVP Response Latency.

RVP—Response discriminability. We did not find a significant Time x Group interaction, $F(1,75) = 0.99$, $p = 0.32$. There were no main effects of Time or Group, although visual inspection shows Excess weight participants performed better than Normal weight participants in both pre- and post-TSST measures (Fig 3).

Decision-making—IGT. We did not find a significant Time x Group interaction, $F(1,77) = 0.005$, $p = 0.94$. There was a significant main effect of Time, $F(1,77) = 6.01$, $p = 0.02$, indicating that both groups exhibited significantly poorer performance after the TSST (Fig 3). There was no significant correlation between T2—T1 cortisol levels and T2—T1 IGT performance.

Correlations between biological and psychological measures and cognitive performance in T2—T1. We found a positive correlation between levels of uric acid and change in RVSP response latency performance between T2 and T1, Spearman’s $Rho = 0.46$, $p = 0.0001$, and a negative correlation between thyroxine levels and change in Iowa Gambling Task performance between T2 and T1, Spearman’s $Rho = -0.27$, $p = 0.03$. We also found a negative correlation




Fig 3. Cognitive performance in adolescents with excess weight and adolescents with normal weight before and after exposure to the Trier Social Stress Task (TSST). Top panel Y axes represent time in milliseconds. The Y axis in the bottom-left panel represents signal detection derived Beta scores, ranging from 0 to 1. The Y axis in the bottom-right panel represents Iowa Gambling Task net scores, ranging from -60 to +60.

doi:10.1371/journal.pone.0123565.g003

between scores of external eating and RVSP response latency performance between T2 and T1, Spearman's $Rho = -0.27$, $p = 0.02$.

Visual Analogue Scales (VAS). We did not find a significant Time x Group interactions on VAS of arousal or stress but results were in the expected direction, with both groups showing more subjective arousal and stress after the TSST (S2 Fig).

Post-hoc analyses in the subsample of participants showing enhanced cortisol response. The primary analyses indicated that in the normal weight group cortisol levels did not change after stress, and therefore there is a concern that cognitive changes were due to spurious factors. To address this issue, we run additional analyses in the subsample of participants who showed sizeable increments in cortisol levels after stress, including 24 participants of the Excess weight group (57% of the original sample) and 20 participants of the Normal weight group (48% of the original sample). The results of these analyses were coherent with the main findings. We found a significant Time x Group interaction on RVP's latency scores, $F(1,41) = 6.17$, $p = 0.02$, whereby a drop in performance was only observed in the Excess weight group (See S2 Fig). Moreover, there was a significant correlation between T2—T1 cortisol levels and T2—T1 RVP Response Latency (Spearman's $Rho = 0.25$, $p_{unilateral} = 0.05$) (Fig 4).

Discussion

We show that social stress specifically increases cortisol levels and hinders attentional response latency in adolescents with excess weight. Conversely, social stress failed to show significant effects on attention response discriminability. Moreover, both excess weight and normal weight adolescents displayed poorer decision-making performance after the social stressor. These




Fig 4. Correlation between between T2—T1 cortisol levels (X Axis) and T2—T1 RVP Response Latency (Y Axis) within the subsample of participants showing TSST-induced increases in cortisol levels.

doi:10.1371/journal.pone.0123565.g004

findings indicate that adolescents who are overweight and obese have enhanced stress reactivity in response to social stressors, which selectively impacts on attentional skills. Since adolescents with excess weight are markedly exposed to social stressors during everyday lives, our findings suggest that stress immunization strategies should be put in place to prevent the harmful impact of social stress on cognition and therefore on progression of obesity.

In agreement with our primary hypothesis, social stress induced greater cortisol response in overweight and obese adolescents. The effect was mild but the specific impact on participants with excess weight agrees with the notion that repetitive social stress may induce sensitization of the hypothalamic-pituitary-adrenal (HPA) axis [22] and purportedly of the HPA axis associations with fronto-limbic systems [23–25]. The discrepancy between our finding of cortisol increase and a previous negative finding in obese adults [26] suggests that adolescence compared to adulthood is a more sensitive time period for abnormal sensitization of stress systems, likely due to ongoing neural maturation of these systems [7,27]. Further, both preclinical and clinical evidence shows that social stressors such as social evaluation and social exclusion are particularly challenging for adolescents [8,28,29]. The potential mechanisms for the specific impact of social stress on stress reactivity in adolescents with excess weight include the additive or synergistic interactions between social stress and inflammation [30,31] and/or between social stress and obesity-related neuroadaptations in anterior cingulate and limbic regions that are essential for stress regulation [32,33]. Our finding is particularly relevant in view of the significant association between cortisol reactivity and obesity-related behaviours [34,35], and of the emerging evidence suggesting that high levels of stress can longitudinally predict the progression of obesity [36].

We also showed a significant impact of social stress on attentional performance in adolescents with excess weight. The effect was again mild and pointed to stress-related hindering of

the capacity to get benefit from a repeated administration of the task. Previous findings indicate that repeated administration of CANTAB attentional tests is associated with significant improvements in performance (of at least 0.3 in Cohen's *d* effect size) [37], and this is what we observed in the control group. However, excess weight adolescents were unable to get benefitted from this repeated administration. The effect was specific for attention-related latency adjustments, but not for psychomotor-related reaction times. Therefore, it suggests a detrimental impact of stress on attention regulation.⁶ This notion is consistent with the neural networks interactions between the HPA axis and medial prefrontal cortex and anterior cingulate cortex regions involved in attention regulation [38–40]. In support, neuroimaging studies have shown that the impact of stress on executive attention is mediated by structural (gray matter) neuroadaptations in prefrontal cortex and anterior cingulate cortex regions [41]. This stress-related attentional hurdle has a high translational value, as individual differences in response latencies to attentional probes are longitudinally associated with increases in BMI [12], implying that adequate control of social stress and/or cognitive boosting of attentional resources may contribute to prevent chronic obesity. This notion is consistent with our finding of significant correlations between less improvement of attentional performance (between T1 and T2) and higher maladaptive eating patterns such as external eating, which reflects attentional bias towards food related cues. Further, both social stress and attentional skills are significantly associated with advantageous social functioning and academic performance [3], and therefore our finding highlights the potential benefit of controlling social stress to improve social and career outcomes in the long-term.

Furthermore, we found poorer decision-making after the social stressor in both adolescents with excess weight and adolescents with normal weight. Since cortisol levels dropped between T3 and T4 (the time window of decision-making task performance) it is unlikely that this finding can be attributed to the effects of acute stress. However, it might be attributed to broader effects of the social stressor, such as the social evaluation context. The latter notion agrees with previous experimental evidence showing that adolescents make riskier choices than young adults or adults when they are under social evaluation [42]. The lack of specificity of our result implies that the impact of social evaluation on decision-making is mediated by neural mechanisms that are similarly sensitized in adolescents regardless of BMI/weight status, or that different neural mechanisms mediate a similar impact of social evaluation on decision-making in excess weight and normal weight adolescents. In favor of the first notion, neuroimaging studies have shown that the impact of social evaluation on decision-making is mediated by increased activation of ventral striatal and orbitofrontal regions [43], which are generally sensitized during adolescence. In favor of the second notion, we have observed that excess weight and normal weight adolescents recruit different brain circuitries during the pondering of social decisions [44]. Future studies are warranted to address this question. In any case, our finding might have general implications for prevention of obesity during adolescence as we know that adolescents who are overweight or obese have higher exposure to social evaluations [3] and that subsequent risky choices are longitudinally associated with weight gain and obesity [11].

We conclude that social stress response is sensitized in adolescents with excess weight, hindering their attentional function. The study has important strengths including the experimental design, the power-informed sample size, the detailed phenotypic characterization and the group matching of excess weight and normal weight adolescents, and the objective measurement of stress reactivity with cortisol biomarkers. However, the results should be as well appraised in light of relevant limitations. It is particularly important to stress that unlike the original TSST [45], the virtual reality TSST was not able to induce significant increases of cortisol levels in the control group. We selected this stressor because it was capable of inducing mild but sizeable stress in the laboratory at the same time that it reduced the ethical implications of

stressing “at risk” obese adolescents [18]. In agreement with this assumption, our results indicate that the stress manipulation was actually more effective in obese adolescents (57% of participants showed increased cortisol levels) than in controls (only 48% of participants showed increased cortisol levels). There are however several factors that may explain the variability in stress induction, such as degree of belief in the cover story or degree of immersion in the virtual reality environment, that were not systematically controlled in this study. Therefore, further studies are warranted to reassess the validity of this virtual reality version, and to replicate our findings using TSST versions that are able to unequivocally reproduce the original TSST stress induction. Moreover, in absence of a “no-stress” control condition, we cannot ascertain a causal link between stress and cognitive performance. However, we base our interpretation on previous evidence showing that improvement (rather than stability or decrease) in performance is typically expected in “no-stress” repeated administration designs [37]. A related limitation is the negative finding in relation to cognitive impulsivity. Since mild arousal improves inhibitory control in adolescents, it is plausible that the mild nature of the stressor fostered cognitive impulsivity increases rather than (expected) decreases after TSST. Future studies are warranted to address these limitations, to expand on the biological, psychological and socio-economic mediators of the impact of social stress on cognition, and to longitudinally assess the relevance of this experimental effect on public health indicators of the progression of obesity.

Supporting Information

S1 Fig. Power analysis calculations.

(TIF)

S2 Fig. Visual Analogue Scales for arousal—left panel- and stress—right panel- in excess and normal weight adolescents before and after the Trier Social Stress Task (TSST).

(TIF)

Author Contributions

Conceived and designed the experiments: AVG. Performed the experiments: MMP EDR JSRV. Analyzed the data: AVG. Contributed reagents/materials/analysis tools: MGR. Wrote the paper: AVG MFS FLT.

References

1. Ludwig DS. Childhood obesity—the shape of things to come. *N Engl J Med* 2007; 357(23): 2325–2327. PMID: [18057334](#)
2. Strauss RS, Pollack HA. Social marginalization of overweight children. *Arch Pediatr Adolesc Med* 2003; 157: 746–752. PMID: [12912779](#)
3. Gunnarsdottir T, Njardvik U, Olafsdottir AS, Craighead L, Bjarnason R. Childhood obesity and co-morbid problems: effects of Epstein’s family-based behavioral treatment in Icelandic sample. *J Eval Clin Pract* 2012; 18(2): 465–472. doi: [10.1111/j.1365-2753.2010.01603.x](#) PMID: [21210895](#)
4. Rouach V, Bloch M, Rosenberg N, Gilad S, Limor R, Stern N, et al. The acute ghrelin response to a psychological stress challenge does not predict the post-stress urge to eat. *Psychoneuroendocrinology* 2007; 32(6): 693–702. PMID: [17560728](#)
5. Therrien F, Drapeau V, Lalonde J, Lupien SJ, Beaulieu S, Tremblay A, et al. Awakening cortisol response in lean, obese, and reduced obese individuals: effect of gender and fat distribution. *Obesity* 2007; 15(2): 377–385. PMID: [17299111](#)
6. Liang J, Matheson BE, Kaye WH, Boutelle KN. Neurocognitive correlates of obesity and obesity-related behaviors in children and adolescents. *Int J Obes* 2014; 38(4):494–506. doi: [10.1038/ijo.2013.142](#) PMID: [23913029](#)
7. Davidson RJ, McEwen BS. Social influences on neuroplasticity: stress and interventions to promote well-being. *Nat neurosci* 2012; 15(5): 689–695. doi: [10.1038/nn.3093](#) PMID: [22534579](#)

8. Tsai SF, Huang TY, Chang CY, Hsu YC, Chen SJ, Yu L, et al. Social instability stress differentially affects amygdalar neuron adaptations and memory performance in adolescent and adult rats. *Front Behav Neurosci* 2014; 8:27. doi: [10.3389/fnbeh.2014.00027](https://doi.org/10.3389/fnbeh.2014.00027) PMID: [24550802](https://pubmed.ncbi.nlm.nih.gov/24550802/)
9. Bechara A. Decision making, impulse control and loss of willpower to resist drugs: a neurocognitive perspective. *Nat neurosci* 2005; 8(11): 1458–1463. PMID: [16251988](https://pubmed.ncbi.nlm.nih.gov/16251988/)
10. Chudasama Y, Passetti F, Rhodes SEV, Lopian D, Desai A, Robbins TW. Dissociable aspects of performance on the 5-choice serial reaction time task following lesions of the dorsal anterior cingulate, infralimbic and orbitofrontal cortex in the rat: differential effects on selectivity, impulsivity and compulsivity. *Behav Brain Res* 2003; 146(1): 105–119.
11. Seeyave DM, Coleman S, Appugliese D, Corwyn RF, Bradley RH, Davidson NS, et al. Ability to delay gratification at age 4 years and risk of overweight at age 11 years. *Arch Pediatr Adolesc Med* 2009; 163(4): 303–308. doi: [10.1001/archpediatrics.2009.12](https://doi.org/10.1001/archpediatrics.2009.12) PMID: [19349558](https://pubmed.ncbi.nlm.nih.gov/19349558/)
12. Frazier-Wood AC, Carnell S, Pena O, Hughes SO, O'Connor TM, Asherson P, et al. Reaction time but not response inhibition shares genetic influences with later BMI in 7–10 years-old. *Obesity* (in press).
13. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 2012; 7(4): 284–294. doi: [10.1111/j.2047-6310.2012.00064.x](https://doi.org/10.1111/j.2047-6310.2012.00064.x) PMID: [22715120](https://pubmed.ncbi.nlm.nih.gov/22715120/)
14. Van den Bos R, Hartevelde M, Stoop H. Stress and decision-making in humans: performance is related to cortisol reactivity, albeit differently in men and women. *Psychoneuroendocrinology* 2009; 34(10): 1449–1458. doi: [10.1016/j.psyneuen.2009.04.016](https://doi.org/10.1016/j.psyneuen.2009.04.016) PMID: [19497677](https://pubmed.ncbi.nlm.nih.gov/19497677/)
15. Preston SD, Buchanan TW, Stansfield RB, Bechara A. Effects of anticipatory stress on decision making in a gambling task. *Behav Neurosci* 2007; 121(2): 257. PMID: [17469915](https://pubmed.ncbi.nlm.nih.gov/17469915/)
16. Van Strien T., Frijters JER, Bergers GPA, Defares PB. The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *International Journal of Eating Disorders* 1986; 5(2): 295–315.
17. Garner DM. EDI 2: inventario de trastornos de la conducta alimentaria: manual. Tea ediciones: Madrid, 1994.
18. Kelly O, Matheson K, Martinez A, Merali Z, Anisman H. Psychosocial stress evoked by a virtual audience: relation to neuroendocrine activity. *CyberPsychol Behav* 2007; 10(5): 655–662. PMID: [17927533](https://pubmed.ncbi.nlm.nih.gov/17927533/)
19. Robbins TW, James M, Owen AM, Sahakian BJ, Lawrence AD, McInnes L, et al. A study of performance on tests from the CANTAB battery sensitive to frontal lobe dysfunction in a large sample of normal volunteers: implications for theories of executive functioning and cognitive aging. *Cambridge Neuropsychological Test Automated Battery. J Int Neuropsychol Soc.* 1998; 4(5): 474–90. PMID: [9745237](https://pubmed.ncbi.nlm.nih.gov/9745237/)
20. Bechara A. Iowa Gambling Task. Boca Raton, FL, USA: Psychological Assessment Resources: 2007.
21. Verdejo-Garcia A, Bechara A, Recknor EC, Perez-Garcia M. Decision-making and the Iowa gambling task: Ecological validity in individuals with substance dependence. *Psychol Belg* 2006; 46(1/2): 55–78.
22. Butler TR, Ariwodola OJ, Weiner JL. The impact of social isolation on HPA axis function, anxiety-like behaviors, and ethanol drinking. *Front Integr Neurosci* 2014; 7:102. doi: [10.3389/fnint.2013.00102](https://doi.org/10.3389/fnint.2013.00102) PMID: [24427122](https://pubmed.ncbi.nlm.nih.gov/24427122/)
23. Jahn AL, Fox AS, Abercrombie HC, Shelton SE, Oakes TR, Davidson RJ, et al. Subgenual prefrontal cortex activity predicts individual differences in hypothalamic-pituitary-adrenal activity across different contexts. *Biol Psychiatry* 2010; 67(2): 175–181. doi: [10.1016/j.biopsych.2009.07.039](https://doi.org/10.1016/j.biopsych.2009.07.039) PMID: [19846063](https://pubmed.ncbi.nlm.nih.gov/19846063/)
24. Liu J, Chaplin TM, Wang F, Sinha R, Mayes LC, Blumberg HP. Stress reactivity and corticolimbic response to emotional faces in adolescents. *J Am Acad Child Adolesc Psychiatry* 2012; 51(3): 304–312. doi: [10.1016/j.jaac.2011.12.014](https://doi.org/10.1016/j.jaac.2011.12.014) PMID: [22365466](https://pubmed.ncbi.nlm.nih.gov/22365466/)
25. Root JC, Tiescher O, Cunningham-Bussell A, Pan H, Epstein J, Altemus M, et al. Frontolimbic function and cortisol reactivity in response to emotional stimuli. *Neuroreport* 2009; 20(4): 429–434. doi: [10.1097/WNR.0b013e328326a031](https://doi.org/10.1097/WNR.0b013e328326a031) PMID: [19225430](https://pubmed.ncbi.nlm.nih.gov/19225430/)
26. Therrien F, Drapeau V, Lalonde J, Lupien SJ, Beaulieu S, Doré J, et al. Cortisol response to the Trier Social Stress Test in obese and reduced obese individuals. *Biol Psychol* 2010; 84(2): 325–329. doi: [10.1016/j.biopsycho.2010.03.013](https://doi.org/10.1016/j.biopsycho.2010.03.013) PMID: [20302906](https://pubmed.ncbi.nlm.nih.gov/20302906/)
27. Rao U, Hammen C, Ortiz LR, Chen LA, Poland RE. Effects of early and recent adverse experiences on adrenal response to psychosocial stress in depressed adolescents. *Biol Psychiatry* 2008; 64(6): 521–526. doi: [10.1016/j.biopsych.2008.05.012](https://doi.org/10.1016/j.biopsych.2008.05.012) PMID: [18597740](https://pubmed.ncbi.nlm.nih.gov/18597740/)
28. Masten CL, Eisenberger NI, Borofsky LA, McNealy K, Pfeifer JH, Dapretto M. Subgenual anterior cingulate responses to peer rejection: a marker of adolescents' risk for depression. *Dev Psychopathol* 2011; 23(1): 283–292. doi: [10.1017/S0954579410000799](https://doi.org/10.1017/S0954579410000799) PMID: [21262054](https://pubmed.ncbi.nlm.nih.gov/21262054/)

29. Masten CL, Eisenberger NI, Pfeifer JH, Dapretto M. Neural responses to witnessing peer rejection after being socially excluded: fMRI as a window into adolescents' emotional processing. *Dev Sci* 2013; 16(5): 743–759 doi: [10.1111/desc.12056](https://doi.org/10.1111/desc.12056) PMID: [24033579](https://pubmed.ncbi.nlm.nih.gov/24033579/)
30. De Santis AS, Diez Roux AV, Hajat A, Aiello AE, Golden SH, Jenny NS, et al. Associations of salivary cortisol levels with inflammatory markers: The Multi-Ethnic Study of Atherosclerosis. *Psychoneuroendocrinology* 2012; 37(7): 1009–1018. doi: [10.1016/j.psyneuen.2011.11.009](https://doi.org/10.1016/j.psyneuen.2011.11.009) PMID: [22178583](https://pubmed.ncbi.nlm.nih.gov/22178583/)
31. Slavich GM, Way BM, Eisenberger NI, Taylor SE. Neural sensitivity to social rejection is associated with inflammatory responses to social stress. *Proc Natl Acad Sci usa* 2010; 107(33): 14817–14822. doi: [10.1073/pnas.1009164107](https://doi.org/10.1073/pnas.1009164107) PMID: [20679216](https://pubmed.ncbi.nlm.nih.gov/20679216/)
32. He Q, Chen C, Dong Q, Xue G, Chen C, Lu ZL, et al. Gray and white matter structures in the midcingulate cortex region contribute to body mass index in Chinese young adults. *Brain Struct Funct* 2013; Oct 22. [Epub ahead of print]
33. Moreno-Lopez L, Soriano-Mas C, Delgado-Rico E, Rio-Valle JS, Verdejo-Garcia A. Brain structural correlates of reward sensitive and impulsivity in adolescents with normal and excess weight. *Plos One* 2012; 7(11):e49185. doi: [10.1371/journal.pone.0049185](https://doi.org/10.1371/journal.pone.0049185) PMID: [23185306](https://pubmed.ncbi.nlm.nih.gov/23185306/)
34. Lu Q, Tao F, Hou F, Zhang Z, Sun Y, Xu Y, et al. Cortisol reactivity, delay discounting and percent body fat in Chinese urban young adolescents. *Appetite* 2014; 72: 13–20. doi: [10.1016/j.appet.2013.09.019](https://doi.org/10.1016/j.appet.2013.09.019) PMID: [24080189](https://pubmed.ncbi.nlm.nih.gov/24080189/)
35. Van Strien T, Roelofs K, de Weerth C. Cortisol reactivity and distress-induced emotional eating. *Psychoneuroendocrinology* 2013; 38(5): 677–684. doi: [10.1016/j.psyneuen.2012.08.008](https://doi.org/10.1016/j.psyneuen.2012.08.008) PMID: [22999262](https://pubmed.ncbi.nlm.nih.gov/22999262/)
36. Kubzansky LD, Bordelois P, Jun HJ, Roberts AL, Cerda M, Bluestone N, et al. The weight of traumatic stress: A prospective study of posttraumatic stress disorder symptoms and weight status in women. *JAMA Psychiatry* 2014; 71(1): 44–51. doi: [10.1001/jamapsychiatry.2013.2798](https://doi.org/10.1001/jamapsychiatry.2013.2798) PMID: [24258147](https://pubmed.ncbi.nlm.nih.gov/24258147/)
37. Lowe C., Rabbitt P. (1998). Test-re-test reliability of the CANTAB and ISPOCD neuropsychological batteries: theoretical and practical issues. *Neuropsychologia*, 36(9), 915–923. PMID: [9740364](https://pubmed.ncbi.nlm.nih.gov/9740364/)
38. Baeken C, Vanderhasselt MA, De Raedt R. Baseline 'state anxiety' influences HPA-axis sensitivity to one sham-controlled HF-rTMS session applied to the right dorsolateral prefrontal cortex. *Psychoneuroendocrinology* 2011; 36(1): 60–67. doi: [10.1016/j.psyneuen.2010.06.006](https://doi.org/10.1016/j.psyneuen.2010.06.006) PMID: [20599325](https://pubmed.ncbi.nlm.nih.gov/20599325/)
39. Kiem SA, Andrade KC, Spoormaker VI, Holsboer F, Czisch M, S amann PG. Resting state functional MRI connectivity predicts hypothalamus-pituitary-axis status in healthy males. *Psychoneuroendocrinology* 2013; 38(8): 1338–1348. doi: [10.1016/j.psyneuen.2012.11.021](https://doi.org/10.1016/j.psyneuen.2012.11.021) PMID: [23279846](https://pubmed.ncbi.nlm.nih.gov/23279846/)
40. Thomason ME, Hamilton JP, Gotlib IH. Stress-induced activation of the HPA axis predicts connectivity between subgenual cingulate and salience network during rest in adolescents. *J Child Psychol Psychiatry* 2011; 52(10): 1026–1034. doi: [10.1111/j.1469-7610.2011.02422.x](https://doi.org/10.1111/j.1469-7610.2011.02422.x) PMID: [21644985](https://pubmed.ncbi.nlm.nih.gov/21644985/)
41. Hanson JL, Chung MK, Avants BB, Rudolph KD, Shirtcliff EA, Gee JC, et al. Structural variations in prefrontal cortex mediate the relationship between early childhood stress and spatial working memory. *J Neurosci* 2012; 32(23): 7917–7925. doi: [10.1523/JNEUROSCI.0307-12.2012](https://doi.org/10.1523/JNEUROSCI.0307-12.2012) PMID: [22674267](https://pubmed.ncbi.nlm.nih.gov/22674267/)
42. Gardner M, Steinberg L. Peer influence on risk taking, risk preference, and risky decision making in adolescence and adulthood: an experimental study. *Dev Psychol* 2005; 41(4): 625. PMID: [16060809](https://pubmed.ncbi.nlm.nih.gov/16060809/)
43. Chein J, Albert D, O'Brien L, Uckert K, Steinberg L. Peers increase adolescent risk taking by enhancing activity in the brain's reward circuitry. *Dev Sci* 2011; 14(2): F1–10. doi: [10.1111/j.1467-7687.2010.01035.x](https://doi.org/10.1111/j.1467-7687.2010.01035.x) PMID: [21499511](https://pubmed.ncbi.nlm.nih.gov/21499511/)
44. Verdejo-Garcia A, Verdejo-Roman J, Lagos FM, Lacomba JA, Schmidt-Rio-Valle J, Soriano-Mas C. Dysfunctional involvement of emotion and reward brain regions on social decision-making in excess weight adolescents. *Hum Brain Mapp* (in press).
45. Kirschbaum C., Pirke K. M., Hellhammer D. H. (1993). The 'Trier Social Stress Test'—a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*, 28(1–2), 76–81. PMID: [8272203](https://pubmed.ncbi.nlm.nih.gov/8272203/)