

RESEARCH ARTICLE

A Novel Pathogenesis-Related Class 10 Protein *Gly m 4l*, Increases Resistance upon *Phytophthora sojae* Infection in Soybean (*Glycine max* [L.] Merr.)

Sujie Fan¹, Liangyu Jiang¹, Junjiang Wu², Lidong Dong¹, Qun Cheng¹, Pengfei Xu^{1*}, Shuzhen Zhang^{1*}

1 Soybean Research Institute, Key Laboratory of Soybean Biology of Chinese Education Ministry, Northeast Agricultural University, Harbin, 150030, Heilongjiang, People's Republic of China, **2** Soybean Research Institute, Heilongjiang Academy of Agricultural Sciences, Collaborative Innovation Center of Grain Production Capacity Improvement in Heilongjiang Province, Harbin, 150086, Heilongjiang, People's Republic of China

* zhangshuzhen@neau.edu.cn (SZZ); xupengfei@neau.edu.cn (PFX)



OPEN ACCESS

Citation: Fan S, Jiang L, Wu J, Dong L, Cheng Q, Xu P, et al. (2015) A Novel Pathogenesis-Related Class 10 Protein *Gly m 4l*, Increases Resistance upon *Phytophthora sojae* Infection in Soybean (*Glycine max* [L.] Merr.). PLoS ONE 10(10): e0140364. doi:10.1371/journal.pone.0140364

Editor: Tianzhen Zhang, Nanjing Agricultural University, CHINA

Received: July 30, 2015

Accepted: September 24, 2015

Published: October 16, 2015

Copyright: © 2015 Fan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The research was supported through funding from the Heilongjiang Province outstanding youth fund (JC201308), NSFC Projects (31071439, 31171577, 31101167), Natural Science Foundation of Heilongjiang Province (C2015010), the Specialized Research Fund for the Doctoral Program of Higher Education (20112325120005), the Science and Technology Innovation Project in Harbin (2012RFQXN011, 2012RFXXN019), and the

Abstract

Phytophthora root and stem rot of soybean, caused by *Phytophthora sojae* (*P. sojae*), is a destructive disease in many soybean planting regions worldwide. In a previous study, an expressed sequence tag (EST) homolog of the major allergen *Pru ar 1* in apricot (*Prunus armeniaca*) was identified up-regulated in the highly resistant soybean ‘Suinong 10’ infected with *P. sojae*. Here, the full length of the EST was isolated using rapid amplification of cDNA ends (RACE). It showed the highest homology of 53.46% with *Gly m 4* after comparison with the eight soybean allergen families reported and was named *Gly m 4-like* (*Gly m 4l*, GenBank accession no. HQ913577.1). The cDNA full length of *Gly m 4l* was 707 bp containing a 474 bp open reading frame encoding a polypeptide of 157 amino acids. Sequence analysis suggests that *Gly m 4l* contains a conserved ‘P-loop’ (phosphate-binding loop) motif at residues 47–55 aa and a Bet v 1 domain at residues 87–120 aa. The transcript abundance of *Gly m 4l* was significantly induced by *P. sojae*, salicylic acid (SA), NaCl, and also responded to methyl jasmonic acid (MeJA) and ethylene (ET). The recombinant *Gly m 4l* protein showed RNase activity and displayed directly antimicrobial activity that inhibited hyphal growth and reduced zoospore release in *P. sojae*. Further analyses showed that the RNase activity of the recombinant protein to degrading tRNA was significantly affected in the presence of zeatin. Over-expression of *Gly m 4l* in susceptible ‘Dongnong 50’ soybean showed enhanced resistance to *P. sojae*. These results indicated that *Gly m 4l* protein played an important role in the defense of soybean against *P. sojae* infection.

Research Fund for Young Teachers through NEAU (2012 RCB 08).

Competing Interests: The authors have declared that no competing interests exist.

Introduction

Pathogenesis-related (PR) proteins were initially described as a class of plant-specific proteins that accumulated in response to pathogen infection; however, subsequent studies showed that these proteins are induced through a variety of biotic and abiotic stresses [1–2]. In addition, PR proteins are reported constitutively expressed in different tissues and organs during growth [3–4] and are currently classified into 17 functional families based on primary structure, serological relationships, and biological activities [1, 4]. Most PR proteins are localized in the vacuole extracellularly or intracellularly. In contrast, PR10 proteins are intracellular and cytoplasmic [5].

Most members of the PR10 protein contain an open reading frame (ORF) of 456 to 696 nucleotides, usually interrupted by an intron of 76–359 nucleotides, which encode a polypeptide of 151–231 amino acids with predicted molecular masses of 15–26 kDa [6–8]. In general, the 3D structures of PR10 proteins contain a 25-amino-acid long α -helix ($\alpha 3$) in C-terminal and two short α -helices ($\alpha 1$ and $\alpha 2$) in N-terminal. The $\alpha 3$ is surrounded by seven anti-parallel β -sheets ($\beta 1$ to $\beta 7$) and the $\alpha 1$ and $\alpha 2$ are located between $\beta 1$ and $\beta 2$ strands [9–10].

The PR10 proteins, first described in cultured parsley (*Petroselinum crispum*) cells [11], have been reported in numerous plant species when induced by various pathogens such as bacteria, fungi, oomycetes, and viruses [12–20]. Moreover, PR10 genes are also induced through various abiotic stresses, such as salt, heat, cold, PEG, dark, drought, ozone, UV irradiation, copper, H_2O_2 and wounding [6, 18, 21–29]. The expression levels of some PR10 genes are regulated following treatments with signaling molecules and plant hormones [30–32].

PR10 proteins are not only involved in plant defense, they also play a role in developmental processes [33] and exhibit enzymatic activities in secondary metabolism [34], suggesting that these proteins constitute a multifunctional protein family [35–36]. The structural similarity between PR10 proteins and ribonucleases from phosphate-starved ginseng (*Panax ginseng*) cells suggests that PR10 proteins function as ribonucleases [37], and several studies have shown that members of the PR10 protein family possess RNA degrading activities [38–39]. Notably, Park et al. [20] showed that CaPR10 from hot pepper (*Capsicum annuum*) exhibited RNase activity and also demonstrated that this protein was directly involved in antiviral and antimicrobial processes *in vitro*. Another study demonstrated that the ribonuclease activity of AhPR10 from peanut (*Arachis hypogaea* L.) was essential for the antifungal activity of this protein, as the loss of the ribonuclease in mutants resulted in a loss of antifungal activity [5]. In addition, the RNase and DNase activities of PBZ1, a PR10 family protein in rice (*Oryza sativa*), and VpPR-10.1, from Chinese wild grape (*Vitis pseudoreticulata*), have been shown to induce programmed cell death and DNA degradation [16, 40]. Most PR10 proteins have also been shown to bind to plant hormones, including cytokinins [41–42], brassinosteroids [10], and hydrophobic ligands [43], which might be important for plant defense responses to pathogens and plant growth and development [10].

In recent years, a variety of PR proteins and homologs causing allergenicity in humans have been isolated and characterized. Bet v 1, belonging to PR10 family, isolated from *Betula verrucosa*, is the main allergen present in pollen grains [44]. Subsequently, a considerable number of diverse plant allergen proteins with sequence similarity to Bet v 1 have been identified, such as Mal d 1 from *Malus domestica* [45], Pru ar 1 from *Prunus armeniaca* (AF020784), Pru av 1 from *Prunus avium* [46], Pyr c 1 from *Pyrus communis* (AF057030), Dau c 1 from *Daucus carota* [47], Pru p 1 from *Prunus persica* [48] and Api g 1 from *Apium graveolens* [49]. In soybean, a variety of allergens have been identified into eight families (<http://www.allergen.org>) till now, namely Gly m 1 [50], Gly m 2 [51], Gly m 3 [52], Gly m 4 [53–55], Gly m 5 [56–57], Gly m 6

[56–57], Gly m 7 and Gly m 8 [58]. Gly m 4, a member of the superfamily of Bet v 1 homologous proteins, are particularly similar to the PR10 proteins of yellow lupine [53–55].

In a previous study, we identified a novel up-regulated sequence tag (EST), homologous to major allergen Pru ar 1 from apricot (*Prunus armeniaca*) (GenBank Accession no. AF020784), in the highly resistant soybean ‘Suinong 10’ after infection with the oomycete *Phytophthora sojae* (*P. sojae*) [59]. In the present study, the full length of the EST was isolated using rapid amplification of cDNA ends (RACE). It showed identifies of 53.46% homology with Gly m 4 protein from *Glycine max*, representing the highest homology among the eight soybean allergen families, and was designated *Gly m 4-like* (*Gly m 4l*). The expression patterns of *Gly m 4l* induced under abiotic and biotic stresses were examined. The RNase activity of Gly m 4l protein and the influence of cytokinins on the RNase activity were established. Then, the antimicrobial activity of Gly m 4l protein against hyphal growth and zoospore release in *P. sojae* were investigated. Moreover, *in vivo*, over-expressing *Gly m 4l* gene in soybean plants under the control of 35S promoter were produced and the antimicrobial activity of transgenic plants were also investigated.

Materials and Methods

Plant Materials and Growth Conditions

‘Suinong 10’, a soybean cultivar with high resistance against the predominant race 1 of *P. sojae* in Heilongjiang, China [60], was used for gene isolation. *P. sojae* race 1, PSR01, which was isolated from infected soybean plants in Heilongjiang [60], was cultivated at 25°C for 7 days on V8 juice agar in a polystyrene dish.

‘Suinong 10’ soybean seeds were planted in pots filled with sterile vermiculite in a growth chamber with a 14 h photoperiod (at a light intensity of 350 mol.m⁻²s⁻¹) at 22°C/ 18°C day/night temperatures and relative humidity of 90 ± 10%.

The seeds of ‘Dongnong 50’ soybean, susceptible to *P. sojae* race 1, were obtained from the Key Laboratory of Soybean Biology in Chinese Ministry of Education, Harbin and used for the gene transformation experiment.

Isolation of the Full-Length *Gly m 4l* cDNA by 5'- and 3'- RACE and Sequence Analysis

In a previous study, a cDNA library of mRNAs encoding expressed sequence tags (ESTs) that showing increased expression during *P. sojae* infection was constructed using suppression subtractive hybridization (SSH) from the leaf tissues of the highly resistant soybean ‘Suinong 10’. A soybean up-regulated EST homolog of the major allergen Pru ar 1 gene in apricot (*Prunus armeniaca*) was identified using microarray analysis and real-time PCR [59]. In the present study, the full-length cDNA was isolated by way of rapid amplification of cDNA ends (RACE), performed using the SMART RACE cDNA amplification Kit (Clontech, CA, USA) according to the manufacturer’s instructions. The gene-specific primers GSP1 for 5'- RACE and GSP2 for 3'- RACE were designed to amplify the antisense and sense strands, respectively (see S1 Table for primer sequences). All RACE PCR reactions were performed at 94°C for 5 min, followed by 35 cycles at 94°C for 30 s, 57°C for 30 s, and 72°C for 2 min, with a final extension at 72°C for 10 min. The PCR products were cloned into pMDTM18-T vector (Promega, USA), then transformed into *E. coli* DH5 α cells (Shanghai Biotech Inc, Shanghai, China) and sequenced (GENEWIZ, Beijing, China). After RACE, a single full-length cDNA sequence was obtained by combining the 5'- RACE fragment, the EST fragment and the 3'-RACE fragment. Then, the full-length cDNA was gotten through the amplification of the cDNA template reversed

transcribed from total RNA using the primer pairs F and R (S1 Table). The full length genomic DNA was also amplified using the primers mentioned above. The amplification products were gel purified and cloned into pMDTM18-T vector and sequenced.

Splicing sequences were corrected with the online BLAST programs (http://www.phytozome.net/Glycine_max; <http://www.ncbi.nlm.nih.gov/BLAST>). Sequence alignments of nucleotides and amino acids sequences were performed using DNAMAN software (<http://www.lynnon.com/>). The evolutionary tree of amino acid sequence of Gly m 4l and other PR10 members was designed to search for homologous proteins using MEGA 5.1 software (<http://www.megasoftware.net>). Three dimensional (3D) structure of Gly m 4l was predicted based on the online program Phyre 2 (<http://www.sbg.bio.ic.ac.uk/phyre2>).

Application of Abiotic and Biotic Stresses

To investigate *Gly m 4l* gene expression profiling, the seedlings of 'Suinong 10' soybean at the first-node stage (V1) [61] were used for various treatments.

To examine the effect of exogenous chemicals and stresses on Gly m 4l expression, the 'Suinong 10' soybean seedlings were treated with 0.5 mM salicylic acid (SA), 100 μ M methyl jasmonic acid (MeJA), 50 mM abscisic acid (ABA), 50 mg.L⁻¹ gibberellic acid (GA₃), 100 mM NaCl, or 20% polyethylene glycol (PEG) (Sigma-Aldrich, USA). The phytohormone was first dissolved in 0.1% (v/v) ethanol (hormone solvent) and subsequently diluted to a working concentration with deionized water. The controls were treated with the same dilutions without phytohormone. For treatment with ethylene (ET), 'Suinong 10' soybean seedlings were maintained in a chamber equilibrated with 5% (v/v) gaseous ethylene. The ethylene concentration was confirmed through gas chromatography. Control experiments were performed in an identical chamber without ethylene. After induction, the soybean seedlings were placed in the appropriate growth chamber with a 14 h photoperiod (at a light intensity of 350 mol. m⁻²s⁻¹) at 22°C/18°C day/night temperatures. For cold induction, the soybean seedlings were maintained in a 4°C incubator, and the control experiments were performed in the growth chamber at 22°C/18°C day/night temperatures. The leaves of soybean seedlings were harvested for RNA isolation at 0, 1, 3, 6, 9, 12, and 24 h after treatments with SA, MeJA, ET, ABA and GA₃. The leaves of soybean seedlings were harvested for RNA isolation at 0, 1, 6, 12, 24, 36, and 72 h after treatments with NaCl, PEG, and cold.

For *P. sojae* treatment, the soybean seedlings were infected with zoospores according to the method of Ward et al. (1979) [62]. After inoculation, the 'Suinong 10' soybean seedlings were kept at 90 ± 10% humidity conditions and the leaves were harvested at 0, 6, 12, 24, 36, and 72 h after the treatment, immediately frozen in liquid nitrogen, and stored at -80°C until RNA extraction and cDNA synthesis.

qPCR Analysis

The expression profiling of *Gly m 4l* was determined through qPCR analyses, performed using a real-time RT-PCR kit according to the manufacturer's instructions (Takara, Japan) on a CFX96 TouchTM Real-Time PCR Detection System (Bio-Rad, USA). Total RNA was extracted using Trizol reagent (Invitrogen, Shanghai, China) following the manufacturer's protocol. The first-strand cDNA was obtained by approximately 1 μ g of total RNA using M-MLV reverse transcriptase kit (Takara, Dalian, China). Each amplification reaction was performed in a total 20 μ L reaction volume containing 10 μ L of 2 \times SYBR Green PCR Master Mix, 150 nmol of each specific primer *Gly m 4l*-qF or *Gly m 4l*-qR (S1 Table), and 10 ng of the synthesized cDNA. The PCR protocol included 95°C for 1 min followed 40 cycles at 95°C for 15 s, 60°C for 15 s, and 72°C for 45 s. The amplification product was confirmed through melting curve

analysis from 95°C to 60°C every one degree. The relative expression value of different tissues was calculated by the $2^{-\Delta\Delta C_t}$ method using the soybean internal control gene *EF1 β* (GenBank accession no. NM_001248778) with the primer pairs *GmEF1 β -F* and *GmEF1 β -R* (S1 Table). The relative expression value for treatments with abiotic and biotic stresses was calculated by the $2^{-\Delta\Delta C_t}$ method using the soybean internal control gene *GmActin 4* (GenBank accession no. AF049106) with the primer pairs *GmActin 4-F* and *GmActin 4-R* (S1 Table). The qPCR analyses were performed on three biological replicates using RNA samples extracted from three independent plants with their respective three technical replicates.

Subcellular Localization of *Gly m 4l* with the Fusion of Green Fluorescent Protein (GFP)

The full-length coding region of *Gly m 4l* was cloned in frame into the 5'-terminus of the GFP coding sequence in 35S:GFP vector using the primer pairs *Gly m 4l-IF* and *Gly m 4l-IR* (S1 Table), generating the fusion construct 35S: Gly m 4l-GFP. The resulting fusion 35S: Gly m 4l-GFP or control 35S: GFP construct was introduced into Arabidopsis protoplasts via the polyethylene glycol (PEG) mediated transformation as described by Yoo et al. [63]. After incubation of transfected Arabidopsis protoplasts cells for 16 h at 25°C, the GFP signal was observed using a Confocal Laser Scanning Microscopy (Leica DMI 6000, Wetzlar, Germany).

Expression and Purification of the Recombinant *Gly m 4l* Protein

The full-length coding region of *Gly m 4l* was inserted in frame into the pET-29b(+) vector (EMD Millipore, USA) to create a pET29b(+)-Gly m 4l. Subsequently, the plasmid pET-29b(+)-Gly m 4l was transformed into the *E. coli* BL21 (DE3) strain, and over-expression of the cloned genes was induced with 1 mM IPTG at 37°C for 5 h. For the recombinant protein purification, the bacterial cells were pelleted after induction, resuspended in 10 mL ice-cold 1×Binding Buffer (0.5 M NaCl, 20 mM Tris-HCl, 5 mM imidazole, pH 7.9), and sonicated on ice for 10 min (30 s pulse/min) until the sample was no longer viscous. Following centrifugation at 1200×g for 15 min at 4°C, the supernatant was harvested and loaded onto a His-bind Resin column (EMD Millipore, USA). The recombinant Gly m 4l protein in the elutes was analyzed through Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and western blotting using anti-His antibody.

RNase and DNase Activity Assays

RNA degradation assays were used to examine whether the recombinant GmPrur 1 protein displayed RNase activity according to the method of Zubini et al. 2009 [48]. The samples containing 10 µg total soybean RNA were incubated with the presence of 10 µg Gly m 4l protein at 37°C for 0, 2 and 4 h in 50 µL of 50 mM buffer at different pHs (citrate for pH 3, MES for pH 5, phosphate for pH 7, and CHES (N-cyclohexyl-2-amino-ethanesulfonic acid) for pH 9). At pH 7, RNase activity of the recombinant Gly m 4l protein was also checked in the presence of zeatin. Samples containing 10 µg total soybean RNA were incubated with 10 µg recombinant Gly m 4l protein and 10 mM zeatin at 37°C for 30 min, 1, 2, 3 and 4 h. The samples incubated with the recombinant Gly m 4l protein alone or boiled Gly m 4l protein and zeatin were used as the negative controls. The result was examined by 1% agarose gel electrophoresis (AGE) then visualized under UV light.

Meanwhile, the RNase activity of the recombinant Gly m 4l protein on yeast tRNA (Sigma, USA) was also assayed by determining the generation of acid-soluble, UV-absorbing species as described by Wang and Ng [64]. Briefly, 10 µg recombinant Gly m 4l protein was incubated with 100 µg tRNA in 75 µL of phosphate buffer (pH 7.0) at 37°C for 30 min, and the reaction

was terminated by introduction of 12 μ L of 3.4% ice-cold perchloric acid, then the samples were centrifuged at 1,500 \times g for 15 min at 4°C. The OD₂₆₀ of the supernatant was read after appropriate dilution. One unit of RNase activity is defined as the amount of enzyme that brings about an increase in OD₂₆₀ of one per minute in the acid-soluble fraction per milliliter of reaction mixture. Negative controls were performed using buffer alone or the boiled recombinant Gly m 4l protein.

The DNase activity assay was performed following the methods as described by Zubini et al. [48] and He et al. [65] with some modifications. Briefly, 10 μ g total DNA prepared from soybean leaves was incubated with 10 μ g recombinant Gly m 4l protein in 50 μ L reaction mixtures in the presence of 50 mM buffer under different pH conditions incubated for 4 h at 37°C. Negative control was performed using buffer alone for 4 h at 37°C. RNase and DNase activity assays were performed on three technical replicates.

In Vitro Antimicrobial Activity of the Recombinant *Gly m 4l* Protein

The antimicrobial activity of the recombinant Gly m 4l protein was assayed by the hyphal growth inhibition method as described by Schlumbaum et al. [66] with some modifications. The pathogen *P. sojae* had been germinated on V8 juice agar plates at 25°C for 72 h until the colonies reached a diameter of 3.5 cm, and sterile filter paper discs of 0.6 cm diameter were placed at 1 cm from the growing front, followed by the application of 15 or 25 μ g of the recombinant Gly m 4l protein onto the discs. After incubation at 25°C for 24 h, the pathogen growth zones were measured. Negative controls were performed using 25 μ g of boiled Gly m 4l protein or buffer alone.

The *P. sojae* zoospores were developed according to the procedure of Ward et al. [62] with minor modifications. Ten colonies with a diameter of 0.8 cm of *P. sojae* were removed and suspended in 10 mL sterile water to develop zoospores. After the fourth replacement of water, 100 μ g of the recombinant Gly m 4l protein was mixed with 10 mL of the supernatant and incubated at 25°C in the dark. After incubation in the dark for 16 h, the zoospore release was observed using a Leica DMI 6000 microscope (Wetzlar, Germany) and the concentration was estimated using a hemacytometer. Negative controls were performed using 100 μ g of boiled recombinant Gly m 4l protein or buffer alone. Antimicrobial activity assay of the recombinant Gly m 4l protein was performed on three technical replicates.

Vector Construction and Soybean Transformation

To over-express *Gly m 4l* gene under the cauliflower mosaic virus (CaMV) 35S promoter of pCAMBIA3301 vector (www.camia.org) containing the *bar* gene as the selective marker, the full-length coding region of *Gly m 4l* was cloned in frame into the pCAMBIA3301 vector with the primer pairs *Gly m 4l*-oF and *Gly m 4l*-oR (S1 Table). The recombinant construct 35S: *Gly m 4l* was introduced into *Agrobacterium tumefaciens* strain LBA4404 using the freeze-thaw method as described by Holsters et al. [67]. The *Agrobacterium*-mediated soybean transformation [68] was performed using cotyledonary nodes of 'Dongnong 50' soybean as explants. After darkness culture, shoot proliferation, elongation induction, root differentiation and plantlet regeneration, the regeneration plants were transferred into pots and grown in the greenhouse.

Pathogen Response Assays of Transgenic Soybean Plants

The seeds of the T₀ transgenic soybean plants were collected, dried at 25°C, and grown on soil. The T₁ transgenic soybean plants were identified using PCR amplification with the primer pairs *bar*-F and *bar*-R to amplify regions of the *bar* reporter gene (see S1 Table for primer sequences). Then, the T₁ transgenic soybean plants were further identified using Southern

hybridization with the DIG High Prime DNA Labeling and Detection Starter Kit II (Roche Cat., Germany), and the probe used for Southern hybridization was a nucleotides of 403 bp from *bar* gene.

The fully expanded leaves of T₂ transgenic soybean plants, of which the T₁ plants were tested through PCR and Southern hybridization, were further identified with the primer pairs *Gly m 4l*-qF or *Gly m 4l*-qR using qPCR (see [S1 Table](#) for primer sequences) and performed to screen on resistance to *P. sojae* according to the procedure of Kim et al. [69] with minor modifications. Briefly, the living leaves inoculated with *P. sojae* were covered with a polytene bag to maintain the relative humidity and the soybean seedlings were incubated in a mist chamber at 25°C, with 90 ± 10% relative humidity under a 16 h photoperiod at a light intensity of 350 mol m⁻² s⁻¹ for investigation. After 48 h and 96 h, the disease symptoms on each leaf sample were observed and photographed using Canon IXUS 860IS camera. To further investigate the responses of *Gly m 4l*-overexpressing plants to *P. sojae* ingress, the seedlings of T₃ transgenic soybean plants were identified with the primer pairs *Gly m 4l*-qF or *Gly m 4l*-qR using qPCR (see [S1 Table](#) for primer sequences), and the relative biomass of *P. sojae* in infected cotyledons of the selected T₃ transgenic plants at the first-node stage (V1) [61] were assessed after 48 h incubation with zoospores suspension of *P. sojae*. The *P. sojae* zoospores were developed according to the procedure of Ward et al. [62] with minor modifications and the concentration was 8 × 10⁵ mL⁻¹ estimated using a hemacytometer. The assessment of biomass of *P. sojae* was based on the transcript level of *P. sojae* *TEF1* (GenBank accession no. EU079791) in reference to soybean *EF1β* according to the method of Chacón et al. [70] (see [S1 Table](#) for *TEF1* and *EF1β* primer sequences). The pathogen response assays were performed on three biological replicates with their respective three technical replicates.

Results

Isolation and Sequence Analysis of *Gly m 4l*

The full-length cDNA sequence was obtained from 'Suinong 10' soybean using RACE. Alignment and phylogenetic tree analysis of the full-length amino acids sequence with the eight soybean allergen families reported showed identities of 53.46% homology with Gly m 4 protein, representing the highest homology ([S1 Fig](#)), and it was designated Gly m 4-like (Gly m 4l, GenBank accession no. HQ913577.1). As shown in [Fig 1](#), the 707 bp *Gly m 4l* gene contains a single open reading frame (ORF) of 474 nucleotides encoding a polypeptide of 157 amino acids, with a calculated molecular mass of 17.14 kDa and a theoretical PI of 4.98. The nucleotide sequence showed a 5' untranslated region (5' UTR) of 88 nucleotides and a 3' UTR of 145 nucleotides, along with a poly A signal (AATAAA) at 656–661 bp. The genomic *Gly m 4l* DNA sequence was also amplified, showing that *Gly m 4l* had an intron of 341 bps at 184 bp. The results of the database search revealed that Gly m 4l contained no signal peptide. The predicted structure of Gly m 4l protein included a conserved 'P-loop' (phosphate-binding loop) motif (GXGGXGXXK at residues 47–55 aa) and a Bet v 1 domain comprising 34 amino acids (at residues 87–120 aa), present in many PR 10 proteins. A neighbor-joining (NJ) phylogram was used to construct a phylogenetic tree based on the deduced sequence of Gly m 4l that contained other members of the PR10 family ([Fig 2A](#)), indicating that these proteins might share a common ancestor and display similar functions. The homology analysis of Gly m 4l with five nearby PR10 protein sequences showed that it shared 56.9% identity with PdPR10 (*Prunus domestica*) (ABW99634.1), representing the highest homology ([Fig 2B](#)). The prediction of the three dimensional (3D) structure of Gly m 4l based on the data from Phyre (<http://www.sbg.bio.jc.ac.uk/phyre/>), showed that this protein was very similar to that of the other PR10 proteins ([Fig 2C](#)). Gly m 4l contained a 24-amino-acid long α -helix (α 2) at the C-terminus and a

```

1      GGACACCCTAAGAGAGCAAAATATTTTCTTCTATCTCATCCTTTCTTTTAGTTTTATTT
61     TATTTTTTTATCTTGGGTAAAAAATCATGGGTGTTTCACTTCTGAAAGCGAACACGT
1      M G V F T S E S E H V
                                           beta 1
121    TTCCCCGTCTCTGCTGCAAAATTATACAAAGCTATTGTCCTAGATGCCAGCAATGTCTT
12     S P V S A A K L Y K A I V L D A S N V F
           L1             alpha 1             L2
181    CCCAAAAGCATTGCCAAATTTCAATTAAGAGCGTAGAAACCATTGAAGGAGATGGAGGCC
32     P K A L P N F I K S V E T I E G D G G P
                                           beta 2             L3 P-loop motif
241    AGGAACCATTAAGAAGCTTACTCTTGCTGAAGGTTTAGGTTATGTGAAGCACCACGTAGA
52     G T I K K L T L A E G L G Y V K H H V D
           beta 3             L4             beta 4
301    TACAATTGACACAGAAAATATGTGTACAATATAGTGTGATTGAAGGCAGTGCATTGTC
72     T I D T E N Y V Y N Y S V I E G S A L S
           L5             beta 5             L6
361    GGAGCCATTGGAGAAGATATGTTATGAGTACAACTGGTGGCAACCCAGATGGAGGATC
92     E P L E K I C Y E Y K L V A T P D G G S
           beta 6             Bet v 1 motif             L7
421    CATTGTGAAGTCCACAAGCAAATACTATACCAAAGGTGATGAGCAACTCGCCGAAGAATA
112    I V K S T S K Y Y T K G D E Q L A E E Y
           beta 7             L8
481    GTGAAGACTGGCAAGGAGAGATCTGCAGTTTCACCAAGGCTATTGAGGATTTCATTCA
132    V K T G K E R S A G F T K A I E D F I Q
                                           alpha 2
541    GGCTAATCCTGATTACAACTAATAAAGTCACCAATTATAATTATGTCTCAGTTTTATTAA
152    A N P D Y N *

601    TTATGGTCATTTGTGTTTTTTTTCCCTTCTTGTGCGTGTGCTTAAATTTCCAAATAA
661    AATTATGTTTGACTTTGCTTCATGAGAAAAGTTTGAGATTACT

```

Fig 1. Nucleotide and amino acid sequence of *Gly m 4l*. The P-loop motif and Bet v 1 motif are shown in shadow. The α -helices and β -sheets are marked underlined.

doi:10.1371/journal.pone.0140364.g001

short α -helix ($\alpha 1$) at the N- terminus. The $\alpha 2$ was surrounded by seven anti-parallel β -sheets ($\beta 1$ to $\beta 7$) and the $\alpha 1$ was located between the $\beta 1$ and $\beta 2$ strands. The short loop structures were named L1 to L8 located between the α -helix and β - sheet (Fig 2A). The results suggested that *Gly m 4l* belonged to PR 10 protein family.

The Transcript Abundance Patterns of *Gly m 4l*

Quantitative real-time PCR was performed to assess the tissue-specific expression of *Gly m 4l* in ‘Suinong 10’ soybean. The results showed that *Gly m 4l* was constitutively and highly expressed in the roots, followed by the leaves and stems (Fig 3A). The transcript abundance of *Gly m 4l* was investigated in the fully expanded unifoliolate leaves of ‘Suinong 10’ soybean in response to biotic and abiotic stresses. For *P. sojae* infection, a significant induction of *Gly m 4l* was detected from 6–72 h after the treatment, and the transcripts reached a maximum level at 24 h (Fig 3B). The transcript abundance of *Gly m 4l* under abiotic stresses was also investigated,




Fig 2. Characterization analysis of Gly m 4l. (A) Phylogeny analysis of Gly m 4l with 20 other PR10 proteins. The GenBank Accession numbers are as follows: PfPR10 (AAB07447.1), PsPR10 (AAA90954.1), MtPR10 (AES65085.1), MsPR10 (CAC37691.1), GmPR10 (XP_006582821.1), Gly m 4 (CAA42646.1), VvPR10.2 (CAC16165.1), VvPR10.3 (CBJ49381.1), Ypr10a (CAB94733.1), ZPR10 (AGL07712.1), PdPR10 (ABW99634.1), VpPR10 (ABC86747.1), GhPR10 (AAG18454.1), NtPR10 (AEY11296.1), CaPR10 (ABC74798.1), SIPR10 (AHC08074.1), RaPR10 (ACH63224.1), ThPR10 (ACK38253.1), SbPR10a (AAW83207.1), and OsPR10 (BAD03969.1). (B) Alignment of amino acid sequences of Gly m 4l and the nearby 5 PR10 proteins. The 'P-loop' motif and the Bet v 1 motifs are marked underlined. (C) The tertiary structure of Gly m 4l protein and the comparison with that of the other five PR10 proteins.

doi:10.1371/journal.pone.0140364.g002

such as NaCl, PEG, and cold (Fig 3C). NaCl treatment induced a significant up-regulation of *Gly m 4l* transcripts from 12–72 h. Cold induced an upregulation of *Gly m 4l* expression significantly only at 72 h after the treatment. There was little change for *Gly m 4l* transcript under the PEG stress. Meanwhile, the transcript abundance of *Gly m 4l* in response to exogenous chemicals including SA, MeJA, ET, ABA or GA₃ was then carried out (Fig 3D). Sprayed with SA, the transcript of *Gly m 4l* was induced significantly from 3–24 h after the treatment, and reached a maximum level at 6 h. The *Gly m 4l* transcripts were also induced by MeJA and ET. There was almost down-regulation of the *Gly m 4l* transcript with the treatments of ABA and GA₃.

Subcellular Localization of *Gly m 4l* Protein

To test the subcellular localization, the Gly m 4l protein was fused to the GFP coding sequences under the control of the CaMV 35S promoter (Fig 4A). Then, the newly constructed vector 35S: Gly m 4l-GFP was introduced into Arabidopsis protoplasts cells using PEG-mediated transient expression. As shown in Fig 4B, confocal microscopic observations showed that GFP was dispersed throughout the entire cells bombarded with the control plasmid 35S: GFP and




Fig 3. Expression patterns analysis of *Gly m 4l* by quantitative real-time PCR. (A) The transcript abundance of *Gly m 4l* in the root, stem and leaf. The Ct value of each sample was normalized to the Ct value of *GmEF1B*. (B) The transcript abundance of *Gly m 4l* in response to *P. sojae* infection. The Ct value of each sample was normalized to the Ct value of *GmActin4*, and the relative expression of *Gly m 4l* was compared with mock plants at the same time point. (C) and (D), The transcript abundance of *Gly m 4l* in response to various stresses. Exogenous chemicals are SA (0.5 mM), MeJA (100 μ M), ET (0.2 mM ethephon), ABA (50 mM), GA₃ (50 mg.L⁻¹), NaCl (100 mM), PEG (20%), and cold (4°C). The Ct value of each sample was normalized to the Ct value of *GmActin4* and the relative expression of *Gly m 4l* was compared with mock plants at the same time point. The experiments were performed on three biological replicates with their respective three technical replicates and statistically analysed using Student's t-test (*P<0.05, **P<0.01). Bars indicate standard error of the mean (SE).

doi:10.1371/journal.pone.0140364.g003

the fusion Gly m 4l-GFP protein was localized exclusively to cell membrane, indicating that Gly m 4l was a cell membrane-localized protein.

Purification of the Recombinant *Gly m 4l* Protein

The expression of the recombinant Gly m 4l protein was remarkably enhanced after induction with 0.5 mM IPTG at 37°C for 1–5 h, reaching the maximum expression at 5 h, and the recombinant Gly m 4l protein was not detected in the control groups (Fig 5A). The recombinant Gly m 4l protein was purified using His-Bind Kits (EMD Millipore, USA), and the molecular weight of the purified protein was about 17 kDa in SDS-PAGE, consistent with the calculated molecular mass (17.14 kDa) (Fig 5B). Meanwhile, the Gly m 4l protein was also immunoprecipitated using anti-His antibody, and a single band appeared on the X-ray film in the regular western blotting (Fig 5C).

RNase and DNase Activities of *Gly m 4l* Protein

To examine the RNase activity of the recombinant Gly m 4l protein, 10 μ g total RNA from ‘Sui-nong 10’ soybean was incubated with 10 μ g purified Gly m 4l protein at different pH conditions. As shown in Fig 6A, the RNase activities of Gly m 4l protein increased at 2 h and 4 h with the increase of pH from 3 to 7. At pH 9, almost no RNA digestion was detected at 2 h,




Fig 4. Subcellular localization analysis of *Gly m 4l*-GFP protein in *Arabidopsis* protoplasts. (A) Flow chart of construction 35S: *Gly m 4l*-GFP for subcellular localization analysis. (B) Images visualized by a Confocal Laser Scanning Microscopy. The images of bright-field (a and b), the GFP fluorescence (green) only (c and d), the chlorophyll autofluorescence (red) only (e and f) and combined ones (g and h) are shown. All scale bars indicate 10 μ m.

doi:10.1371/journal.pone.0140364.g004

while RNA was almost totally degraded at 4 h. To evaluate the effect of cytokinin on the RNase activity of the *Gly m 4l* protein, we examined the RNase activity in the presence of 10 mM zeatin at pH 7. The RNA digestion was delayed by 30 min in the presence of zeatin compared to that without zeatin (Fig 6B). Meanwhile, RNase activity of *Gly m 4l* was also checked using yeast tRNA, and the results showed that RNase activity of *Gly m 4l* towards yeast tRNA with the presence of zeatin was significantly lower than that without zeatin (Fig 6C).

Furthermore, the DNase activity of the *Gly m 4l* protein was also examined, and no hydrolysis was tested after 4 h of incubation at different pH conditions (Fig 6D).

In Vitro Antimicrobial Activity of the Recombinant *Gly m 4l* Protein

In vitro, inhibition activity of the recombinant *Gly m 4l* protein against hyphal growth of *P. sojae* was assayed, and hyphal growth was monitored from 24 to 72 h. After incubation for 72 h, 2–3 mm and 4–6 mm zones with inhibited hyphal growth were detected with 15 μ g and 25 μ g of the recombinant protein, respectively (Fig 7A). No inhibition of the growth of *P. sojae* was observed using discs containing elution buffer or boiled protein (Fig 7A). Antimicrobial activity of the recombinant *Gly m 4l* protein against *P. sojae* zoospores was also assayed. The amount of zoospores was significantly decreased with the presence of 100 μ g recombinant *Gly m 4l* protein, compared to that with the presence of 100 μ g boiled *Gly m 4l* protein or buffer alone (Fig 7B). These results suggested that the recombinant *Gly m 4l* protein possessed antibiotic activity against *P. sojae* through the inhibition of hyphal growth and zoospore release.

Over-Expression of *Gly m 4l* in Soybean Enhances Resistance to *P. sojae*

To investigate whether over-expression of *Gly m 4l* enhances resistance in transgenic plants, the T₁ transgenic soybean plants, confirmed through PCR and Southern hybridization (S2




Fig 5. Analysis of the purified recombinant *Gly m 4l* protein. (A) The recombinant *Gly m 4l* protein induced with 0.5 mM IPTG at 37°C for 1–5 h in *E.coli* BL21 (DE3). (B) SDS-PAGE analysis of the purified recombinant *Gly m 4l* protein using His-Bind Kits. (C) The purified recombinant *Gly m 4l* protein analyzed by immunoblotting using anti-His antibody.

doi:10.1371/journal.pone.0140364.g005

Fig), were developed to T₂ transgenic soybean plants and selected by qPCR (Fig 8A) to assay the pathogen response (S3 Fig). After 96 h incubation with *P. sojae*, the leaves of the non-transgenic soybean plants exhibited clear and large lesions compared to those of the transgenic plants (Fig 8B), and the lesion area of the transgenic soybean lines is significantly ($P < 0.01$) smaller than that of non-transgenic soybean plants after 96 h incubation with *P. sojae* (Fig 8C). Moreover, the T₃ transgenic soybean plants were confirmed by qPCR (Fig 8D), and the relative biomass of *P. sojae* in infected cotyledons after 48 h incubation with zoospores suspension of *P. sojae* was also analyzed. The results indicated that the biomass of *P. sojae* based on the




Fig 6. RNase and DNase activities of the recombinant *Gly m 4l* protein. (A) RNase activity of Gly m 4l protein under different pH conditions. Samples containing 10 µg total soybean RNA were incubated with the presence of 10 µg Gly m 4l protein at 37°C for 0, 2 and 4 h at pH 3, 5, 7 and 9 from left to right. (B) RNase activity of the recombinant Gly m 4l protein analyzed in the presence of zeatin using total soybean RNA at pH 7. Samples containing 10 µg total soybean RNA were incubated with the presence of 10 µg Gly m 4l protein and 10 mM zeatin at 37°C for 30 min, 1, 2, 3 and 4 h. The samples incubated with the presence of the recombinant Gly m 4l protein or boiled Gly m 4l protein and zeatin were used as negative controls. (C) RNase activity of the recombinant Gly m 4l protein analyzed in the presence of zeatin using yeast tRNA. Samples containing 100 µg yeast tRNA were incubated with the presence of 10 µg recombinant Gly m 4l protein and 10 mM zeatin at 37°C for 30 min. The samples incubated with the presence of Gly m 4l protein alone or boiled Gly m 4l protein and zeatin were used as controls. The results were calculated by the $2^{-\Delta\Delta Ct}$ method. (D) DNase activity of the recombinant Gly m 4l protein under different pH conditions. Samples containing 10 µg total soybean DNA and 10 µg Gly m 4l protein were incubated at 37°C for 0 and 4 h at pH 3, 5, 7 and 9 from left to right. The experiments were performed on three technical replicates and statistically analysed using Student's t-test (*P<0.05, **P<0.01). Bars indicate standard error of the mean (SE).

doi:10.1371/journal.pone.0140364.g006

transcript level of the *P. sojae TEF1* gene was significantly (P<0.01) lower in *Gly m 4l*-overexpressing transgenic plants than that in non-transgenic ones (Fig 8E). These results indicated that the over-expression of *Gly m 4l* in soybean plants improved resistance to *P. sojae*.




Fig 7. Antimicrobial activity of the recombinant *Gly m 4l* protein. (A) The antimicrobial activity of the recombinant *Gly m 4l* protein against the hyphal growth of *P. sojae*. 1, 25 μg boiled *Gly m 4l* protein. 2, 15 μg *Gly m 4l* protein; 3, Elution buffer. 4, 25 μg *Gly m 4l* protein. (B) The antimicrobial activity of the recombinant *Gly m 4l* protein against zoospore release of *P. sojae*. The experiments were performed on three technical replicates and statistically analysed using Student's t-test (* $P < 0.05$, ** $P < 0.01$). Bars indicate standard error of the mean (SE).

doi:10.1371/journal.pone.0140364.g007




Fig 8. Response of *Gly m 4l* transgenic soybean plants to *P. sojae*. (A) Quantitative real-time PCR of the T_2 transgenic soybean plants. (B) Disease symptoms on the leaves of the transgenic lines and non-transgenic lines treated with a *P. sojae* race 1 inoculum at 48 h and 96 h. (C) The lesion area of the transgenic lines and non-transgenic lines were detected after 96 h of incubation with *P. sojae*. (D) Quantitative real-time PCR of the T_3 transgenic soybean plants. (E) Quantitative real-time PCR analysis of *P. sojae* relative biomass based on the transcript level of the *P. sojae TEF1* gene. The experiment was performed on three biological replicates with their respective three technical replicates and statistically analysed using Student's t-test (* $P < 0.05$, ** $P < 0.01$). Bars indicate standard error of the mean (SE).

doi:10.1371/journal.pone.0140364.g008

Discussion

Until recently, a variety of allergens have been identified into eight families in soybean (<http://www.allergen.org>), namely Gly m 1, Gly m 2, Gly m 3, Gly m 4, Gly m 5, Gly m 6, Gly m 7 and Gly m 8. Gly m 4 is a member of the superfamily of PR10 proteins [53–55], but there is almost no study on the disease resistance of this protein. Although soybean PR10 genes were first identified in 1992 [71], there is limited information concerning the potential functions of these proteins *in vitro* and *in vivo*. In a previous study, we functionally characterized *GmPR10* (GenBank accession no. FJ960440) in soybean [14]. In the present study, we provide the first description of Gly m 4l in soybean (*Glycine max* [L.] Merr.), showing identities of 53.46% homology with Gly m 4 protein, a member of the PR10 family, containing a Bet v1-motif and P-loop motif. We also isolated and characterized the PR10 member Gly m 4l in response to pathogen (*P. sojae*) infection in soybean plants. The amino acid sequence, structure, subcellular location and other biological functions of PR10 proteins indicated that the major members of the PR10 family can be grouped into two different classes: intracellular pathogenesis-related proteins (IPR) with homology to ribonucleases [1] and (S)-norcochlorine synthases (NCS) [72]. The sequence analysis indicated that Gly m 4l contained no signal peptide and the subcellular localization analysis showed that Gly m 4l protein was localized at cell membrane, suggesting that Gly m 4l is an intracellular protein similar to IPRs of the PR10 family. Gly m 4l also shared several conserved features of known IPR PR10 proteins, such as a small molecular mass, acidic pI and an intron of 76–359 nucleotides [8, 73]. Most *PR10* genes are clustered in chromosomes [8, 14], and the analysis of *Gly m 4l* based on data obtained from the internet (<http://soybase.org/GlycineBlastPages/>) indicated that a total of nine genes were clustered into nine linkage groups, namely Gm 01, 07, 08, 09, 10, 14, 15, 17, and 18.

In this study, our results indicated that mRNA transcripts of *Gly m 4l* were remarkably increased by SA stress, but relatively low under MeJA and ET treatments, and almost decreased with ABA and GA₃ treatments (Fig 3). Therefore, we speculated that *Gly m 4l* might play a key role in soybean plants resistance to *P. sojae* mainly depending on SA signaling, which is one of the important component of signal transduction and could activate plant defense responses against pathogen attack [74]. In addition, the expression of *Gly m 4l* was also induced by NaCl and cold. It is likely that high-salinity and low-temperature resulted in increased cytosolic Ca²⁺ [75–76], and Ca²⁺ was the second messenger for hypersensitive response induction or defense gene expression and leads to up-regulation of PR proteins [77].

Some of PR10 proteins have previously been shown to possess RNase or DNase activity and performed a link to plant defense strategy repertoire, such as JIOsPR10 from *Oryza sativa* [39], VpPR10.2 from *Vitis pseudoreticulata* [65], JcPR10a from *Jatropha curcas* [78], VpPR10.1 from *Vitis pseudoreticulata* [16], VpPR10.4 and VpPR10.7 from *Vitis pseudoreticulata* [79]. On the basis of these findings, it supports the hypothesis that RNase activity was potentially crucial for plant defense, such as activating the death of infected cells to limit pathogen invasion, regulating and controlling mRNA transcription upon pathogen infection or other stresses, and directly degrading pathogenic RNA [40, 48]. Previous reports showed that a limited number of proteins have been targeted to non-classical secretory pathway without an N-terminal signal peptide [80]. Although Gly m 4l contained no signal peptide (Fig 1) and the subcellular localization analysis showed that Gly m 4l protein was localized at cell membrane (Fig 4), whether or not Gly m 4l is a secretory protein is unclear yet. Under *P. sojae* stress, one possibility is that Gly m 4l protein in soybean could be targeted to non-classical secretory pathway and secreted into the intercellular space, then it performed antimicrobial function by directly degrading pathogenic RNA to inhibit hyphal growth or zoospore release. Another is that Gly m 4l protein could not be targeted to non-classical secretory pathway, then it

performed antimicrobial functional role through degrading RNA of haustorium, which is formed and stick into cytoplasm of susceptible soybean with *P. sojae* infection [81], to limit pathogen invasion.

Members of PR10 family have been reported that they have antimicrobial activities in transgenic plants, as examples, *VpPR10.2* gene from *Vitis pseudoreticulata* responded to *Plasmopara viticola* infection and over-expression of *VpPR10.2* in the host plant enhanced resistance to *Plasmopara viticola* [65], the over-expression of *PgPR10-2* and *PgPR10-4* conferred a tolerance against fungal infection [69, 82], and over-expression of *GmPR10* in soybean enhanced resistance to *P. sojae* [14]. In the present study, *Gly m 4l* over-expression transgenic soybean plants enhanced resistance to *P. sojae in vivo* and showed a lower biomass of *P. sojae* than that in non-transgenic soybean plants, suggesting that *Gly m 4l* could inhibit *P. sojae* infection and might be a useful tool for managing Phytophthora root and stem rot in soybean plants.

Conclusions

In conclusion, a new member of the PR10 protein family, *Gly m 4l*, was isolated from resistant soybean 'Suinong 10', showed significant increased transcript abundance with *P. sojae* inoculum, and also induced by SA, NaCl, MeJA and ET. The recombinant *Gly m 4l* protein showed RNase activity and displayed directly antimicrobial activity that inhibited hyphal growth and reduction zoospore release in *P. sojae*. Further analyses showed that the RNase activity of the recombinant *Gly m 4l* protein on tRNA was significantly inhibited in the presence of zeatin. Over-expression of *Gly m 4l* in susceptible 'Dongnong 50' soybean showed enhanced resistance to *P. sojae*. These results indicated that the *Gly m 4l* protein played an important role in the defense of soybean against *P. sojae* infection.

Supporting Information

S1 Fig. Alignment and phylogenetic tree analysis of the full-length amino acids sequence with other soybean allergens.

(TIF)

S2 Fig. Southern blot analysis of transgenic soybean plants. control, non-transgenic soybean plant; ox2 and ox7, T₁ independently transgenic soybean plants; 3301, pCAMBIA3301 vector.

(TIF)

S3 Fig. Disease symptoms on the leaves of the transgenic lines and non-transgenic lines treated with a *P. sojae* race 1 inoculum.

(TIF)

S1 Table. Oligonucleotide primers used in this study.

(DOC)

Acknowledgments

The research was supported through funding from the Heilongjiang Province outstanding youth fund (JC201308), NSFC Projects (31071439, 31171577, 31101167), Natural Science Foundation of Heilongjiang Province (C2015010), the Specialized Research Fund for the Doctoral Program of Higher Education (20112325120005), the Science and Technology Innovation Project in Harbin (2012RFQXN011, 2012RFXXN019), and the Research Fund for Young Teachers through NEAU (2012 RCB 08).

Author Contributions

Conceived and designed the experiments: PFX SZZ. Performed the experiments: SJF LYJ LDD QC. Analyzed the data: JJW. Contributed reagents/materials/analysis tools: SZZ PFX. Wrote the paper: SJF LYJ LDD QC.

References

1. van Loon LC, Pierpoint WS, Boller T, Conejero V. Recommendations for naming plant pathogenesis-related proteins. *Plant Mol Biol Rep*. 1994; 12(3): 245–264. doi: [10.1007/BF02668748](https://doi.org/10.1007/BF02668748)
2. van Loon LC, Rep M, Pieterse CM. Significance of inducible defense-related proteins in infected plants. *Annu Rev Phytopathol*. 2006; 44: 135–162. doi: [10.1146/annurev.phyto.44.070505.143425](https://doi.org/10.1146/annurev.phyto.44.070505.143425) PMID: [16602946](https://pubmed.ncbi.nlm.nih.gov/16602946/)
3. Agarwal P, Agarwal P. Pathogenesis related-10 proteins are small, structurally similar but with diverse role in stress signaling. *Mol Biol Rep*. 2014; 41: 599–611. doi: [10.1007/s11033-013-2897-4](https://doi.org/10.1007/s11033-013-2897-4) PMID: [24343423](https://pubmed.ncbi.nlm.nih.gov/24343423/).
4. Christensen AB, Cho BH, Naesby M, Gregersen PL, Brandt J, Madriz-Ordeñana K et al. The molecular characterization of two barley proteins establishes the novel PR-17 family of pathogenesis-related proteins. *Mol Plant Pathol*. 2002; 3(3): 135–144. doi: [10.1046/j.1364-3703.2002.00105.x](https://doi.org/10.1046/j.1364-3703.2002.00105.x) PMID: [20569319](https://pubmed.ncbi.nlm.nih.gov/20569319/).
5. Chadha P, Das RH. A pathogenesis related protein, AhPR10 from peanut: an insight of its mode of anti-fungal activity. *Planta*. 2006; 225(1): 213–222. doi: [10.1007/s00425-006-0344-7](https://doi.org/10.1007/s00425-006-0344-7) PMID: [16832688](https://pubmed.ncbi.nlm.nih.gov/16832688/).
6. Liu XJ, Huang BB, Lin J, Fei J, Chen ZH, Pang YZ et al. A novel pathogenesis-related protein (SsPR10) from *Solanum surattense* with ribonucleolytic and antimicrobial activity is stress- and pathogen-inducible. *J Plant Physiol*. 2006; 163: 546–556. doi: [10.1016/j.jplph.2005.04.031](https://doi.org/10.1016/j.jplph.2005.04.031) PMID: [16473659](https://pubmed.ncbi.nlm.nih.gov/16473659/).
7. Hoffmann-Sommergruber K, Vanek-Krebitz M, Radauer C, Wen J, Ferreira F, Scheiner O et al. Genomic characterization of members of the Bet v 1 family: genes coding for allergens and pathogenesis-related proteins share intron positions. *Gene*. 1997; 197: 91–100. doi: [10.1016/S0378-1119\(97\)00246-1](https://doi.org/10.1016/S0378-1119(97)00246-1) PMID: [9332353](https://pubmed.ncbi.nlm.nih.gov/9332353/).
8. Gao ZS, van de Weg WE, Schaart JG, Schouten HJ, Tran DH, Kodde LP. Genomic cloning and linkage mapping of the Mal d1 (PR-10) gene family in apple (*Malus domestica*). *Theor Appl Genet*. 2005; 111: 171–183. doi: [10.1007/s00122-005-2018-4](https://doi.org/10.1007/s00122-005-2018-4) PMID: [15883791](https://pubmed.ncbi.nlm.nih.gov/15883791/).
9. Biesiadka J, Bujacz G, Sikorski MM, Jaskolski M. Crystal structures of two homologous pathogenesis-related proteins from yellow lupine. *J Mol Biol*. 2002; 319(5): 1223–1234. doi: [10.1016/S0022-2836\(02\)00385-6](https://doi.org/10.1016/S0022-2836(02)00385-6) PMID: [12079359](https://pubmed.ncbi.nlm.nih.gov/12079359/).
10. Markovic-Housley Z, Degano M, Lamba D, von Roepenack-Lahaye E, Clemens S, Susani M et al. Crystal structure of a hypoallergenic isoform of the major birch pollen allergen Bet v 1 and its likely biological function as a plant steroid carrier. *J Mol Biol*. 2003; 325: 123–133. PMID: [12473456](https://pubmed.ncbi.nlm.nih.gov/12473456/).
11. Somssich IE, Schmelzer E, Bollmann J, Hahlbrock K. Rapid activation by fungal elicitor of genes encoding “pathogenesis-related proteins” in cultured parsley cells. *Proc Natl Acad Sci USA*. 1986; 83(8): 2427–2430. PMID: [16593684](https://pubmed.ncbi.nlm.nih.gov/16593684/).
12. Kim ST, Kim SG, Hwang DH, Kang SY, Kim HJ, Lee BH et al. Proteomic analysis of pathogen-responsive proteins from rice leaves induced by rice blast fungus, *Magnaporthe grisea*. *Proteomics*. 2004; 4: 3569–3578. doi: [10.1002/pmic.200400999](https://doi.org/10.1002/pmic.200400999) PMID: [15478215](https://pubmed.ncbi.nlm.nih.gov/15478215/).
13. Walter MH, Liu JW, Grand C, Lamb CJ, Hess D. Bean pathogenesis-related (PR) proteins deduced from elicitor-induced transcripts are members of a ubiquitous new class of conserved PR proteins including pollen allergens. *Mol Gen Genet*. 1990; 222(2–3): 353–360. PMID: [2274036](https://pubmed.ncbi.nlm.nih.gov/2274036/).
14. Xu PF, Jiang LY, Wu JJ, Li W, Fan S, Zhang S. Isolation and characterization of a pathogenesis-related protein 10 gene (*GmPR10*) with induced expression in soybean (*Glycine max*) during infection with *Phytophthora sojae*. *Mol Biol Rep*. 2014; 41(8): 4899–4909. doi: [10.1007/s11033-014-3356-6](https://doi.org/10.1007/s11033-014-3356-6) PMID: [24737571](https://pubmed.ncbi.nlm.nih.gov/24737571/).
15. Colditz F, Niehaus K, Kajinski F. Silencing of PR-10-like proteins in *Medicago truncatula* results in an antagonistic induction of other PR proteins and an increased tolerance upon infection with the oomycete *Aphanomyces euteiches*. *Planta*. 2007; 226(1): 57–71. doi: [10.1007/s00425-006-0466-y](https://doi.org/10.1007/s00425-006-0466-y) PMID: [17237953](https://pubmed.ncbi.nlm.nih.gov/17237953/).
16. Xu TF, Zhao XC, Jiao YT, Wei JY, Wang L, Xu Y. A pathogenesis related protein, VpPR-10.1, from *Vitis pseudoreticulata*: an insight of its mode of antifungal activity. *PLoS One*. 2014; 9(4): e95102 doi: [10.1371/journal.pone.0095102](https://doi.org/10.1371/journal.pone.0095102) PMID: [24759805](https://pubmed.ncbi.nlm.nih.gov/24759805/)

17. Maillot P, Lebel S, Schellenbaum P, Jacques A, Walter B. Differential regulation of SERK, LEC1–Like and pathogenesis–related genes during indirect secondary somatic embryogenesis in grapevine. *Plant Physiol Bioch.* 2009; 47(8): 743–752. doi: [10.1016/j.plaphy.2009.03.016](https://doi.org/10.1016/j.plaphy.2009.03.016) PMID: [19406655](https://pubmed.ncbi.nlm.nih.gov/19406655/).
18. Bahramnejad B, Goodwin PH, Zhang J, Atnaseo C, Erickson LR. A comparison of two class 10 pathogenesis–related genes from alfalfa and their activation by multiple stresses and stress–related signaling molecules. *Plant Cell Rep.* 2010; 29(11): 1235–1250. doi: [10.1007/s00299-010-0909-6](https://doi.org/10.1007/s00299-010-0909-6) PMID: [20703880](https://pubmed.ncbi.nlm.nih.gov/20703880/).
19. Polesani M, Desario F, Ferrarini A, Zamboni A, Pezzotti M, Kortekamp A et al. cDNA–AFLP analysis of plant and pathogen genes expressed in grapevine infected with *Plasmopara viticola*. *BMC Genomics.* 2008; 9: 142–155. doi: [10.1186/1471-2164-9-142](https://doi.org/10.1186/1471-2164-9-142) PMID: [18366764](https://pubmed.ncbi.nlm.nih.gov/18366764/).
20. Park CJ, Kim KJ, Shin R, Park JM, Shin YC, Paek KH. Pathogenesis–related protein 10 isolated from hot pepper functions as a ribonuclease in an antiviral pathway. *Plant J.* 2004; 37(2): 186–198. doi: [10.1046/j.1365-3113X.2003.01951.x](https://doi.org/10.1046/j.1365-3113X.2003.01951.x) PMID: [14690503](https://pubmed.ncbi.nlm.nih.gov/14690503/).
21. Jellouli N, Ben Jouira H, Skouri H, Ghorbel A, Gourgouri A, Mliki A. Proteomic analysis of Tunisian grapevine cultivar Razegui under salt stress. *J Plant Physiol.* 2008; 165(5): 471–481. doi: [10.1016/j.jplph.2007.02.009](https://doi.org/10.1016/j.jplph.2007.02.009) PMID: [17942184](https://pubmed.ncbi.nlm.nih.gov/17942184/).
22. Wisniewski M, Bassett C, Arora R. Distribution and partial characterization of seasonally expressed proteins in different aged shoots and roots of 'Loring' peach (*Prunus persica*). *Tree Physiol.* 2004; 24(3): 339–345. PMID: [14704143](https://pubmed.ncbi.nlm.nih.gov/14704143/).
23. Borsics T, Lados M. Dodder infection induces the expression of a pathogenesis–related gene of the family PR–10 in alfalfa. *J Exp Bot.* 2002; 53(375): 1831–1832. doi: [10.1093/jxb/erf039](https://doi.org/10.1093/jxb/erf039) PMID: [12147735](https://pubmed.ncbi.nlm.nih.gov/12147735/).
24. Walter MH, Liu JW, Wunn J, Hess D. Bean ribonuclease–like pathogenesis–related protein genes (Ypr10) display complex patterns of developmental, dark–induced and exogenous–stimulus–dependent expression. *Eur J Biochem.* 1996; 239: 281–293. doi: [10.1111/j.1432-1033.1996.0281u.x](https://doi.org/10.1111/j.1432-1033.1996.0281u.x) PMID: [8706731](https://pubmed.ncbi.nlm.nih.gov/8706731/).
25. Hashimoto M, Kisseleva L, Sawa S, Furukawa T, Komatsu S, Koshiba T. A novel rice PR10 protein, RSOsPR10, specifically induced in roots by biotic and abiotic stresses, possibly via the jasmonic acid signaling pathway. *Plant Cell Physiol.* 2004; 45(4): 550–559. doi: [10.1093/pcp/pch063](https://doi.org/10.1093/pcp/pch063) PMID: [15169937](https://pubmed.ncbi.nlm.nih.gov/15169937/).
26. Agrawal GK, Rakwal R, Yonekura M, Kubo A, Saji H. Proteome analysis of differentially displayed proteins as a tool for investigating ozone stress in rice (*Oryza sativa* L.) seedlings. *Proteomics.* 2002; 2(8): 947–959. doi: [10.1002/1615-9861\(200208\)2:8<947::AID-PROT947>3.0.CO;2-J](https://doi.org/10.1002/1615-9861(200208)2:8<947::AID-PROT947>3.0.CO;2-J) PMID: [12203890](https://pubmed.ncbi.nlm.nih.gov/12203890/).
27. Rakwal R, Agrawal GK, Yonekura M. Separation of proteins from stressed rice (*Oryza sativa* L.) leaf tissues by two–dimensional polyacrylamide gel electrophoresis: induction of pathogenesis–related and cellular protect proteins by jasmonic acid, UV irradiation and copper chloride. *Electrophoresis.* 1999; 20(17): 3472–3478. doi: [10.1002/\(SICI\)1522-2683\(19991101\)20:17<3472::AID-ELPS3472>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1522-2683(19991101)20:17<3472::AID-ELPS3472>3.0.CO;2-0) PMID: [10608717](https://pubmed.ncbi.nlm.nih.gov/10608717/).
28. Utriainen M, Kokko H, Auriola S, Sarrazin O, Kärenlampi S. PR–10 protein is induced by copper stress in roots and leaves of a Cu/Zn tolerant clone of birch, *Betula pendula*. *Plant Cell Environ.* 1998; 21(8): 821–828. doi: [10.1046/j.1365-3040.1998.00326.x](https://doi.org/10.1046/j.1365-3040.1998.00326.x)
29. Liu JJ, Ekramoddoullah AKM, Yu X. Differential expression of multiple PR10 proteins in western white pine following wounding, fungal infection and cold–hardening. *Physiol Plant.* 2003; 119: 544–553. doi: [10.1046/j.1399-3054.2003.00200.x](https://doi.org/10.1046/j.1399-3054.2003.00200.x)
30. Rakwal R, Agrawal GK, Yonekura M. Light–dependent induction of *OsPR10* in rice (*Oryza sativa* L.) seedlings by the global stress signaling molecule jasmonic acid and protein phosphatase 2A inhibitors. *Plant Sci.* 2001; 161(3): 469–479. doi: [10.1016/S0168-9452\(01\)00433-2](https://doi.org/10.1016/S0168-9452(01)00433-2)
31. Poupard P, Brunel N, Leduc N, Viemont J, Strullu D, Simoneau P. Expression of a Bet v 1 homologue gene encoding a PR 10 protein in birch roots: induction by auxin and localization of the transcripts by *in situ* hybridization. *Aust J Plant Physiol.* 2001; 28(1): 57–63. doi: [10.1071/PP00040](https://doi.org/10.1071/PP00040)
32. Jwa NS, Kumar Agrawal G, Rakwal R, Park CH, Prasad Agrawal V. Molecular cloning and characterization of a novel Jasmonate inducible pathogenesis–related class 10 protein gene, *JIOsPR10*, from rice (*Oryza sativa* L.) seedling leaves. *Biochem Bioph Res Co.* 2001; 286(5): 973–983. doi: [10.1006/bbrc.2001.5507](https://doi.org/10.1006/bbrc.2001.5507) PMID: [11527396](https://pubmed.ncbi.nlm.nih.gov/11527396/).
33. Sikorski MM, Biesiadka J, Kasperska AE, Kopcińska J, Lotocka B, Golinowski W. Expression of genes encoding PR10 class pathogenesis–related proteins is inhibited in yellow lupine root nodules. *Plant Sci.* 1999; 149: 125–137. doi: [10.1016/S0168-9452\(99\)00148-X](https://doi.org/10.1016/S0168-9452(99)00148-X)
34. Koistinen KM, Soininen P, Venalainen TA, Hayrinen J, Laatikainen R, Perakyla M, et al. Birch PR–10c interacts with several biologically important ligands. *Phytochemistry.* 2005; 66(21): 2524–2533. PMID: [16246382](https://pubmed.ncbi.nlm.nih.gov/16246382/).

35. Wen J, Vanek-Krebitz M, Hoffmann-Sommergruber K, Scheiner O, Breiteneder H. The potential of Betv1 homologues, a nuclear multigene family, as phylogenetic markers in flowering plants. *Mol Phylogenet Evol.* 1997; 8(3): 317–333. doi: [10.1006/mpev.1997.0447](https://doi.org/10.1006/mpev.1997.0447) PMID: [9417891](https://pubmed.ncbi.nlm.nih.gov/9417891/).
36. Radauer C, Lackner P, Breiteneder H. The Bet v 1 fold: an ancient, versatile scaffold for binding of large, hydrophobic ligands. *BMC Evol Biol.* 2008; 8: 286. doi: [10.1186/1471-2148-8-286](https://doi.org/10.1186/1471-2148-8-286) PMID: [18922149](https://pubmed.ncbi.nlm.nih.gov/18922149/).
37. Fernandes H, Michalska K, Sikorski M, Jaskolski M. Structural and functional aspects of PR-10 proteins. 2013; 280: 1169–1199. doi: [10.1111/febs.12114](https://doi.org/10.1111/febs.12114) PMID: [23289796](https://pubmed.ncbi.nlm.nih.gov/23289796/).
38. Bantignies B, Seguin J, Muzac I, Dédaldéchamp F, Gulick P, Ibrahim R. Direct evidence for ribonucleolytic activity of a PR-10-like protein from white lupin roots. *Plant Mol Biol.* 2000; 42(6): 871–881. PMID: [10890534](https://pubmed.ncbi.nlm.nih.gov/10890534/).
39. Kim ST, Yu S, Kang YH, Kim SG, Kim JY, Kim SH et al. The rice pathogen-related protein 10 (JIOsPR10) is induced by abiotic and biotic stresses and exhibits ribonuclease activity. *Plant Cell Rep.* 2008; 27(3): 593–603. doi: [10.1007/s00299-007-0485-6](https://doi.org/10.1007/s00299-007-0485-6) PMID: [18074138](https://pubmed.ncbi.nlm.nih.gov/18074138/).
40. Kim SG, Kim ST, Wang Y, Yu S, Choi IS, Kim YC et al. The RNase activity of rice probenazole-induced protein1 (PBZ1) plays a key role in cell death in plants. *Mol Cells.* 2011; 31(1): 25–31. doi: [10.1007/s10059-011-0004-z](https://doi.org/10.1007/s10059-011-0004-z) PMID: [21110127](https://pubmed.ncbi.nlm.nih.gov/21110127/).
41. Fujimoto Y, Nagata R, Fukasawa H, Yano K, Azuma M, Iida A et al. Purification and cDNA cloning of cytokinin-specific binding protein from mung bean (*Vigna radiata*). *Eur J Biochem.* 1998; 258(2): 794–802. doi: [10.1046/j.1432-1327.1998.2580794.x](https://doi.org/10.1046/j.1432-1327.1998.2580794.x) PMID: [9874249](https://pubmed.ncbi.nlm.nih.gov/9874249/).
42. Pasternak O, Bujacz GD, Fujimoto Y, Hashimoto Y, Jelen F, Otlewski J et al. Crystal structure of *Vigna radiata* cytokinin-specific binding protein in complex with zeatin. *Plant Cell.* 2006; 18(10): 2622–2634. doi: [10.1105/tpc.105.037119](https://doi.org/10.1105/tpc.105.037119) PMID: [16998071](https://pubmed.ncbi.nlm.nih.gov/16998071/).
43. Mogensen JE, Wimmer R, Larsen JN Spangfort MD, Otzen DE. The major birch allergen, Bet v 1, shows affinity for a broad spectrum of physiological ligands. *J Biol Chem.* 2002; 277(26): 23684–23692. PMID: [11953433](https://pubmed.ncbi.nlm.nih.gov/11953433/).
44. Gajhede M, Osmark P, Poulsen FM, Ipsen H, Larsen JN, Joost van Neerven RJ et al. X-ray and NMR structure of Bet v 1, the origin of birch pollen allergy. *Nat Struct Biol.* 1996; 3(12): 1040–1045. PMID: [8946858](https://pubmed.ncbi.nlm.nih.gov/8946858/).
45. Vanek-Krebitz M, Hoffmann-Sommergruber K, Laimer da Camara Machado M, Susani M, Ebner C, Kraft D et al. Cloning and sequencing of Mal d 1, the major allergen from apple (*Malus domestica*), and its immunological relationship to Bet v 1, the major birch pollen allergen. *Biochem Biophys Res Commun.* 1995; 214(2): 538–551. doi: [10.1006/bbrc.1995.2320](https://doi.org/10.1006/bbrc.1995.2320) PMID: [7677763](https://pubmed.ncbi.nlm.nih.gov/7677763/).
46. Scheurer S, Metzner K, Haustein D, Vieths S. Molecular cloning, expression and characterization of Pru a 1, the major cherry allergen. *Mol Immunol.* 1997; 34(8–9): 619–629. PMID: [9393965](https://pubmed.ncbi.nlm.nih.gov/9393965/).
47. Peters S, Imani J, Mahler V, Foetisch K, Kaul S, Paulus KE et al. Dau c 1.01 and Dau c 1.02-silenced transgenic carrot plants show reduced allergenicity to patients with carrot allergy. *Transgenic Res.* 2011; 20(3): 547–556. doi: [10.1007/s11248-010-9435-0](https://doi.org/10.1007/s11248-010-9435-0) PMID: [20798987](https://pubmed.ncbi.nlm.nih.gov/20798987/).
48. Zubini P, Zambelli B, Musiani F, Ciurli S, Bertolini P, Baraldi E. The RNA hydrolysis and the cytokinin binding activities of PR-10 proteins are differently performed by two isoforms of the Pru p 1 peach major allergen and are possibly functionally related. *Plant Physiol.* 2009; 150(3): 1235–1247. doi: [10.1104/pp.109.139543](https://doi.org/10.1104/pp.109.139543) PMID: [19474212](https://pubmed.ncbi.nlm.nih.gov/19474212/).
49. Breiteneder H, Ebner C. Molecular and biochemical classification of plant-derived food allergens. *J Allergy Clin Immunol.* 2000; 106: 27–36. doi: [10.1067/mai.2000.106929](https://doi.org/10.1067/mai.2000.106929) PMID: [10887301](https://pubmed.ncbi.nlm.nih.gov/10887301/).
50. Codina R, Lockey RF, Fernández-Caldas E, Rama R. Identification of the soybean hull allergens responsible for the Barcelona asthma outbreaks. *Int Arch Allergy Immunol.* 1999; 119(1): 69–71. doi: [10.1159/000024178](https://doi.org/10.1159/000024178) PMID: [10341324](https://pubmed.ncbi.nlm.nih.gov/10341324/).
51. Codina R, Lockey RF, Fernández-Caldas E, Rama R. Purification and characterization of a soybean hull allergen responsible for the Barcelona asthma outbreaks. II. Purification and sequencing of the Gly m 2 allergen. *Clin Exp Allergy.* 1997; 27(4): 424–430. PMID: [9146936](https://pubmed.ncbi.nlm.nih.gov/9146936/).
52. Rihs HP, Chen Z, Ruëff F, Petersen A, Rozynek P, Heimann H et al. IgE binding of the recombinant allergen soybean profilin (rGly m 3) is mediated by conformational epitopes. *J Allergy Clin Immunol.* 1999; 104(6): 1293–1301. PMID: [10589015](https://pubmed.ncbi.nlm.nih.gov/10589015/).
53. Geng T, Liu K, Frazier R, Shi L, Bell E, Glenn K et al. Development of a Sandwich ELISA for Quantification of Gly m 4, a Soybean Allergen. *J Agric Food Chem.* 2015; 63(20): 4947–53. doi: [10.1021/acs.jafc.5b00792](https://doi.org/10.1021/acs.jafc.5b00792) PMID: [25946567](https://pubmed.ncbi.nlm.nih.gov/25946567/).
54. Julka S, Kuppanan K, Karnoup A, Dielman D, Schafer B, Young SA. Quantification of Gly m 4 protein, a major soybean allergen, by two-dimensional liquid chromatography with ultraviolet and mass spectrometry detection. *Anal Chem.* 2012; 84(22): 10019–30. doi: [10.1021/ac3024685](https://doi.org/10.1021/ac3024685) PMID: [23106440](https://pubmed.ncbi.nlm.nih.gov/23106440/).

55. Berkner H, Neudecker P, Mittag D, Ballmer-Weber BK, Schweimer K, Vieths S. Cross-reactivity of pollen and food allergens: soybean Gly m 4 is a member of the Bet v 1 superfamily and closely resembles yellow lupine proteins. *Biosci Rep*. 2009; 29: 183–192. doi: [10.1042/BSR20080117](https://doi.org/10.1042/BSR20080117) PMID: [18834331](https://pubmed.ncbi.nlm.nih.gov/18834331/)
56. Holzhauser T, Wackermann O, Ballmer-Weber BK, Bindslev-Jensen C, Scibilia J, Perono-Garoffo L et al. Soybean (*Glycine max*) allergy in Europe: Gly m 5 (b-conglycinin) and Gly m 6 (glycinin) are potential diagnostic markers for severe allergic reactions to soy. *J Allergy Clin Immunol*. 2009; 123: 452–458. doi: [10.1016/j.jaci.2008.09.034](https://doi.org/10.1016/j.jaci.2008.09.034) PMID: [18996574](https://pubmed.ncbi.nlm.nih.gov/18996574/).
57. Ito K. Grain and legume allergy. *Chem Immunol Allergy*. 2015; 101: 145–51. doi: [10.1159/000375468](https://doi.org/10.1159/000375468) PMID: [26022874](https://pubmed.ncbi.nlm.nih.gov/26022874/).
58. Kattan JD, Sampson HA. Soy reactivity may be better identified by component testing with Gly m 8 than traditional testing methods. 2015; Feb 21. Available: <https://aaaai.confex.com/aaaai/2015/webprogram/Paper19970.html>.
59. Xu PF, Wu JJ, Allen X, Li WB, Chen WY, Lv HY et al. Differentially expressed genes of soybean during infection by *Phytophthora sojae*. *J Integr Agr*. 2012; 11(3): 368–377.
60. Zhang SZ, Xu PF, Wu JJ, Allen X, Zhang JX, Li WB et al. Races of *Phytophthora sojae* and their virulences on commonly grown soybean varieties in Heilongjiang, China. *Plant Dis*. 2010; 94: 87–91. doi: [10.1094/PDIS-94-1-0087](https://doi.org/10.1094/PDIS-94-1-0087)
61. Fehr WR, Caviness CE, Burmood DT, Pennington JS. Stage of development descriptions for soybeans, *Glycine max* (L.) Merrill. *Crop Sci*. 1971; 11(6): 929–931. doi: [10.2135/cropsci1971.0011183X001100060051x](https://doi.org/10.2135/cropsci1971.0011183X001100060051x)
62. Ward EWB, Lazarovits G, Unwin CH, Buzzell RI. Hypocotyl reactions and glyceollin in soybeans inoculated with zoospores of *Phytophthora megaspoma* var. *sojae*. *Phytopathology*. 1979; 69: 951–955. doi: [10.1094/Phyto-69-951](https://doi.org/10.1094/Phyto-69-951)
63. Yoo SD, Cho YH, Sheen J. Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat Protoc*. 2007; 2: 1565–1572. doi: [10.1038/nprot.2007.199](https://doi.org/10.1038/nprot.2007.199) PMID: [17585298](https://pubmed.ncbi.nlm.nih.gov/17585298/).
64. Wang HX, Ng, TB. Purification and characterization of a potent homodimeric guanine-specific ribonuclease from fresh mushroom (*Pleurotus tuberregium*) *sclerotia*. *Int J Biochem Cell B*. 2001; 33: 483–490. doi: [10.1016/S1357-2725\(01\)00038-3](https://doi.org/10.1016/S1357-2725(01)00038-3) PMID: [11331203](https://pubmed.ncbi.nlm.nih.gov/11331203/).
65. He MY, Xu Y, Cao JJ, Zhu Z, Jiao Y, Wang Y et al. Subcellular localization and functional analyses of a PR10 protein gene from *Vitis pseudoreticulata* in response to *Plasmopara viticola* infection. *Protoplasma*. 2013; 250: 129–140. doi: [10.1007/s00709-012-0384-8](https://doi.org/10.1007/s00709-012-0384-8) PMID: [22327469](https://pubmed.ncbi.nlm.nih.gov/22327469/).
66. Schlumbaum A, Mauch F, Vogeli U, Boller T. Plant chitinases are potent inhibitors of fungal growth. *Nature*. 1986; 324: 365–367.
67. Holsters M, de Waele D, Depicker A, Messens E, van Montagu M, Schell J. Transfection and transformation of *Agrobacterium tumefaciens*. *Mol Gen Genet*. 1978; 163(2): 181–187. doi: [10.1007/BF00267408](https://doi.org/10.1007/BF00267408) PMID: [355847](https://pubmed.ncbi.nlm.nih.gov/355847/)
68. Paz MM, Shou HX, Guo ZB, Zhang ZY, Banerjee AK, Wang K. Assessment of conditions affecting *Agrobacterium*-mediated soybean transformation using the cotyledonary node explants. *Euphytica*. 2004; 136(2): 167–179. doi: [10.1023/B:EUPH.0000030669.75809.dc](https://doi.org/10.1023/B:EUPH.0000030669.75809.dc)
69. Kim YJ, Jang MG, Lee HJ, Jang GH, Sukweenadhi J, Kwon WS et al. Functional characterization of the pathogenesis-related protein family 10 gene, PgPR10-4, from *Panax ginseng* in response to environmental stresses. *Plant Cell Tiss Org*. 2014; 118: 531–543. doi: [10.1007/s11240-014-0505-5](https://doi.org/10.1007/s11240-014-0505-5)
70. Chacón O, González M, López Y, Portieles R, Pujol M, González E et al. Over-expression of a protein kinase gene enhances the defense of tobacco against *Rhizoctonia solani*. *Gene*. 2010; 452: 54–62. doi: [10.1016/j.gene.2009.11.011](https://doi.org/10.1016/j.gene.2009.11.011) PMID: [20004236](https://pubmed.ncbi.nlm.nih.gov/20004236/).
71. Crowell DN, John ME, Russell D, Amasino RM. Characterization of a stress-induced, developmentally regulated gene family from soybean. *Plant Mol Biol*. 1992; 18(3): 459–466. PMID: [1371403](https://pubmed.ncbi.nlm.nih.gov/1371403/).
72. Samanani N, Liscombe DK, Facchini PJ. Molecular cloning and characterization of norcochlorine synthase, an enzyme catalyzing the first committed step in benzyloquinoline alkaloid biosynthesis. *Plant J*. 2004; 40(2): 302–313. doi: [10.1111/j.1365-3113X.2004.02210.x](https://doi.org/10.1111/j.1365-3113X.2004.02210.x) PMID: [15447655](https://pubmed.ncbi.nlm.nih.gov/15447655/).
73. Robert N, Ferran J, Breda C. Molecular characterization of the incompatible interaction of *Vitis vinifera* leaves with *Pseudomonas syringae* pv. *pisi*: Expression of genes coding for stilbene synthase and class 10 PR protein. *Eur J Plant Pathol*. 2001; 107(2): 249–261.
74. Mellersh DG, Foulds IV, Higgins VJ, Heath MC. H₂O₂ plays different roles in determining penetration failure in three diverse plant-fungal interactions. *Plant J*. 2002; 29: 257–268. doi: [10.1046/j.0960-7412.2001.01215.x](https://doi.org/10.1046/j.0960-7412.2001.01215.x) PMID: [11844104](https://pubmed.ncbi.nlm.nih.gov/11844104/).

75. Knight H, Trewavas AJ, Knight MR. Calcium signaling in *Arabidopsis thaliana* responding to drought and salinity. *Plant J.* 1997; 12(5): 1067–1078. doi: [10.1046/j.1365-313X.1997.12051067.x](https://doi.org/10.1046/j.1365-313X.1997.12051067.x) PMID: [9418048](https://pubmed.ncbi.nlm.nih.gov/9418048/).
76. Wood NT, Allan AC, Haley A, Viry-Moussaïd M, Trewavas AJ. The characterization of differential calcium signaling in tobacco guard cells. *Plant J.* 2000; 24(3): 335–344. doi: [10.1046/j.1365-313x.2000.00881.x](https://doi.org/10.1046/j.1365-313x.2000.00881.x) PMID: [11069707](https://pubmed.ncbi.nlm.nih.gov/11069707/).
77. Odjakova D, Hadjiivanova C. The complexity of pathogen defense in plants. *Bulg. J. Plant physiol.* 2001; 27(1–2): 101–109.
78. Agarwal P, Bhatt V, Singh R, Das M, Sopory SK, Chikara J. Pathogenesis-related gene, *JcPR-10a* from *Jatropha curcas* exhibit RNase and antifungal activity. *Mol. Biotechnol.* 2013; 54(2): 412–415. doi: [10.1007/s12033-012-9579-7](https://doi.org/10.1007/s12033-012-9579-7) PMID: [22763562](https://pubmed.ncbi.nlm.nih.gov/22763562/).
79. Wang L, Wei JY, Zou Y, Xu K, Wang Y, Cui L et al. Molecular Characteristics and biochemical functions of VpPR10s from *Vitis pseudoreticulata* associated with biotic and abiotic stresses. *Int J Mol Sci.* 2014; 15: 19162–19182. doi: [10.3390/ijms151019162](https://doi.org/10.3390/ijms151019162) PMID: [25340981](https://pubmed.ncbi.nlm.nih.gov/25340981/).
80. Bendtsen JD, Lars Juhl Jensen LJ, Blom N, Von Heijne G, Brunak S. Feature-based prediction of non-classical and leaderless protein secretion. *Protein Eng Des Sel.* 2004; 17: 349–356. doi: [10.1093/protein/gzh037](https://doi.org/10.1093/protein/gzh037) PMID: [15115854](https://pubmed.ncbi.nlm.nih.gov/15115854/).
81. Enkerli K, Hahn MG, Mims CW. Ultrastructure of compatible and incompatible interactions of soybean roots infected with the plant pathogenic oomycete *Phytophthora sojae*. *Can J Bot.* 1997; 75(9): 1493–1508. doi: [10.1139/b97-864](https://doi.org/10.1139/b97-864)
82. Lee OR, Pulla RK, Kim YJ, Balusamy SR, Yang DC. Expression and stress tolerance of PR10 genes from *Panax ginseng* C. A Meyer. *Mol Biol Rep.* 2013; 39: 2365–2374. doi: [10.1007/s11033-011-0987-8](https://doi.org/10.1007/s11033-011-0987-8) PMID: [21667108](https://pubmed.ncbi.nlm.nih.gov/21667108/).