

RESEARCH ARTICLE

# Persistent Postural-Perceptual Dizziness: A Matter of Higher, Central Dysfunction?

Dagny Holle<sup>1\*</sup>, Benedict Schulte-Steinberg<sup>1</sup>, Sebastian Wurthmann<sup>1</sup>, Steffen Naegel<sup>1</sup>, Ilya Ayzenberg<sup>2</sup>, Hans-Christoph Diener<sup>1</sup>, Zaza Katsarava<sup>3</sup>, Mark Obermann<sup>1</sup>

**1** Department of Neurology, University of Duisburg-Essen, Essen, Germany, **2** Department of Neurology, Ruhr University Bochum, Bochum, Germany, **3** Department of Neurology, Evangelical hospital Unna, Unna, Germany

\* [dagny.holle@uk-essen.de](mailto:dagny.holle@uk-essen.de)



CrossMark  
click for updates

OPEN ACCESS

**Citation:** Holle D, Schulte-Steinberg B, Wurthmann S, Naegel S, Ayzenberg I, Diener H-C, et al. (2015) Persistent Postural-Perceptual Dizziness: A Matter of Higher, Central Dysfunction? PLoS ONE 10(11): e0142468. doi:10.1371/journal.pone.0142468

**Editor:** Simon J Cropper, University of Melbourne, AUSTRALIA

**Received:** June 16, 2015

**Accepted:** October 22, 2015

**Published:** November 16, 2015

**Copyright:** © 2015 Holle et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** The authors have no support or funding to report.

**Competing Interests:** Dagny Holle has received a research grant from Grünenthal, Allergan and the German Ministry of Education and Research (BMBF). Benedict Schulte-Steinberg reports nothing to disclose. Sebastian Wurthmann reports nothing to disclose. Steffen Naegel reports nothing to disclose. Ilya Ayzenberg reports nothing to disclose. Hans-Christoph Diener has received honoraria for

## Abstract

### Objective

Persistent postural-perceptual dizziness (PPPD) is the most common vestibular disorder in the age group between 30 and 50 years. It is considered to be based on a multisensory maladjustment involving alterations of sensory response pattern including vestibular, visual and motion stimuli. Previous data supported a link between vestibular and pain mechanism. The aim of the study was to investigate whether other sensory inputs such as pain stimuli might be altered in terms of a more widespread central perception dysfunction in this disorder.

### Methods

Nociceptive blink reflex was measured in 27 patients with PPPD and compared with 27 healthy, age and gender matched controls. The habituation of the R2 component of the blink reflex was evaluated as the percentage area-under-the curve (AUC) decrease in ten consecutive blocks of five averaged rectified responses. Additionally, clinical characteristics were evaluated.

### Results

In patients with PPPD a lack of habituation was observed compared to healthy controls. Relative AUC decreased between the first and the tenth block by 19.48% in PPPD patients and by 31.63% ( $p = 0.035$ ) in healthy controls. There was no correlation between clinical data (course of disease, comorbid depression, medication, trigger factors) or electrophysiological data (perception threshold, pain threshold, stimulus intensity) and habituation pattern. No trigeminal sensitization in terms of facilitation of absolute values could be detected.

### Conclusion

Our study results supports the hypothesis of the multisensory dimension of impaired sensory processing in patients with PPPD extends beyond vestibular/visual motion stimuli and

participation in clinical trials, contribution to advisory boards or lectures from Addex Pharma, Allergan, Almirall, AstraZeneca, Bayer Vital, Berlin Chemie, Coherex Medical, CoLucid, Böhringer Ingelheim, Bristol-Myers Squibb, GlaxoSmithKline, Grünenthal, Janssen-Cilag, Lilly, La Roche, 3M Medica, Minster, MSD, Novartis, Johnson & Johnson, Pierre Fabre, Pfizer, Schaper and Brümmer, SanofiAventis, and Weber & Weber; received research support from Allergan, Almirall, AstraZeneca, Bayer, Galaxo-Smith-Kline, Janssen-Cilag, and Pfizer. Headache research at the Department of Neurology in Essen is supported by the German Research Council (DFG), the German Ministry of Education and Research (BMBF), and the European Union. Zaza Katsarava: has received research grants and honoraria from Allergan, Bayer, Biogen and Merck, and is an advisory board member for Allergan. Headache research at the Department of Neurology in Essen is supported by the German Research Council (DFG), the German Ministry of Education and Research (BMBF), and the European Union. Mark Obermann: has received scientific grants by the German Federal Ministry of Education and Research BMBF 01EM 0513. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

reflexive postural/oculomotor control mechanisms to other sensory inputs such as pain perception in terms of a more generalized disturbed habituation pattern.

## Introduction

Persistent postural-perceptual dizziness (PPPD) is one of the most common causes of chronic dizziness in middle-aged patients and the second most common diagnosis in patients presenting with vestibular symptoms[1] with a high impact on functioning in daily living and quality of life.[2] The main clinical characteristics of this disorder include persisting subjective non-rotational vertigo or dizziness, hypersensitivity to motion stimuli, including the patients own movement and motion of objects in the visual surround, as well as difficulties with precision visual tasks.[3] Typically these patients have normal values in clinical balance tests.[4] Objective tests to prove the diagnosis of PPPD do not exist.

The diagnosis of PPPD is based on a clinical condition named phobic postural vertigo (PPV, Phobischer Schwankschwindel) which was initially described by Brandt and Dieterich in the 1980s. [5] In the early 2000s Staab et al. refined and updated the concept of PPV and renamed it chronic subjective dizziness (CSD). [6] In 2014 new consensus criteria were established and the clinical condition was renamed PPPD. PPPD has now been included by the World Health Organization (WHO) in its draft list to be part of the upcoming International Classification of Diseases (ICD-11) in 2017.

The underlying pathophysiologic mechanisms of PPPD are still enigmatic. PPPD often follows an acute vestibular disorder e.g. vestibular neuritis. Although patients recover from the initial acute disorder and clinical and diagnostics tests become normal again PPPD patients keep on having symptoms of dizziness. Newer concepts, therefore, favor a failure of readaptation/habituation of the postural control system in response to acute symptoms of dizziness and vertigo as the main driving force of the disorder. This maladjustment might lead to a misprediction of the sensory consequences of one's own action explaining the main clinical symptom of dizziness and balance problems in these patients.[4,5] Initially useful high-risk postural control strategies (e.g. moving cautiously when suffering from an acute vestibular neuritis) fail to revert to normal when the initial disorder resolves or is compensated.

Clinical and epidemiological observations as well as previous scientific studies support the concept of an overlap between vestibular and pain mechanisms that might be involved in the pathophysiology of PPPD. On one hand pain disorders such as migraine and balance disorders show a high comorbidity.[7,8] On the other hand migrainous vertigo is the second most frequent cause of recurrent vertigo.[9] Optokinetic stimulation -a primarily vestibular stimulus-intensifies triggered trigeminal pain, and pain sensitivity in the fingers in migraineurs [10] and makes migraineurs more susceptible to develop migraine attacks [11]. Painful trigeminal stimulation itself worsens motion sickness [12] and increases nystagmus in migraineurs.[13] Additive effects of processing afferent vestibular and pain pathways in pre-parabrachial and pre-thalamic areas to the amygdala and cerebral cortex might be responsible for these observations. [14,15] Despite central pathways peripheral mechanism might be additionally involved. For example, similar receptor expressions regarding serotonergic, TRPV1 and purinergic receptors can be found in trigeminal, vestibular and spiral ganglion cells (for review [14]).

The aim of this study was to investigate whether hypothesized maladjusted readaptation/habituation in PPPD extends beyond vestibular/visual motion stimuli and reflexive postural/oculomotor control to other sensory inputs such as pain perception in terms of a more generalized disturbed phenomenon. Therefore, we investigated pain stimulation associated

habituation of the R2 component of the nociceptive blink, which reflects short-time adaptation to sensory stimuli,

## Methods

Written and informed consent was obtained from all patients and healthy volunteers before entering the study. The study was performed according the Declaration of Helsinki and the study protocol was approved by the Ethics Committee of the Medical Faculty of the University of Duisburg-Essen, Germany (Approval Number: 13-5482-BO).

## Subjects

Twenty-seven subjects (10 ♂, 17♀) with chronic subjective dizziness according to the diagnostic criteria of Staab et al.[3] ([www.who.int/classifications/icd/revision/en/](http://www.who.int/classifications/icd/revision/en/)) as well as 27 age- and gender-matched healthy volunteers (HV) were included into the study (Table 1).

Patients with prior diagnosis of CSD were recruited from the Dizziness and Vertigo Center Essen. They were then re-evaluated to match the new diagnostic criteria of persistent postural-perceptual dizziness (PPPD). All patients fulfilled both diagnostic criteria (CSD and PPPD). PPPD displays the latest consensus for diagnosis of this clinical condition and the World Health Organization has included PPPD in its draft list of diagnoses to be added to the next edition of the International Classification of Diseases (ICD-11) in 2017.[16]

All participants were assessed face to face and demographic data were obtained with a standardised questionnaire including questions from the German version of the Center for Epidemiologic Studies Depression Scale (CES-D) called "Allgemeine Depressionsskala" (ADS). Migraine was diagnosed based on the current diagnostic criteria (ICHD (International Classification of Headache disorders) 3 beta) by an experienced medical doctor. Depression was diagnosed based on the current ICD-10 criteria and ADS by an experienced medical doctor.

One patient suffered from migraine in his childhood but had no migraine attacks after the age of 20. Analysis without this patient did not change study results significantly. Four patients suffered from mild depression. Analysis without these patients did not change study results significantly.

All subjects were instructed not to drink caffeine or alcohol containing beverages within 4 hours before the recording to avoid contamination of study results.

## Electrophysiological settings

The nociceptive blink reflex (nBR) was elicited as described previously.[17,18] Two planar concentric electrodes (inomed Medizintechnik GmbH Lübeck; [www.walter-graphtek.com](http://www.walter-graphtek.com)) were attached 10mm above the entry zone of the right supraorbital nerve approximately two cm apart. To avoid unnecessary discomfort for the patients, pain stimulation was restricted to the right side. Stimulation was applied with a triple monopolar square wave pulse lasting 0.5ms. The pulse interval was set at 5 ms, interstimulus interval was 12–18 seconds in a pseudorandomized order. Intensity for stimulation was 2 times individual pain threshold, which was determined with two ascending and descending sequences of successive current intensities between 0.2 and 2 mA in 0.2 mA intensity steps. Additionally, the perception threshold was determined. Surface electrodes recorded rectified EMG-responses over the right orbicularis oculi muscle, referenced to the ipsilateral orbital rim (recording parameters: bandwidth 1Hz to 1 kHz, sampling rate 2.5 kHz, sweep length 300ms (1401 plus, Signal, Cambridge Electronic Design UK)). Ten blocks each of 6 stimuli were applied with an interval of 2 minutes between each block. The 2-min interblock interval was chosen as it has been shown to detect the most reliable

**Table 1. Clinical characteristics of patients with persistent postural-perceptual dizziness.**

	Healthy Volunteers (n = 27)	PPPD atien (n = 27)
Age (y)	39.48± 10.20	39.00 ± 8.48
Gender	10 ♂, 17 ♀	10 ♂, 17 ♀
Previous history of migraine	0 (0%)	1 (4%)
Depression	0 (0%)	4 (15%)
Constant dizziness/unsteadiness*	n/a	27 (100%)
Postural relationship**	n/a	27 (100%)
Symptoms since	n/a	
> 3 month		3 (11%)
> 1 year		10 (37%)
2–5 years		7 (26%)
> 5 years		7 (26%)
Context-dependent symptom exacerbation ***	n/a	27 (100%)
Triggering factors (Prior neuro-otologic diseases (e.g. benign paroxysmal positional vertigo, vestibular neuritis), Prior medical problems (e.g. mild traumatic brain injury), Prior psychiatric disorders (e.g. anxiety disorder)	n/a	27 (100%)
Normal Physical examination or minor non-diagnostic abnormalities (e.g. scar-related hypesthesia)	27 (100%)	27 (100%)
Normal vestibular laboratory testing****	27 (100%)	27 (100%)
Normal MRI or minor non-diagnostic abnormalities (e.g. small arachnoidal cyst, not significant white matter lesions)	n/a	27 (100%)
Medication SSRI or SNRI	0%	4 (15%)

Values are means ± SD or n (%) of all patients/healthy volunteers

SSRI: Selective Serotonin Reuptake Inhibitor; SNRI: Serotonin-Noradrenalin Reuptake Inhibitor; n/a: not applicable; SD: standard deviation; n: number of patients or healthy volunteers

\* Dizziness is present throughout the day but fluctuates in severity

\*\* Symptoms are most severe when walking or standing, less severe when sitting, and absent or very minor when recumbent

\*\*\* Provocative factors include active or passive motion that is not related to a specific direction or position, exposure to large-field moving visual stimuli or complex visual stimulation, performance of small-field, precision visual activities such as reading or using a computer

\*\*\*\* Vestibular laboratory testing includes: Videonystagmography (VNG), Vestibular-evoked myogenic potentials (VEMP)

doi:10.1371/journal.pone.0142468.t001


habituation in normal subjects.[19] This habituation paradigm has shown efficacy in previous studies on habituation pattern in migraine patients.[20]

### Data processing

Signal analysis was performed by an investigator blinded to the diagnosis. The first sweep of each block was excluded from further analysis to avoid contamination by the startle response. Area under the curve (AUC) of the R2 response (RA: response area of the R2 response) was analysed in each sweep offline after demeaning, rectification and averaging between 27 and 87ms. Each block was analysed separately, calculating mean values for each block separately. Habituation was defined as percentage change of the AUC between the first and the 10<sup>th</sup> block of recordings.

### Statistical analysis

One-way analysis of variance (ANOVA) was used to calculate group differences in mean pain threshold, mean stimulus intensities and clinical characteristics. ANOVA for repetitive measurements was applied for measurement of habituation. Therefore, different blocks (1 to 10)



**Fig 1. Habituation pattern of the nociceptive blink reflex.** Habituation pattern of the nociceptive blink reflex comparing patients with persistent postural-perceptual dizziness (PPPD) and healthy volunteers (HV) showing reduced habituation in patients with PPPD. ANOVA \*  $p < 0.05$ , \*\*  $p < 0.005$ . Habituation of the R2 response area of right-sided nBR in 10 blocks of five averagings. Second to tenth block are expressed as percentage of the first block

doi:10.1371/journal.pone.0142468.g001

were considered as within subject factors and the diagnostic group (HV and PPPD) as between subject factors. Results are presented as means  $\pm$  standard deviations except in Fig 1 where standard errors were used for better visualization of the different curves. Significance level was set as  $p < 0.05$ .

## Results

There were no significant differences in mean perception threshold and mean pain threshold as well as mean stimulation intensities (2x pain threshold) comparing PPPD patients and HV (Table 2). We found no significant differences in regard to the R2 latency and area under the curve comparing both groups (Table 2).

Habituation of the nBR RA was significantly reduced in patients with PPPD compared to healthy volunteers (HV) (Fig 1).

In HV a decrease in amplitude of 31.63% between the first and the 10<sup>th</sup> block of five averagings was detected whereas PPPD patients showed a decrease in amplitude of 19.48%. Habituation gradually increased over the ten blocks in both groups. The difference of the degree of nBR RA habituation between PPPD patients and HV was significant from the 7<sup>th</sup> to the 10<sup>th</sup> block of stimulation. No correlation between habituation pattern and other clinical characteristics (course of disease, depression, medication, trigger factors) or electrophysiological results (perception threshold, pain threshold, stimulus intensity) could be detected (data not shown).

**Table 2. Mean values of electrophysiological findings (block 1).**

	Healthy Volunteers	PPPD Patients	Significance level
Perception threshold [mA]	0.33 ± 0.13	0.33 ± 0.19	n.s.
Pain threshold [mA]	0.97 ± 0.52	1.05 ± 0.61	n.s.
Stimulation intensities [mA]	1.67 ± 0.62	1.73 ± 0.55	n.s.
AUC [(x10-3)(μV x ms)]	101.19 ± 31.51	125.54 ± 55.42	n.s.
N2 latency [ms]	38.90 ± 5.83	40.86 ± 6.05	n.s.

Values are means ± SD

mA: Milliampere; AUC: Area under the curve; μV: Mikrovolt; ms: Milliseconds; n.s.: not significant

PPPD: Persistent postural-perceptual dizziness

doi:10.1371/journal.pone.0142468.t002

## Discussion

Patients with PPPD showed a habituation deficit of the nBR RA of the nociceptive blink reflex suggesting insufficient sensory information processing that goes beyond vestibular system dysfunction supporting the concept of a multisensory maladjustment in this disorder.

This multisensory approach of the underlying pathophysiology of this clinical condition was initially described by Brandt and Dieterich. They postulated in 1986 that PPV (Phobic postural vertigo) is caused by an inadequate compensation of self-induced sensory stimulation.[5] Due to this pathophysiological concept patients predict sensory consequences of their own motoric actions inadequately leading to a disturbed perceptual stability. Clinically, patients feel a reduced balance and gait control with a fear of falling. This subjective impression of balance instability was detected using a pressure sensitive mat (GAITRite®). Patients with PPV showed an inadequate gait control, which was characterized by a slower walk velocity, reduced cadence and stride length compared to healthy controls.[21] Symptoms even increase when visual control is absent. Furthermore, inappropriate postural strategies involving supraspinal control resulting in increased body sway during normal stance conditions seem also to be involved in this disorder.[22–24] Patients seem to use inappropriate balance strategies during normal stance which are usually used only during difficult and demanding balance tasks, probably due to anxious control of posture.[25,26] Interestingly, during demanding postural tasks such as standing with closed eyes, balance control is comparable to healthy subjects.

Balaban et al. developed in this context the concept of the balance-migraine-anxiety syndrome [15] which might be a suitable pathophysiological framework for PPPD as well and underscores multisensory maladjustment dimension in this clinical condition complex. Based on the clinical observation of increased comorbidity of balance disorders (PPPD/CSD/PPV and neuro-otologic diseases) migraine [27–29] and psychiatric disorders [6,30,31] Balaban et al. hypothesized that additive effects of processing afferent vestibular and pain information in pre-parabrachial and pre-thalamic pathway components to the amygdala and cerebral cortex might be the pathophysiological correlate. [14,15] In line with this hypothesis voxel-based morphometry data showed alterations in vestibular migraine that resemble those of migraine but also extent to areas of the vestibular system.[32] Additionally, functional imagine data of acute migraine attacks also show that the areas of activation also compromise areas of the vestibular central network especially within the dorsal and dorsolateral pons.[33] Despite central effects, peripheral mechanism might also be involved in the overlap of these clinical conditions. Patterns of serotonergic, TRPV1 and purinergic receptor expression are similar in trigeminal, vestibular and spiral (cochlear) ganglion cells.[15] This might be an explanation why similar treatment regimens can be used in this clinical disorders (e.g. antidepressants such as SNRI

and SSRI in PPPD. [34,35], migraine [36] and depression[37]; triptans in migraine[38] and motion sickness [39]).

However, only one of our PPPD patients was suffering from migraine in the past. Therefore, not all patients seem to display the complete clinical picture of a balance-migraine-anxiety syndrome, although pathways within the underlying pathophysiology of this syndrome might be disturbed. Patients may suffer clinically only from one part of the syndrome e.g. balance disorders, but electrophysiological investigation shows alterations extending beyond vestibular/visual motion stimuli and reflexive postural/oculomotor control to other sensory inputs such as pain perception.

NBR habituation is not a pathognomonic feature of PPPD, but can be observed in other disorders such as migraine. [19,20]. Interestingly, lack of habituation in migraineurs is not restricted to pain modalities, but can also be detected in other sensory modalities such as visually evoked potentials[40], somatosensory evoked potentials (SEP)[41], auditory evoked potentials (AEP)[42] and the vestibulo-collic reflex.[43] It was suggested that this observation reflects an alteration of cortical excitability in general, which goes beyond pure pain processing deficits in migraine. Similar mechanisms might be involved in PPPD as well. Our data support the hypothesis, that higher cortical regions might be involved in the underlying pathophysiology of PPPD, regulating perception of sensory information in a more general way as multiple modalities are involved.

In comparison to other nBR studies, especially to Di Clemente et al, who used the same stimulation paradigm[20], the habituation deficit in PPPD patients seems less pronounced than in pain disorders and can only be detected in the final stimulation blocks (7 to 10). In contrast, in migraineurs, the differences of habituation can be observed as early as the second block of averages. In migraineurs it was shown, that habituation deficit normalizes with increasing attack frequency[20,40]. Our PPPD patients were almost constantly suffering from clinical symptoms of dizziness and balance problems, which might be an explanation why habituation is less pronounced than in migraine patients without headaches. Additionally, one could hypothesize that general central maladjustment only leads as a kind of “side effect” to impaired pain habituation in PPPD compared to the main effect, which would be impaired vestibular processing. In contrast, the main effect in pain disorders such as migraine would strike pain perception and would therefore be more pronounced than in PPPD.

The exact physiological mechanism of habituation and its impairment are unknown, yet. Some authors suggest a lower pre-activation level of the sensory cortex to be a pathophysiological correlate especially in migraineurs, who need a larger activation level after repetitive stimulation which was termed “ceiling effect”. [44] Healthy controls, in contrast, reach the top of response activity and, therefore, habituate faster. Other authors hypothesize that the neurotransmitter dopamine might be involved in the mechanism of habituation. This hypothesis is based on the observation that the habituation deficit which can be detected in Parkinson’s disease is reversible with dopaminergic but not with anticholinergic therapy.[45] Therefore, a dopaminergic deficit has been discussed as pathophysiological correlate of a habituation deficit reflecting sensory disinhibition in this disorder. Animal findings indicate that the dopamine D<sub>1</sub> receptor plays a role in the generation of habituation to repetitive stimuli whereas D<sub>2</sub> and D<sub>3</sub> receptors act to inhibit the habituation process.[46] To which extent a dopaminergic deficit might play a role in PPPD and if affected patients might benefit from dopamine substitution is unknown. In therapy, up to now only SSRIs and SNRIs have been investigated in clinical trials and have shown efficacy in PPPD. [34,35] These substances increase dopamine levels despite their intrinsic serotonergic properties.[47]

Some limitations of this study have to be addressed. We did investigate patients only once. Information on the course of disease, especially regarding a possible regeneration of

habituation deficit after the dizziness resolved would be an interesting question. Additionally, analysis of larger subgroups regarding triggering factors, risk factors, medication influences, comorbidities, etc. would be interesting for further analysis as our small patient number per subgroup would probably be too small to detect more subtle differences. Four patients suffered from depression, which might influence habituation. However, exclusion of these patients did not alter the results.

In summary, our study supports the hypothesis of a multisensory dimension of impaired sensory processing in patients with PPPD not only confined to vestibular/visual motion stimuli and reflexive postural/oculomotor control mechanisms but also extended more generally to other sensory inputs and reflexive responses such as pain perception in terms of a disturbed habituation pattern. PPPD might therefore serve as a model supporting the physiological link between vestibular and pain mechanism.

## Supporting Information

**S1 File. Data set.** Raw values of study results of all study participants. (SAV)

## Author Contributions

Conceived and designed the experiments: DH MO HCD. Performed the experiments: DH BSS SW. Analyzed the data: DH SN MO IA. Contributed reagents/materials/analysis tools: ZK IA. Wrote the paper: DH HCD MO SW.

## References

1. Strupp M, Glaser M, Karch C, Rettinger N, Dieterich M, Brandt T. [The most common form of dizziness in middle age: phobic postural vertigo]. *Nervenarzt*. 2003; 74: 911–914. doi: [10.1007/s00115-003-1567-5](https://doi.org/10.1007/s00115-003-1567-5) PMID: [14551699](https://pubmed.ncbi.nlm.nih.gov/14551699/)
2. Best C, Eckhardt-Henn A, Diener G, Bense S, Breuer P, Dieterich M. Interaction of somatoform and vestibular disorders. *J Neurol Neurosurg Psychiatry*. 2006; 77: 658–664. doi: [10.1136/jnnp.2005.072934](https://doi.org/10.1136/jnnp.2005.072934) PMID: [16614028](https://pubmed.ncbi.nlm.nih.gov/16614028/)
3. Staab JP. Chronic subjective dizziness. *Continuum (Minneapolis, Minn)*. 2012; 18: 1118–1141. doi: [10.1212/01.CON.0000421622.56525.58](https://doi.org/10.1212/01.CON.0000421622.56525.58)
4. Brandt T. Phobic postural vertigo. *Neurology*. 1996; 46: 1515–1519. PMID: [8649539](https://pubmed.ncbi.nlm.nih.gov/8649539/)
5. Brandt T, Dieterich M. Phobischer Attacken-Schwankschwindel, ein neues Syndrom. *Munch Med Wochenschr*. 1986; 128: 247–250.
6. Staab JP, Ruckenstein MJ. Expanding the differential diagnosis of chronic dizziness. *Arch Otolaryngol Head Neck Surg*. 2007; 133: 170–176. doi: [10.1001/archotol.133.2.170](https://doi.org/10.1001/archotol.133.2.170) PMID: [17309987](https://pubmed.ncbi.nlm.nih.gov/17309987/)
7. Neuhauser H, Leopold M, von Brevern M, Arnold G, Lempert T. The interrelations of migraine, vertigo, and migrainous vertigo. *Neurology*. 2001; 56: 436–441. PMID: [11222783](https://pubmed.ncbi.nlm.nih.gov/11222783/)
8. Lempert T, Neuhauser H. Epidemiology of vertigo, migraine and vestibular migraine. *J Neurol*. 2009; 256: 333–338. doi: [10.1007/s00415-009-0149-2](https://doi.org/10.1007/s00415-009-0149-2) PMID: [19225823](https://pubmed.ncbi.nlm.nih.gov/19225823/)
9. Neuhauser HK. Epidemiology of vertigo. *Curr Opin Neurol*. 2007; 20: 40–46. doi: [10.1097/WCO.0b013e328013f432](https://doi.org/10.1097/WCO.0b013e328013f432) PMID: [17215687](https://pubmed.ncbi.nlm.nih.gov/17215687/)
10. Drummond PD. Motion sickness and migraine: optokinetic stimulation increases scalp tenderness, pain sensitivity in the fingers and photophobia. *Cephalalgia*. 2002; 22: 117–124. PMID: [11972579](https://pubmed.ncbi.nlm.nih.gov/11972579/)
11. Granston A, Drummond PD. Painful stimulation of the temple during optokinetic stimulation triggers migraine-like attacks in migraine sufferers. *Cephalalgia*. 2005; 25: 219–224. doi: [10.1111/j.1468-2982.2004.00844.x](https://doi.org/10.1111/j.1468-2982.2004.00844.x) PMID: [15689198](https://pubmed.ncbi.nlm.nih.gov/15689198/)
12. Drummond PD, Granston A. Facial pain increases nausea and headache during motion sickness in migraine sufferers. *Brain*. 2004; 127: 526–534. doi: [10.1093/brain/awh061](https://doi.org/10.1093/brain/awh061) PMID: [14749288](https://pubmed.ncbi.nlm.nih.gov/14749288/)
13. Marano E, Marcelli V, Di Stasio E, Bonuso S, Vacca G, Manganeli F, et al. Trigeminal stimulation elicits a peripheral vestibular imbalance in migraine patients. *Headache*. 2005; 45: 325–331. doi: [10.1111/j.1526-4610.2005.05069.x](https://doi.org/10.1111/j.1526-4610.2005.05069.x) PMID: [15836568](https://pubmed.ncbi.nlm.nih.gov/15836568/)

14. Balaban CD. Migraine, vertigo and migrainous vertigo: Links between vestibular and pain mechanisms. *J Vestib Res.* 2011; 21: 315–321. doi: [10.3233/VES-2011-0428](https://doi.org/10.3233/VES-2011-0428) PMID: [22348936](https://pubmed.ncbi.nlm.nih.gov/22348936/)
15. Balaban CD, Jacob RG, Furman JM. Neurologic bases for comorbidity of balance disorders, anxiety disorders and migraine: neurotherapeutic implications. *Expert Rev Neurother.* 2011; 11: 379–394. doi: [10.1586/em.11.19](https://doi.org/10.1586/em.11.19) PMID: [21375443](https://pubmed.ncbi.nlm.nih.gov/21375443/)
16. ICD-11 Beta Draft—Joint Linearization for Mortality and Morbidity Statistics [Internet]. [cited 17 Aug 2015]. Available: <http://apps.who.int/classifications/icd11/browse/l-m/en#/http://id.who.int/icd/entity/2005792829>
17. Kaube H, Katsarava Z, Käufer T, Diener H, Ellrich J. A new method to increase nociception specificity of the human blink reflex. *Clin Neurophysiol.* 2000; 111: 413–416. PMID: [10699400](https://pubmed.ncbi.nlm.nih.gov/10699400/)
18. Katsarava Z, Ellrich J, Diener HC, Kaube H. Optimized stimulation and recording parameters of human “nociception specific” blink reflex recordings. *Clin Neurophysiol.* 2002; 113: 1932–1936. PMID: [12464330](https://pubmed.ncbi.nlm.nih.gov/12464330/)
19. Katsarava Z, Giffin N, Diener H-C, Kaube H. Abnormal habituation of “nociceptive” blink reflex in migraine—evidence for increased excitability of trigeminal nociception. *Cephalalgia.* 2003; 23: 814–819. PMID: [14510928](https://pubmed.ncbi.nlm.nih.gov/14510928/)
20. Di Clemente L, Coppola G, Magis D, Fumal A, De Pasqua V, Di Piero V, et al. Interictal habituation deficit of the nociceptive blink reflex: an endophenotypic marker for presymptomatic migraine? *Brain.* 2007; 130: 765–770. doi: [10.1093/brain/awl351](https://doi.org/10.1093/brain/awl351) PMID: [17251239](https://pubmed.ncbi.nlm.nih.gov/17251239/)
21. Schniepp R, Wuehr M, Huth S, Pradhan C, Brandt T, Jahn K. Gait characteristics of patients with phobic postural vertigo: effects of fear of falling, attention, and visual input. *J Neurol.* 2014; 261: 738–746. doi: [10.1007/s00415-014-7259-1](https://doi.org/10.1007/s00415-014-7259-1) PMID: [24519356](https://pubmed.ncbi.nlm.nih.gov/24519356/)
22. Holmberg J, Tjernström F, Karlberg M, Fransson PA, Magnusson M. Reduced postural differences between phobic postural vertigo patients and healthy subjects during a postural threat. *J Neurol.* 2009; 256: 1258–1262. doi: [10.1007/s00415-009-5110-x](https://doi.org/10.1007/s00415-009-5110-x) PMID: [19363634](https://pubmed.ncbi.nlm.nih.gov/19363634/)
23. Kapfhammer HP, Mayer C, Hock U, Huppert D, Dieterich M, Brandt T. Course of illness in phobic postural vertigo. *Acta Neurol Scand.* 1997; 95: 23–28. PMID: [9048981](https://pubmed.ncbi.nlm.nih.gov/9048981/)
24. Krafczyk S, Schlamp V, Dieterich M, Haberhauer P, Brandt T. Increased body sway at 3.5–8 Hz in patients with phobic postural vertigo. *Neurosci Lett.* 1999; 259: 149–152. PMID: [10025580](https://pubmed.ncbi.nlm.nih.gov/10025580/)
25. Schniepp R, Wuehr M, Pradhan C, Novozhilov S, Krafczyk S, Brandt T, et al. Nonlinear variability of body sway in patients with phobic postural vertigo. *Front Neurol.* 2013; 4: 115. doi: [10.3389/fneur.2013.00115](https://doi.org/10.3389/fneur.2013.00115) PMID: [23966974](https://pubmed.ncbi.nlm.nih.gov/23966974/)
26. Wuehr M, Pradhan C, Novozhilov S, Krafczyk S, Brandt T, Jahn K, et al. Inadequate interaction between open- and closed-loop postural control in phobic postural vertigo. *J Neurol.* 2013; 260: 1314–1323. doi: [10.1007/s00415-012-6797-7](https://doi.org/10.1007/s00415-012-6797-7) PMID: [23263595](https://pubmed.ncbi.nlm.nih.gov/23263595/)
27. Furman JM, Balaban CD, Jacob RG, Marcus DA. Migraine-anxiety related dizziness (MARD): a new disorder? *J Neurol Neurosurg Psychiatr.* 2005; 76: 1–8. doi: [10.1136/jnnp.2004.048926](https://doi.org/10.1136/jnnp.2004.048926) PMID: [15607984](https://pubmed.ncbi.nlm.nih.gov/15607984/)
28. Furman JM, Marcus DA, Balaban CD. Migrainous vertigo: development of a pathogenetic model and structured diagnostic interview. *Curr Opin Neurol.* 2003; 16: 5–13. doi: [10.1097/01.wco.0000053582.70044.e2](https://doi.org/10.1097/01.wco.0000053582.70044.e2) PMID: [12544852](https://pubmed.ncbi.nlm.nih.gov/12544852/)
29. Marcus DA, Furman JM, Balaban CD. Motion sickness in migraine sufferers. *Expert Opin Pharmacother.* 2005; 6: 2691–2697. doi: [10.1517/14656566.6.15.2691](https://doi.org/10.1517/14656566.6.15.2691) PMID: [16316307](https://pubmed.ncbi.nlm.nih.gov/16316307/)
30. Furman JM, Jacob RG. A clinical taxonomy of dizziness and anxiety in the otoneurological setting. *J Anxiety Disord.* 2001; 15: 9–26. PMID: [11388360](https://pubmed.ncbi.nlm.nih.gov/11388360/)
31. Staab JP. Chronic dizziness: the interface between psychiatry and neuro-otology. *Curr Opin Neurol.* 2006; 19: 41–48. PMID: [16415676](https://pubmed.ncbi.nlm.nih.gov/16415676/)
32. Obermann M, Wurthmann S, Steinberg BS, Theysohn N, Diener H-C, Naegel S. Central vestibular system modulation in vestibular migraine. *Cephalalgia.* 2014; 34: 1053–1061. doi: [10.1177/0333102414527650](https://doi.org/10.1177/0333102414527650) PMID: [24662323](https://pubmed.ncbi.nlm.nih.gov/24662323/)
33. May A, Matharu M. New insights into migraine: application of functional and structural imaging. *Curr Opin Neurol.* 2007; 20: 306–309. doi: [10.1097/WCO.0b013e328136c20b](https://doi.org/10.1097/WCO.0b013e328136c20b) PMID: [17495625](https://pubmed.ncbi.nlm.nih.gov/17495625/)
34. Ruckenstein MJ, Staab JP. Chronic subjective dizziness. *Otolaryngol Clin North Am.* 2009; 42: 71–77, ix. doi: [10.1016/j.otc.2008.09.011](https://doi.org/10.1016/j.otc.2008.09.011) PMID: [19134491](https://pubmed.ncbi.nlm.nih.gov/19134491/)
35. Staab JP. Clinical clues to a dizzying headache. *J Vestib Res.* 2011; 21: 331–340. doi: [10.3233/VES-2011-0426](https://doi.org/10.3233/VES-2011-0426) PMID: [22348938](https://pubmed.ncbi.nlm.nih.gov/22348938/)

36. Jackson JL, Cogbill E, Santana-Davila R, Eldredge C, Collier W, Gradall A, et al. A Comparative Effectiveness Meta-Analysis of Drugs for the Prophylaxis of Migraine Headache. *PLoS ONE*. 2015; 10: e0130733. doi: [10.1371/journal.pone.0130733](https://doi.org/10.1371/journal.pone.0130733) PMID: [26172390](https://pubmed.ncbi.nlm.nih.gov/26172390/)
37. Linde K, Kriston L, R ucker G, Jamil S, Schumann I, Meissner K, et al. Efficacy and acceptability of pharmacological treatments for depressive disorders in primary care: systematic review and network meta-analysis. *Ann Fam Med*. 2015; 13: 69–79. doi: [10.1370/afm.1687](https://doi.org/10.1370/afm.1687) PMID: [25583895](https://pubmed.ncbi.nlm.nih.gov/25583895/)
38. Marmura MJ, Silberstein SD, Schwedt TJ. The acute treatment of migraine in adults: the american headache society evidence assessment of migraine pharmacotherapies. *Headache*. 2015; 55: 3–20. doi: [10.1111/head.12499](https://doi.org/10.1111/head.12499) PMID: [25600718](https://pubmed.ncbi.nlm.nih.gov/25600718/)
39. Furman JM, Marcus DA, Balaban CD. Rizatriptan reduces vestibular-induced motion sickness in migraineurs. *J Headache Pain*. 2011; 12: 81–88. doi: [10.1007/s10194-010-0250-z](https://doi.org/10.1007/s10194-010-0250-z) PMID: [20862509](https://pubmed.ncbi.nlm.nih.gov/20862509/)
40. Di Clemente L, Coppola G, Magis D, Fumal A, De Pasqua V, Schoenen J. Nociceptive blink reflex and visual evoked potential habituations are correlated in migraine. *Headache*. 2005; 45: 1388–1393. doi: [10.1111/j.1526-4610.2005.00271.x](https://doi.org/10.1111/j.1526-4610.2005.00271.x) PMID: [16324171](https://pubmed.ncbi.nlm.nih.gov/16324171/)
41. Restuccia D, Vollono C, Virdis D, del Piero I, Martucci L, Zanini S. Patterns of habituation and clinical fluctuations in migraine. *Cephalalgia*. 2014; 34: 201–210. doi: [10.1177/0333102413508241](https://doi.org/10.1177/0333102413508241) PMID: [24121287](https://pubmed.ncbi.nlm.nih.gov/24121287/)
42. Sand T, Zhitniy N, White LR, Stovner LJ. Brainstem auditory-evoked potential habituation and intensity-dependence related to serotonin metabolism in migraine: a longitudinal study. *Clin Neurophysiol*. 2008; 119: 1190–1200. doi: [10.1016/j.clinph.2008.01.007](https://doi.org/10.1016/j.clinph.2008.01.007) PMID: [18316245](https://pubmed.ncbi.nlm.nih.gov/18316245/)
43. Allena M, Magis D, De Pasqua V, Schoenen J, Bisdorff AR. The vestibulo-colic reflex is abnormal in migraine. *Cephalalgia*. 2007; 27: 1150–1155. doi: [10.1111/j.1468-2982.2007.01414.x](https://doi.org/10.1111/j.1468-2982.2007.01414.x) PMID: [17784856](https://pubmed.ncbi.nlm.nih.gov/17784856/)
44. Schoenen J. Deficient habituation of evoked cortical potentials in migraine: a link between brain biology, behavior and trigeminovascular activation? *Biomed Pharmacother*. 1996; 50: 71–78. doi: [10.1016/0753-3322\(96\)84716-0](https://doi.org/10.1016/0753-3322(96)84716-0) PMID: [8761712](https://pubmed.ncbi.nlm.nih.gov/8761712/)
45. Penders CA, Delwaide PJ. Blink reflex studies in patients with Parkinsonism before and during therapy. *J Neurol Neurosurg Psychiatr*. 1971; 34: 674–678. PMID: [5158781](https://pubmed.ncbi.nlm.nih.gov/5158781/)
46. Halberstadt AL, Geyer MA. Habituation and Sensitization of Acoustic Startle: Opposite Influences of Dopamine D1 and D2-family Receptors. *Neurobiol Learn Mem*. 2009; 92: 243–248. doi: [10.1016/j.nlm.2008.05.015](https://doi.org/10.1016/j.nlm.2008.05.015) PMID: [18644244](https://pubmed.ncbi.nlm.nih.gov/18644244/)
47. Zhou F-M, Liang Y, Salas R, Zhang L, De Biasi M, Dani JA. Corelease of dopamine and serotonin from striatal dopamine terminals. *Neuron*. 2005; 46: 65–74. doi: [10.1016/j.neuron.2005.02.010](https://doi.org/10.1016/j.neuron.2005.02.010) PMID: [15820694](https://pubmed.ncbi.nlm.nih.gov/15820694/)