

RESEARCH ARTICLE

Escherichia coli Heat-Stable Enterotoxin Mediates Na⁺/H⁺ Exchanger 4 Inhibition Involving cAMP in T₈₄ Human Intestinal Epithelial Cells

Ana R. Beltrán^{1,2}, Luciene R. Carraro-Lacroix³, Camila N. A. Bezerra³, Marcelo Cornejo¹, Katrina Norambuena¹, Fernando Toledo⁴, Joaquín Araos⁷, Fabián Pardo⁷, Andrea Leiva⁷, Carlos Sanhueza⁷, Gerhard Malnic³, Luis Sobrevia^{5,6,7*}, Marco A. Ramírez^{1,7*}

1 Cellular Physiology Laboratory, Biomedical Department, Faculty of Health Sciences, Universidad de Antofagasta, Antofagasta 1270300, Chile, **2** Department of Education, Faculty of Education, Universidad de Antofagasta, Antofagasta 1270300, Chile, **3** Department of Physiology and Biophysics, Institute of Biomedical Sciences, University of São Paulo, São Paulo 3550308–1009, Brazil, **4** Department of Basic Sciences, Faculty of Sciences, Universidad del Bío-Bío, Chillán 3780000, Chile, **5** Department of Physiology, Faculty of Pharmacy, Universidad de Sevilla, Seville E-41012, Spain, **6** University of Queensland Centre for Clinical Research (UQCCR), Faculty of Medicine and Biomedical Sciences, University of Queensland, Herston, QLD 4029, Queensland, Australia, **7** Cellular and Molecular Physiology Laboratory (CMPL), Division of Obstetrics and Gynecology, Faculty of Medicine, School of Medicine, Pontificia Universidad Católica de Chile, Santiago 8330024, Chile

* marco.ramirez@uantof.cl (MAR); sobrevia@med.puc.cl (LS)



OPEN ACCESS

Citation: Beltrán AR, Carraro-Lacroix LR, Bezerra CNA, Cornejo M, Norambuena K, Toledo F, et al. (2015) *Escherichia coli* Heat-Stable Enterotoxin Mediates Na⁺/H⁺ Exchanger 4 Inhibition Involving cAMP in T₈₄ Human Intestinal Epithelial Cells. PLoS ONE 10(12): e0146042. doi:10.1371/journal.pone.0146042

Editor: Xiaodong Cheng, The University of Texas Health Science Center, UNITED STATES

Received: October 5, 2015

Accepted: December 11, 2015

Published: December 29, 2015

Copyright: © 2015 Beltrán et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: Semillero Dirección de Investigación Universidad de Antofagasta (5309, 5313), Chile, <http://www.uantof.cl>. Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP, Brazil, <http://www.fapesp.br>. Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT 1150377, 1150344, 3140516, 11150083), Chile, <http://www.conicyt.cl/fondecyt/>.

Abstract

The enterotoxigenic *Escherichia coli* strains lead to diarrhoea in humans due to heat-labile and heat-stable (STa) enterotoxins. STa increases Cl⁻ release in intestinal cells, including the human colonic carcinoma T₈₄ cell line, involving increased cGMP and membrane alkalization due to reduced Na⁺/H⁺ exchangers (NHEs) activity. Since NHEs modulate intracellular pH (pH_i), and NHE1, NHE2, and NHE4 are expressed in T₈₄ cells, we characterized the STa role as modulator of these exchangers. pH_i was assayed by the NH₄Cl pulse technique and measured by fluorescence microscopy in BCECF–preloaded cells. pH_i recovery rate (*dpH_i/dt*) was determined in the absence or presence of 0.25 μmol/L STa (30 minutes), 25 μmol/L HOE-694 (concentration inhibiting NHE1 and NHE2), 500 μmol/L sodium nitroprusside (SNP, spontaneous nitric oxide donor), 100 μmol/L dibutyryl cyclic GMP (db-cGMP), 100 nmol/L H89 (protein kinase A inhibitor), or 10 μmol/L forskolin (adenylyl cyclase activator). cGMP and cAMP were measured in cell extracts by radioimmunoassay, and buffering capacity (*β_i*) and H⁺ efflux (*J_{H⁺}*) was determined. NHE4 protein abundance was determined by western blotting. STa and HOE-694 caused comparable reduction in *dpH_i/dt* and *J_{H⁺}* (~63%), without altering basal pH_i (range 7.144–7.172). STa did not alter *β_i* value in a range of 1.6 pH_i units. The *dpH_i/dt* and *J_{H⁺}* was almost abolished (~94% inhibition) by STa + HOE-694. STa effect was unaltered by db-cGMP or SNP. However, STa and forskolin increased cAMP level. STa–decreased *dpH_i/dt* and *J_{H⁺}* was mimicked by forskolin, and STa + HOE-694 effect was abolished by H89. Thus, incubation of T₈₄ cells with STa results in reduced NHE4 activity

Competing Interests: The authors have declared that no competing interests exist.

leading to a lower capacity of pH_i recovery requiring cAMP, but not cGMP. STa effect results in a causal phenomenon (STa/increased cAMP/increased PKA activity/reduced NHE4 activity) ending with intracellular acidification that could have consequences in the gastrointestinal cells function promoting human diarrhoea.

Introduction

Intestinal colon cells are polarized epithelial cells that express a wide range of plasma membrane transporters for a variety of substrates. Membrane transporters at the apical border of these cells promote absorption and release of nutrients, electrolytes and water from and to the intestinal lumen. However, membrane transporters at the basolateral border maintain cell homeostasis by the release of these and other nutrients to the interstitium. The apical membrane of intestinal colon cells is directly exposed to agents and toxins, including the enterotoxigenic *Escherichia coli* (ETEC) strains, an intestinal agent leading to diarrhoea in humans [1]. ETEC colonizes host intestines and releases heat-labile and/or heat-stable (STa) enterotoxins. STa causes secretory diarrhoea and is responsible for about half of all ETEC-related diarrhoeal diseases, including traveller's diarrhoea and epidemic diarrhoea of the newborn [1–5].

STa binds to guanylyl cyclase-C (GC-C) receptors expressed in intestine, kidney, testis and lung, leading to an increase in the intracellular cGMP level [6–8]. STa also increases chloride secretion in a cAMP-dependent manner via the cystic fibrosis transmembrane conductance regulator (CFTR) channels in rat jejunum [9]. In an early study, STa was shown to cause mucosal alkalization due to inhibition of the Na^+/H^+ exchange in rat duodenum [10,11]. However, there are not reports addressing whether this enterotoxin modulates intracellular pH (pH_i), and whether this phenomenon would involve Na^+/H^+ exchangers (NHEs) activity. Since both cGMP and cAMP decrease NHEs activity [12,13], an increase in the intracellular pH (pH_i) in response to STa is expected.

NHEs are key in the modulation of intracellular pH (pH_i), and are differentially expressed and regulated in intestine epithelial cells [14–17]. At least 11 isoforms of the NHEs family have been identified, out of which NHE1, 2, 3, and 4 are expressed in gastrointestinal membranes [16,17]. NHE4 is highly expressed in the stomach, renal cortex and medulla, ureter, skeletal muscle, heart, liver, and spleen [18]. NHE4 is involved in gastric secretion [19] and plays a large role in controlling pH_i [20]. Indeed, NHE4 was identified in the human colon carcinoma cell line T₈₄ [21] and in human colonic crypts [13]. This exchanger isoform modulates plays a determinant role in maintaining pH_i homeostasis; however, nothing is known about the regulation of NHE4 activity in T₈₄ cells by ETEC-released STa. Since T₈₄ cells express the GC-C receptors for STa [22], we hypothesize that STa modulates NHE4 activity and the signalling pathways involved in this phenomenon in this cell type. Our findings suggest that STa decreases NHE4 activity, without altering its protein expression via a mechanism that requires cAMP. This could be determinant in the planning of future therapies for human diarrhoea.

Materials and Methods

Cell culture

The cell line T₈₄ derived from colonic adenocarcinoma of male adult human were purchased from the American Type Culture Collection (ATCC, Rockville, MD, USA) and used for the experiments. T₈₄ cells in culture (5% CO₂, 37°C, pH 7.4) were maintained in Dulbecco's

modified Eagle's medium F12 (DMEM/F12, Gibco, Grand Island, NY, USA) containing low (5 mmol/L) D-glucose and supplemented with 14.5 mmol/L NaHCO₃, 3.2 mmol/L D-glutamine, 15 mmol/L HEPES, 5% foetal calf serum (FCS), 100 IU/mL penicillin and 100 mg/mL streptomycin (hereafter referred as primary culture medium (PCM)) as described [21]. Cells were harvested with trypsin/EGTA (0.25/0.2%, 3 minutes, 37°C) and seeded on sterile glass coverslips or 24 well plates for further 72 hours culture until confluence. Cells were then rinsed (3 times) with PCM containing 0.2% FCS (low-FCS/PCM) and cultured in this medium for further 48 hours in order to obtain a cell cycle synchronized culture.

Measurement of pH_i

T₈₄ cell monolayers in a glass coverslip were mounted in a thermoregulated chamber on an inverted microscope (Nikon Diaphot-TMD, Tokyo, Japan). The cells were incubated for 10 minutes at 37°C with the fluorescent pH sensitive probe 2,7-bicarboxyethyl-5,6-carboxy-fluorescein acetoxymethyl ester (BCECF-AM, 12 μmol/L) (Molecular Probes, Eugene, OR, USA), as described [21]. Cells were then superfused by gravity at 3 mL/minute (37°C) with the control solutions (CS) ((mmol/L) NaCl 141, KCl 5, CaCl₂ 1, KH₂PO₄ 0.4, MgCl₂ 0.5, MgSO₄ 0.4, Na₂HPO₄ 0.3, HEPES 10, D-glucose 0.6 (pH 7.4, 37°C)) using an electromechanic switching system (Heater and Valve Controller, Yale University Electronics Shop, New Haven, CT, USA). The pH_i was calculated from fluorescence ratios measured at excitation of 495/440 nm and emission at 520 nm using a Georgia Instruments PMT-400 photomultiplier system, as described [23]. An area of 260 μm diameter was read, including approximately 200–300 cells. Measurements were performed at 2.5–seconds interval for a period of 300 milliseconds per measurement. The pH_i was calibrated using 10 μmol/L nigericin in a calibrating solution ((mmol/L) KCl 130, NaCl 20, CaCl₂ 1, MgCl₂ 1, HEPES 5 (pH 6.0, 7.0 and 8.0)) as described [21].

pH_i recovery

The pH_i recovery was examined by applying the NH₄Cl pulse technique [21,23,24]. In brief, BCECF-AM loaded cells were superfused with CS until the basal pH_i was stabilized (~15 minutes). T₈₄ cells were preincubated with 0.1, 0.25 or 0.75 μmol/L STa for 30 minutes in the presence of 25 μmol/L HOE-694 (a concentration that inhibits NHE1 and NHE2 activity), as described [21,25,26]. The cells were then exposed (2 minutes) to CS supplemented with NH₄Cl (NH₄Cl/CS solution) ((mmol/L) NaCl 121, KCl 5.4, CaCl₂ 1, KH₂PO₄ 0.4, MgCl₂ 0.5, MgSO₄ 0.4, Na₂HPO₄ 0.3, HEPES 10, D-glucose 0.6, NH₄Cl 20 (pH 7.4, 37°C)). After this incubation period the NH₄Cl/CS solution was replaced by rinsing the cells with CS free of NH₄Cl, without or with 25 μmol/L HOE-694, 500 μmol/L sodium nitroprusside (SNP, spontaneous nitric oxide donor) [27], 100 μmol/L dibutyryl cyclic GMP (db-cGMP), 100 nmol/L H89 (a protein kinase A inhibitor) [28] or 10 μmol/L forskolin (an activator of adenylyl cyclase) [29].

Initial rates of pH_i recovery (dpH_i/dt) were calculated from data collected for the first 60 seconds of the recovery (i.e., after removing the NH₄Cl load) and fitted by a first order lineal regression as described [21,24]. The results were expressed in pH_i units/minute. The fraction of dpH_i/dt mediated by NHE4 ($^{NHE4}dpH_i/dt$) was estimated by the expression:

$$^{NHE4}dpH_i/dt = (^{Total}dpH_i/dt) - (^{HOE}dpH_i/dt)$$

where $^{Total}dpH_i/dt$ is the dpH_i/dt estimated in the absence of HOE-694 (i.e., total initial rate), and $^{HOE}dpH_i/dt$ is the dpH_i/dt estimated in the presence of HOE-694, i.e., under inhibition of NHE1 and NHE2 [21]. The relative effect of STa on $^{NHE4}dpH_i/dt$ (STa^{RE}) was determined by

the expression:

$$STa^{RE} = 100 \times \left(\frac{^{STaNHE4} dpHi/dt}{^{NHE4} dpHi/dt} \right)$$

where $^{STa-NHE4} dpHi/dt$ is $^{NHE4} dpHi/dt$ measured in the presence of STa.

Intrinsic buffering capacity

The ability of intrinsic cellular components to buffer changes in pH_i , i.e., intracellular buffer capacity (β_i), was measured as described [21,24]. After determining the basal pH_i the cells were incubated in a 0.5 mmol/L KCl-containing Na^+ -free CS (Na^+ /CS) ((mmol/L) N-methyl-D-glucamine (NMDG) 120, KCl 5, $CaCl_2$ 1.8, $MgCl_2$ 1, HEPES 30, D-glucose 5 (pH 7.4, 37°C)). Cells were then incubated in the latter solution containing decreasing concentrations of NH_4Cl (50, 20, 10, 5, 2.5 or 1 mmol/L). The β_i ($Beta(i)$) was calculated from the expression:

$$Beta(i) = \frac{\text{change } [NH_4^+]_i}{\text{change } (pH_i)}$$

where the intracellular NH_4^+ concentration ($[NH_4^+]_i$) was obtained from the Henderson-Hasselbalch equation on the assumption that $[NH_3]_i$ (intracellular NH_3) was equivalent to $[NH_3]_o$ (extracellular NH_3), and $\text{change } (pH_i)$ is the fraction of change in units of pH_i value. Knowing the $dpHi/dt$ and β_i values, the rate of overall transmembrane H^+ flux (J_{H^+}) was calculated from the following expression:

$$J_{H^+} = Beta(i) \times \left(\frac{dpHi}{dt} \right)$$

cAMP and cGMP determination

T_{84} cells were cultured to confluence in 98-well plates. Cells were first treated for 10 minutes with 1 mmol/L 3-isobutyl-1-methylxanthine (IBMX) (Sigma-Aldrich, St. Louis, MO, USA) and next incubated for another 10 minutes with culture medium containing IBMX or IBMX and STa or forskolin. cAMP and cGMP levels were measured by enzyme immunoassay (cAMP or cGMP Direct Biotrak EIA, GE Healthcare, PA, USA) according to manufacturer's instructions. Values of cAMP or cGMP were normalized to total cell protein per well.

Western blotting

Total protein was obtained from confluent T_{84} cells rinsed (x2) with ice-cold PBS and harvested in 100 μ L of lysis buffer (10% SDS, 20% glycerol, 100 mmol/L dithiothreitol, 2.9 mmol/L Tris (pH 6.8), 0.1% bromophenol blue) (63.7 mmol/L Tris/HCl (pH 6.8), 10% glycerol, 2% sodium dodecylsulphate, 1 mmol/L Na_3VO_4 , 50 mg/mL leupeptin, 5% β -mercaptoethanol) as described [21,27]. Cells were sonicated (6 cycles, 5 seconds, 100 W, 4°C) and total protein was isolated by centrifugation (13500 g, 15 minutes, 4°C). Proteins (50 μ g) were separated by polyacrylamide gel (7.5%) electrophoresis, transferred to Immobilon-P polyvinylidene difluoride membranes (BioRad Laboratories, Hertfordshire, UK) and probed with primary monoclonal rabbit *anti*-NHE1 (1:1000 dilution, 12 hours, 4°C), primary polyclonal rabbit *anti*-NHE2 (1:1000 dilution, 12 hours, 4°C) (Abcam, Cambridge, UK), primary rat *anti*-NHE4 antibody (11H11, amino acids 565–675, ~55 kDa) (kindly donated by Dr Daniel Biemesderfer from Yale School of Medicine, New Haven, CT, USA) (1:1000 dilution, 2 hours, 22°C), or monoclonal mouse *anti*- β -actin (1:5000 dilution, internal reference) (Santa Cruz Biotechnology, Santa

Cruz, CA, USA) antibodies. The membranes were rinsed in Tris buffer saline-0.1% Tween 20 (TBS-T) and further incubated (1 hour) in TBS-T/0.2% bovine serum albumin (BSA) containing secondary horseradish peroxidase-conjugated goat *anti-rat* or *anti-mouse* antibodies (Thermo Scientific, Rockford, IL, USA). Proteins were detected by enhanced chemiluminescence (film exposure time was 1 minute) in a ChemiDoc-It 510 Imagen System (UVP, LCC Upland, CA, USA) and quantified by densitometry [27,30].

Statistical analysis

The values are mean \pm S.E.M., where n indicates number of different cell cultures ($n = 27$ for STa-untreated (i.e., control) and 25 STa-treated cells) with 3–4 replicates per experiment. The normality of the data (i.e., parametric) was confirmed with Kolmogorov-Smirnov's test. The variances across the control and STa-treated cells under Bartlett's test were homogeneous. Comparisons between two groups were performed by means of Student's unpaired t -test. The difference between more than two groups were performed by analysis of variance (ANOVA, one or two-ways). If the ANOVA demonstrated a significant interaction between variables, *post hoc* analyses were performed by the multiple-comparison Bonferroni test. The experimenter running the assays was blinded to the groups allocation before and during the experiments, and when assessing the outcome (i.e., around 30 days). The statistical software GraphPad InStat 3.0b and GraphPad Prism 7.0a.65 (GraphPad Software Inc., San Diego, CA, USA) were used for data analysis. $P < 0.05$ was considered statistically significant.

Results

Effect of STa on pH_i values

Basal pH_i in T_{84} cells detected in this study was comparable to previous reports in this cell type [21,31,32] and was unaltered in cells preincubated with STa (Table 1, Fig 1). Following the NH_4Cl pulse the acidic pH_i values detected in the cells exposed to STa or HOE-694 were partially restored (27 ± 3 or $55 \pm 6\%$, respectively) compared with cells in the absence of these agents (Fig 1). When cells were coincubated with STa + HOE-694 the NH_4Cl -induced acidic pH_i was only minimally restored ($9 \pm 1\%$).

Effect of STa on pH_i recovery kinetics

Since T_{84} cells express NHE1, NHE2 and NHE4, but not NHE3 forms [21,33], we assayed which of these forms was involved in STa effect on dpH_i/dt . The dpH_i/dt values in the presence of STa or HOE-694 were lower (65 ± 7 or $62 \pm 6\%$, respectively) when compared with cells in the absence of these molecules (Table 1). Coincubation of cells with STa + HOE-694 resulted in higher reduction ($90 \pm 6\%$) in the dpH_i/dt compared with the reduction seen in cells treated with STa or HOE-694 alone.

Effect of STa on β_i and J_{H^+}

The β_i value detected in T_{84} cells in the absence of STa (31.1 ± 2.5 (mmol/L)/intracellular pH units) was similar to that previously reported for this cell type under the same culture and measurement conditions (~ 31 (mmol/L)/intracellular pH units) [21]. Change in β_i value was not significantly altered by $0.25 \mu\text{mol/L}$ STa in a range of 1.6 pH_i units in T_{84} cells. Parallel assays show that cells treated with STa exhibit decreased J_{H^+} ($60 \pm 7\%$) compared with cells in the absence of this toxin (Fig 2A). Since maximal inhibitory effect on this parameter was achieved with $0.25 \mu\text{mol/L}$ STa in the presence of $25 \mu\text{mol/L}$ HOE-694 (Fig 2B), this concentration was used in all subsequent experiments. HOE-694 caused a decrease in J_{H^+} ($56 \pm 7\%$) that was

Table 1. Modulation of intracellular pH by STa in T₈₄ cells.

	<i>pHi</i>	<i>dpHi/dt</i>
Control	7.170 ± 0.028	0.133 ± 0.009
STa	7.144 ± 0.019	0.046 ± 0.009 *
HOE-694	7.172 ± 0.034	0.051 ± 0.010 *
HOE-694 + STa	7.130 ± 0.046	0.014 ± 0.001 *†
Forskolin	7.156 ± 0.021	0.051 ± 0.010 *
Forskolin + STa	7.171 ± 0.030	0.048 ± 0.009 *
Forskolin + HOE-694	7.161 ± 0.050	0.017 ± 0.003 *‡
Forskolin + HOE-694 + STa	7.125 ± 0.061	0.016 ± 0.002 *‡
H89 + HOE-694 + STa	7.143 ± 0.038	0.046 ± 0.008 *
db-cGMP	7.21 ± 0.054	0.110 ± 0.012
db-cGMP + STa	7.10 ± 0.021	0.050 ± 0.012 *§
db-cGMP + HOE-694	7.19 ± 0.053	0.057 ± 0.002 *§
db-cGMP + HOE-694 + STa	7.14 ± 0.051	0.015 ± 0.001 *§¶
SNP	7.16 ± 0.026	0.123 ± 0.009
SNP + STa	7.15 ± 0.021	0.045 ± 0.011 *&
SNP + HOE-694	7.11 ± 0.024	0.047 ± 0.009 *&
SNP + HOE-694 + STa	7.14 ± 0.052	0.015 ± 0.012 *&§

The intracellular pH (*pHi*) was measured in BCECF-AM–preloaded T₈₄ cells as described in Methods. Cells were also subjected to an acid pulse (NH₄Cl assay) and the initial rates of *pHi* recovery (*dpHi/dt*) was measured in cells in the absence (Control) or presence (30 minutes) of 0.25 μmol/L heat-stable (STa) enterotoxin, 25 μmol/L HOE-694 (Na⁺/H⁺ exchangers inhibitor), 10 μmol/L forskolin, 100 nmol/L H89 (protein kinase A inhibitor), 100 μmol/L dibutyl cyclic GMP (db-cGMP), or 500 μmol/L sodium nitroprusside (SNP). STa at 0.1 and 0.75 μmol/L did not alter *pHi* values (7.121 ± 0.011 and 7.160 ± 0.014, respectively; *P*>0.05, *n* = 4). STa at 0.1 μmol/L partially reduced *dpHi/dt* value (0.098 ± 0.005 *pHi* units/minute, *P*<0.05, *n* = 4), and inhibition at 0.75 μmol/L (0.056 ± 0.007 *pHi* units/minute) was similar (*P*>0.05, *n* = 4) to 0.25 μmol/L STa (see also Fig 2B).

**P*<0.04 versus Control

†*P*<0.03 versus STa or HOE-694

‡*P*<0.03 versus Forskolin, Forskolin + STa, and H89 + HOE-694 + STa,

§*P*<0.05 versus db-cGMP

¶*P*<0.05 versus db-cGMP + STa and db-cGMP + HOE-694, &*P*<0.05 versus db-cGMP

§*P*<0.03 versus SNP + STa and SNP + HOE-694

doi:10.1371/journal.pone.0146042.t001

similar (*P*>0.05) to that in cells in the presence of STa. Coincubation of cells with STa + HOE-694 resulted in a decrease in *J_H⁺* (89 ± 6%) that was higher compared with the effect seen in cells treated with STa or HOE-694 alone.

NHE1, NHE2 and NHE4 protein abundance

To address whether STa-associated decrease in *J_H⁺* was due to lower protein abundance of NHE4, or whether this toxin alters NHE1 or NHE2 protein abundance, the protein level of these membrane transporters was assayed. The results show that incubation of T₈₄ cells with STa did not alter NHE1, NHE2 or NHE4 protein abundance (Fig 3).

cGMP and cAMP involvement on NHE4–mediated *pHi* recovery kinetics

STa is shown to increase the cGMP level in T₈₄ cells [34]; however, the role of cGMP as modulator of NHE4 activity is not addressed [17]. Thus, we next investigated whether STa effect on




Fig 1. Effect of STa on pH_i recovery. T₈₄ cells were preloaded with BCECF-AM in the absence or presence (30 minutes) of 0.25 μmol/L heat-stable (STa) enterotoxin. After transferring the cells into a spectrofluorometer the basal pH_i was stabilized and then exposed (1.5–2 minutes) to a control solution containing 20 mmol/L NH₄Cl (+ NH₄Cl). Cells were then rinsed with a NH₄Cl-free solution (–NH₄Cl) and left in this medium without (Control) or with 0.25 μmol/L STa, 25 μmol/L HOE-694, or both (STa + HOE-694) (see Methods). Initial rates of pH_i recovery were calculated from data collected for the first 60 seconds after removing the NH₄Cl load. Values are mean ± S.E.M. (n = 25–27).

doi:10.1371/journal.pone.0146042.g001

NHE4-dependent dpH_i/dt in this cell type was modulated by direct administration of exogenous cGMP. The results show that dpH_i/dt and basal pH_i (Table 1), and J_{H^+} (Fig 4) were unaltered in T₈₄ cells exposed to db-cGMP in the absence of HOE-694 or STa. However, the reduction in dpH_i/dt and J_{H^+} seen in response to STa, HOE-694, or STa + HOE-694 was unaltered by db-cGMP. When cells were incubated with SNP (a spontaneous NO donor) [27] the results were similar to those in the presence of db-cGMP (Table 1, Fig 4). Parallel results show that cGMP intracellular level was increased by STa and SNP, confirming previous reports in T₈₄ cells [35] and rat distal colon crypts [36], but it was unaltered by HOE-694 (not shown).

We next assayed whether cAMP was involved in the response of T₈₄ cells to STa-reduced NHE4-mediated pH_i recovery kinetics. Cells incubated with forskolin (adenylyl cyclase activator) [29] in the absence of HOE-694 resulted in a decrease in dpH_i/dt (Table 1) and J_{H^+} (Fig 5A) that was of a similar magnitude to the decrease seen in cells incubated with STa in the absence or presence of this activator. However, in the presence of HOE-694 or STa + HOE-694, forskolin caused a reduction in these parameters that was similar to that seen in cells coincubated with STa + HOE-694 in the absence of this activator. Parallel results show that intracellular level of cAMP increased by STa (4.9 ± 0.5 fold) and forskolin (8.9 ± 1.5 fold) (Fig 5B). Additionally, pre-incubation of cells with H89 (inhibitor of PKA) [28] reversed the decrease in dpH_i/dt and J_{H^+} caused by STa + HOE-694 to values that are comparable to STa or HOE-694 alone.

Discussion

This study shows that the enterotoxigenic *Escherichia coli* (ETEC) released heat-stable (STa) enterotoxin decreases the pH_i recovery kinetics in the human col carcinoma T₈₄ cell line. This




Fig 2. Effect of STa on J_{H^+} . The overall transmembrane H^+ flux rates (J_{H^+}) were calculated from initial rates of pH_i recovery and the intrinsic buffer capacity (β_i) values (see [Methods](#)). **A**, T_{84} cells were exposed to culture medium without (–, Control) or with (+) 0.25 $\mu\text{mol/L}$ heat-stable (STa) enterotoxin, 25 $\mu\text{mol/L}$ HOE-694, or both (see [Methods](#)). **B**, T_{84} cells were exposed to increasing concentrations of STa in the presence of 25 $\mu\text{mol/L}$ HOE-694 as in A. In A, * $P < 0.05$ versus Control, † $P < 0.05$ versus STa or HOE-694. In B, * $P < 0.05$ versus Control, † $P < 0.05$ versus other values in STa + HOE-694. Values are mean \pm S.E.M. ($n = 25\text{--}27$).

doi:10.1371/journal.pone.0146042.g002

phenomenon results from a lower activity of NHE4 without altering its protein expression. STa effect depends on the level of cAMP, but not cGMP, and PKA activation. These findings represent a novel mechanism of pH_i homeostasis by STa that could have consequences in the physiology of gastrointestinal cells leading to human diarrhoea.

STa modulation of NHEs activity

STa is an enterotoxin that causes gastrointestinal electrolyte imbalance characterized by a higher Cl^- release to the gastrointestinal lumen, a phenomenon that ends in diarrhoea in humans [1,3–5]. One of the potential mechanisms for these adverse effects of STa is a mucosal alkalization due to lower activity of plasma membrane mechanisms involved in maintaining transmembrane distribution of H^+ , including NHEs activity [10,11]. Our results show that STa caused a decrease




Fig 3. Effect of STa on NHE4 protein abundance. Western blot for NHE4 protein abundance in whole extracts of T₈₄ cells exposed for 30 minutes in the absence (Control) or presence (STa) of 0.25 μmol/L heat-stable (STa) enterotoxin. Lower panel: NHE4/β-actin ratio densitometries normalized to 1 in Control. β-Actin is internal reference. Values are mean ± S.E.M. (n = 15).

doi:10.1371/journal.pone.0146042.g003

in NHEs activity resulting in lower H⁺ efflux (i.e., J_{H⁺}). This phenomenon may be responsible for the observed reduction in the capacity to restore the p_{H_i} recovery kinetics after an acid pulse. This possibility is supported by the findings showing that STa caused a similar reduction in dp_{H_i}/dt and J_{H⁺} (reduction in dp_{H_i}/dt / reduction in J_{H⁺} = 1.1), thus, making possible that alterations in the p_{H_i} recovery rate caused by STa was due to reduced H⁺ efflux kinetics. In addition, since the intrinsic buffering capacity (β_i) values were unaltered by STa (β_i with STa/β_i without STa = 1), it is unlikely that these alterations were the result of an altered β_i in T₈₄ cells. Indeed, in cells incubated with STa the p_{H_i} value was not significantly altered (p_{H_i} with STa/p_{H_i} without STa = 0.996) compared with cells in the absence of this toxin.




Fig 4. Involvement of cGMP on STa modulation of J_{H⁺}. T₈₄ cells were exposed for 30 minutes in the absence (-SNP) or presence (+ SNP) of 500 μmol/L sodium nitroprusside (SNP). The overall transmembrane H⁺ flux rates (J_{H⁺}) were calculated from initial rates of p_{H_i} recovery and the intrinsic buffer capacity (β_i) values (see Methods). Cells were exposed to culture medium without (-, Control, red bar) or with (+) 100 μmol/L dibutyl cyclic GMP (db-cGMP), 0.25 μmol/L STa, and/or 25 μmol/L HOE-694 (see Methods). *P<0.05 versus Control or corresponding db-cGMP, †P<0.05 versus corresponding STa or HOE-694 in the presence of db-cGMP. Values are mean ± S.E.M. (n = 25–27).

doi:10.1371/journal.pone.0146042.g004




Fig 5. Involvement of cAMP and PKA on STa modulation of J_{H^+} . A, The overall transmembrane H^+ flux rates (J_{H^+}) were calculated from initial rates of pH_i recovery and the intrinsic buffer capacity (β_i) values (see [Methods](#)). Cells were exposed to culture medium without (-, Control, red bar) or with (+) 10 μ mol/L forskolin, 0.25 μ mol/L heat-stable (STa) enterotoxin, 25 μ mol/L HOE-694, and/or 100 nmol/L H89 (see [Methods](#)). B, cAMP levels in cells in the absence (Control) or presence of STa or forskolin as in A. In A, * $P < 0.05$ versus Control, † $P < 0.05$ versus STa, HOE-694, or STa + HOE-694. In B, * $P < 0.05$ versus Control, † $P < 0.05$ versus STa. Values are mean \pm S.E.M. ($n = 25-27$).

doi:10.1371/journal.pone.0146042.g005

Interestingly, it was initially shown [31] that T_{84} cells express mainly NHEs (NHE1, NHE2 and NHE4) [21], in a minor grade Cl^-/HCO_3^- exchangers and Na^+/HCO_3^- cotransporters, but not other classical mechanisms of H^+ export such as the vacuolar H^+ -ATPases [37] or H^+/K^+ -ATPases [38]. Out of these membrane transport systems, NHEs play a major role in the removal of intracellular H^+ in most cell types maintaining stable pH_i and extracellular pH values [14–17,20,37,38].

NHE4 is an isoform of the NHEs family of membrane exchangers whose function results in the modulation of pH_i in mammalian cells [14,16,17]. This membrane Na^+/H^+ exchanger isoform is expressed in the human gastrointestinal tract, and is co-expressed with NHE1 and

NHE2, but not NHE3, in T₈₄ cells [21,33], as confirmed in this study. Interestingly, cells exposed to HOE-694 show lower dpH_i/dt and J_{H^+} most likely via a mechanism involving lower activity of NHE1 and NHE2 isoforms, since the concentration of this inhibitor used in the present study (25 $\mu\text{mol/L}$) preferentially inhibits these isoforms, but not NHE4 [21,26]. Indeed, cells in the presence of HOE-694 show partial recovery of the pH_i value suggesting that not all the pH_i recovery is mediated by NHE1 and NHE2, but other mechanism(s) is plausible in this cell type.

Since STa in the presence of HOE-694, i.e., where NHE1 and NHE2 were not functional, almost abolished the dpH_i/dt and J_{H^+} (both reduced by ~90%), it is likely that NHE4 isoform was inhibited by this enterotoxin in T₈₄ cells. This possibility is supported when we consider that the concentration of STa used in our study is close to the STa half-maximal stimulatory concentration for cGMP accumulation reported in T₈₄ cells [25]. Additionally, the possibility that STa reduces the dpH_i/dt and J_{H^+} via a mechanism including lower expression of NHE4, or NHE1 or NHE2, is unlikely since the protein abundance for none of these isoforms were altered by the toxin. Thus, STa-reduced H^+ efflux seems to be due to a lower activity rather than expression of NHE4 in this cell type. STa effect in the presence of HOE-694 leads a remaining fraction of pH_i recovery that accounted for 10% of the total recovery after an acid pulse. This finding could result from other mechanisms than inhibition of NHE1, 2 or 4, such as activity of Cl^-/HCO_3^- exchangers and/or Na^+/HCO_3^- cotransporters expressed in T₈₄ cells [31]. Indeed, STa was shown to increase HCO_3^- secretion via a higher Na^+/HCO_3^- activity in duodenal CFRT^{-/-} mice [39]. However, our pH_i recovery assays were performed in the absence of extracellular HCO_3^- in this cell type making the latter unlikely.

Involvement of cAMP on STa effect

It has been shown that STa increases Cl^- secretion in a cAMP- and cGMP-dependent manner via CFTR channels in rat jejunum [9]. Initial reports show that STa-increased cGMP, but unaltered cAMP level in rabbit distal ileum mucosa [40] or reduced cAMP level in mice intestine [41]. Our results show that exposure of T₈₄ cells to STa results in increased cGMP and cAMP levels. Since these nucleotides decrease NHEs activity [12,13], STa-increased levels may have functional consequences on pH_i recovery in T₈₄ cells.

Since incubation of cells with exogenous cGMP (db-cGMP) did not alter basal dpH_i/dt and J_{H^+} in our assays it is likely that this cyclic nucleotide is not involved in the modulation of NHEs activity in T₈₄ cells. Furthermore, the inhibitory effect of STa on dpH_i/dt and J_{H^+} in the presence of HOE-694 was unaltered by db-cGMP, suggesting that NHE4 inhibition by STa was independent of cGMP. This is supported by the findings showing that dpH_i/dt and J_{H^+} inhibition by STa or HOE-694 alone was unaltered when cells were incubated with these molecules and db-cGMP. Additionally, exposure of cells to exogenous NO delivered by SNP, a spontaneous NO donor [27], does not change STa effect in the absence or presence of HOE-694. Since SNP did not alter the reduction in the dpH_i/dt and J_{H^+} caused by HOE-694 itself, NO in this cell type may not alter this inhibitors' effectiveness on NHE1 and NHE2.

It was early shown that forskolin, a potent activator of adenylyl cyclase, has a profound effect in T₈₄ transmonolayer net water flux (J_w) [29], suggesting that cAMP could be involved in this phenomenon. Unfortunately, the cAMP level was not determined in the latter study. Additionally, incubation of T₈₄ cells with secretagogues whose actions are mediated by cAMP ends with Cl^- secretion from this cell type [35,42–44]. However, it is paradoxical that even when the level of cAMP was found unaltered in T₈₄ cells in response to STa, this toxin effect on Cl^- secretion closely resembles a cAMP-mediated mechanism in this cell type [35]. Our findings show that cAMP level is increased in T₈₄ cells treated with STa or with forskolin. Since the




Fig 6. Potential involvement of cAMP and PKA on STa modulation of J_{H^+} . In T₈₄ cells the enterotoxigenic *Escherichia coli* (ETEC) released heat-stable enterotoxin (STa) activates guanylyl cyclase-C (GC-C) receptors to generate (green arrow) cyclic GMP (cGMP) increasing (↑) its intracellular level. STa also increases cyclic AMP (cAMP) level via a mechanism that is not well defined in this cell type (?). Increase in cAMP activates protein kinase A (PKA), which could be responsible of a reduced (↓) activity of the Na⁺/H⁺ exchanger isoform 4 (NHE4). The resulting intracellular accumulation of H⁺ leads to intracellular acidification, a phenomenon that, via undefined mechanism, could be responsible for the increase in chloride (Cl⁻) secretion via the cystic fibrosis transmembrane conductance regulator channels (CFRT) reported in this cell type and human diarrhoea.

doi:10.1371/journal.pone.0146042.g006

effect of forskolin alone was to diminish the dpH_i/dt and J_{H^+} in a same magnitude as STa alone or STa + forskolin, it is likely that a higher cAMP level could be involved in downregulation of NHE4 activity in this cell type. Parallel results suggest that NHE1 and NHE2 may not be under modulation by STa–or forskolin–mediated cAMP increase since the inhibition caused by HOE-694 of dpH_i/dt and J_{H^+} by itself or in the presence of STa was unaltered by forskolin. Interestingly, since H89, a PKA inhibitor, resulted in restoration of the reduced dpH_i/dt and J_{H^+} seen in the presence of STa + HOE-694 + forskolin to values that are comparable to those in the presence of these molecules per separate, it is likely that PKA may mediate STa inhibition of NHE4 in T₈₄ cells.

In conclusion, the enterotoxigenic *Escherichia coli* released STa enterotoxin has a deleterious effect on the normal physiology of T₈₄ cells *in vitro*. In terms of its association with human diarrhoea this enterotoxin was found to increase not only cGMP levels, but also the cAMP level, perhaps leading to PKA activation in this cell type. It is proposed that STa reduces the capacity of T₈₄ cells to recover the pH_i after an acid pulse via a mechanism that includes reduced activity of NHE4, but not NHE1 or NHE2, in this cell type. These findings constitute a novel mechanism of pH_i homeostasis by STa in this cell type, and perhaps in the

gastrointestinal epithelium, resulting in a deficient recovery rate and H⁺ efflux after metabolic alterations associated with intracellular acidification. These findings complement the reduced transepithelial electrical resistance caused by STa in T₈₄ cells, indicative of an intestinal barrier dysfunction in addition to STa-induced water secretion [45]. Considering that T₈₄ cells respond with increased Cl⁻ release to STa via cGMP- and cAMP-dependent mechanisms, a role of NHE4 in this phenomenon is proposed. All together the alterations caused by STa in a functional sequence (i.e., STa / increased cAMP / increased PKA activity / decreased NHE4 activity / increased intracellular acidification) (Fig 6) could have consequences in the physiology of gastrointestinal cells promoting human diarrhoea.

Acknowledgments

Authors thank research staff at the Cellular Physiology Laboratory of the Biomedical Department, Faculty of Health Sciences, Universidad de Antofagasta, and from the Cellular and Molecular Physiology Laboratory (CMPL) from Pontificia Universidad Católica de Chile.

Author Contributions

Conceived and designed the experiments: ARB GM LS MAR. Performed the experiments: ARB LRC-L CNAB MC JA FP AL KN. Analyzed the data: ARB FT JA FP AL CS GM LS MAR KN. Contributed reagents/materials/analysis tools: CS FP AL FT GM LS MAR. Wrote the paper: ARB MC LS MAR.

References

1. Qadri F, Svennerholm AM, Faruque AS, Sack RB. Enterotoxigenic *Escherichia coli* in developing countries: epidemiology, microbiology, clinical features, treatment, and prevention. *Clin Microbiol Rev*. 2005; 18: 465–483. PMID: [16020685](#)
2. Gross R. *Escherichia coli* diarrhoea. *J Infect*. 1983; 7: 177–192. PMID: [6363562](#)
3. Cravioto A, Reyes R, Ortega R, Fernandez G, Hernandez R, Lopez D. Prospective study of diarrhoeal disease in a cohort of rural Mexican children: incidence and isolated pathogens during the first two years of life. *Epidemiol Infect*. 1988; 101: 123–134. PMID: [3402544](#)
4. Abu-Elyazeed R, Wierzbka A, Mourad L, Peruski B, Kay BA, Rao M, et al. Epidemiology of enterotoxigenic *Escherichia coli* diarrhea in a pediatric cohort in a periurban area of lower Egypt. *J Infect Dis*. 1999; 179: 382–389. PMID: [9878022](#)
5. Moon C, Zhang W, Sundaram N, Yarlagadda S, Reddy VS, Arora K, et al. Drug-induced secretory diarrhea: A role for CFTR. *Pharmacol Res*. 2015; 102: 107–112. doi: [10.1016/j.phrs.2015.08.024](#) PMID: [26429773](#)
6. Forte LR, Krause WJ, Freeman RH. *Escherichia coli* enterotoxin receptors: localization in opossum kidney, intestine, and testis. *Am J Physiol*. 1989; 257: F874–F881. PMID: [2556042](#)
7. Schulz S, Green CK, Yuen PS, Garbers DL. Guanylyl cyclase is a heat-stable enterotoxin receptor. *Cell* 1990; 63: 941–948. PMID: [1701694](#)
8. Carrithers SL. Diarrhea or colorectal cancer: can bacterial toxins serve as a treatment for colon cancer?. *Proc Natl Acad Sci USA*. 2003; 100: 3018–3020. PMID: [12631696](#)
9. Golin-Bisello F, Bradbury N, Ameen N. STa and cGMP stimulate CFTR translocation to the surface of villus enterocytes in rat jejunum and is regulated by protein kinase G. *Am J Physiol*. 2005; 289: C708–C716.
10. Fawcus K, Gorton VJ, Lucas ML, McEwan GT. Stimulation of three distinct guanylate cyclases induces mucosal surface alkalisation in rat small intestine in vitro. *Comp Biochem Physiol A Physiol*. 1997; 118: 291–295. PMID: [9366058](#)
11. Lucas ML. A reconsideration of the evidence for *Escherichia coli* STa (heat stable) enterotoxin-driven fluid secretion: a new view of STa action and a new paradigm for fluid absorption. *J Applied Microbiol*. 2001; 90: 7–26.
12. Bachmann O, Juric M, Seidler U, Manns M, Yu H. Basolateral ion transporters involved in colonic epithelial electrolyte absorption, anion secretion and cellular homeostasis. *Acta Physiol*. 2011; 201: 33–46.

13. Arena E, Longo W, Roberts K, Geibel P, Nateqi J, Brandstetter M, et al. Functional role of NHE4 as a pH regulator in rat and human colonic crypts. *Am J Physiol*. 2012; 302: C412–C418.
14. Slepov ER, Rainey JK, Sykes BD, Fliegel L. Structural and functional analysis of the Na⁺/H⁺ exchanger. *Biochem J*. 2007; 401: 623–633. PMID: [17209804](#)
15. Orłowski J, Grinstein S. Na⁺/H⁺ exchangers. *Compr Physiol*. 2011; 1: 2083–2100. doi: [10.1002/cphy.c110020](#) PMID: [23733698](#)
16. Provost JJ, Wallert MA. Inside out: targeting NHE1 as an intracellular and extracellular regulator of cancer progression. *Chem Biol Drug Des*. 2013; 81: 85–101. doi: [10.1111/cbdd.12035](#) PMID: [23253131](#)
17. Fuster DG, Alexander RT. Traditional and emerging roles for the SLC9 Na⁺/H⁺ exchangers. *Pflugers Arch*. 2014; 466: 61–76. doi: [10.1007/s00424-013-1408-8](#) PMID: [24337822](#)
18. Pizzonia JH, Biemesderfer D, Abu-Alfa AK, Wu MS, Exner M, Isenring P, et al. Immunohistochemical characterization of Na⁺/H⁺ exchanger isoform NHE4. *Am J Physiol*. 2013; 275: F510–F517.
19. Gawenis LR, Greeb JM, Prasad V, Grisham C, Sanford LP, Doetschman T, et al. Impaired gastric acid secretion in mice with a targeted disruption of the NHE4 Na⁺/H⁺ exchanger. *J Biol Chem*. 2005; 280: 12781–12789. PMID: [15684419](#)
20. Rossmann H, Sonnentag T, Heinzmann A, Seidler B, Bachmann O, Vieillard-Baron D, et al. Differential expression and regulation of Na⁺/H⁺ exchanger isoforms in rabbit parietal and mucous cells. *Am J Physiol*. 2001; 281: G447–G458.
21. Beltran AR, Ramirez MA, Carraro-Lacroix LR, Hiraki Y, Reboucas NA, Malnic G. NHE1, NHE2, and NHE4 contribute to regulation of cell pH in T84 colon cancer cells. *Pflugers Arch*. 2008; 455: 799–810. PMID: [17943310](#)
22. Selvaraj NG, Prasad R, Goldstein JL, Rao MC. Evidence for the presence of cGMP-dependent protein kinase-II in human distal colon and in T84, the colonic cell line. *Biochim Biophys Acta* 2000; 1484: 32–43.
23. Fernandez R, Malnic G. H⁺ ATPase and Cl⁻ interaction in regulation of MDCK cell. *J Membr Biol*. 1998; 163: 137–145. PMID: [9592078](#)
24. Aravena C, Beltran AR, Cornejo M, Torres V, Diaz ES, Guzmán-Gutiérrez, et al. Potential role of sodium-proton exchangers in the low concentration arsenic trioxide-increased intracellular pH and cell proliferation. *Plos One* 2012; 7: e51451. doi: [10.1371/journal.pone.0051451](#) PMID: [23236503](#)
25. Bijvelds MJ, Loos M, Bronsveld I, Hellemaans A, Bongartz JP, Ver Donck L, et al. Inhibition of heat-stable toxin-induced intestinal salt and water secretion by a novel class of guanylyl cyclase C inhibitors. *J Infect Dis*. 2015; 212: 1806–1815. doi: [10.1093/infdis/jiv300](#) PMID: [25999056](#)
26. Ikuma M, Kashgarian M, Binder HJ, Rajendran VM. Differential regulation of NHE isoforms by sodium depletion in proximal and distal segments of rat colon. *Am J Physiol*. 1999; 276: G539–G549. PMID: [9950829](#)
27. Pardo F, Silva L, Sáez T, Salsoso R, Gutiérrez J, Sanhuesa C, et al. Human supraphysiological gestational weight gain and fetoplacental vascular dysfunction. *Int J Obes (Lond)*. 2015; 39: 1264–1273.
28. Ao M, Sarathy J, Domingue J, Alrefai WA, Rao MC. Chenodeoxycholic acid stimulates Cl⁻ secretion via cAMP signaling and increases cystic fibrosis transmembrane conductance regulator phosphorylation in T84 cells. *Am J Physiol*. 2013; 305: C447–C456.
29. Toriano R, Kierbel A, Ramírez MA, Malnic G, Parisi M. Spontaneous water secretion in T84 cells: effects of STa enterotoxin, bumetanide, VIP, forskolin, and A-23187. *Am J Physiol*. 2001; 281: G816–G822.
30. Salsoso R, Guzmán-Gutiérrez E, Sáez T, Bugueño K, Ramírez MA, Fariás M, et al. Insulin restores L-arginine transport requiring adenosine receptors activation in umbilical vein endothelium from late-onset preeclampsia. *Placenta* 2015; 36: 287–296. doi: [10.1016/j.placenta.2014.12.007](#) PMID: [25573092](#)
31. Ramírez MA, Toriano R, Parisi M, Malnic G. Control of cell pH in the T84 colon cell line. *J Membr Biol*. 2000; 177: 149–157. PMID: [11003689](#)
32. Musa-Aziz R, Oliveira-Souza M, Mello-Aires M. Signaling pathways in the biphasic effect of ANG II on Na⁺/H⁺ exchanger in T84 cells. *J Membr Biol*. 2005; 205: 49–60. PMID: [16283585](#)
33. Toriano R, Ozu M, Politi MT, Dorr RA, Curto MA, Capurro C. Uroguanylin regulates net fluid secretion via the NHE2 isoform of the Na⁺/H⁺ exchanger in an intestinal cellular model. *Cell Physiol Biochem*. 2011; 28: 733–742. doi: [10.1159/000335767](#) PMID: [22178885](#)
34. Kots AY, Choi BK, Estrella-Jimenez ME, Warren CA, Gilbertson SR, Guerrant RL, et al. Pyridopyrimidine derivatives as inhibitors of cyclic nucleotide synthesis: Application for treatment of diarrhea. *Proc Natl Acad Sci USA*. 2008; 105: 8440–8445. doi: [10.1073/pnas.0803096105](#) PMID: [18559851](#)

35. Huott PA, Liu W, McRoberts JA, Giannella RA, Dharmasathaphorn K. Mechanism of action of Escherichia coli heat stable enterotoxin in a human colonic cell line. *J Clin Invest.* 1988; 82: 514–523. PMID: [2457034](#)
36. Morel E, Dublineau I, Griffiths NM. Effect of radiation on cAMP, cGMP and Ca²⁺ pathways and their interactions in rat distal colon. *Radiat Res.* 2003; 160: 263–272. PMID: [12926985](#)
37. Casey JR, Grinstein S, Orlowski J. Sensors and regulators of intracellular pH. *Nat Rev Mol Cell Biol.* 2010; 11: 50–61. doi: [10.1038/nrm2820](#) PMID: [19997129](#)
38. Gillies RJ, Raghunand N, Garcia-Martin ML, Gatenby RA. pH imaging. A review of pH measurement methods and applications in cancers. *IEEE Eng Med Biol Mag.* 2004; 23: 57–64.
39. Sellers ZM, Childs D, Chow JYC, Smith AJ, Hogan DL, Isenberg JI, et al. Heat-stable enterotoxin of Escherichia coli stimulate a non-CFTR-mediated duodenal bicarbonate secretory pathway. *Am J Physiol.* 2005; 288: G654–G663.
40. Field M, Graf LH, Laird WJ, Smith PL. Heat-stable enterotoxin of Escherichia coli: in vitro effects on guanylate cyclase activity, cyclic GMP concentration, and ion transport in small intestine. *Proc Natl Acad Sci USA.* 1978; 75: 2800–2804. PMID: [26915](#)
41. Giannella RA, Drake KW. Effect of purified Escherichia coli heat-stable enterotoxin on intestinal cyclic nucleotide metabolism and fluid secretion. *Infect Immun.* 1979; 24: 19–23. PMID: [378842](#)
42. Cartwright CA, McRoberts JA, Mandel KG, Dharmasathaphorn K. Synergistic action of cyclic adenosine monophosphate-and calcium-mediated chloride secretion in a colonic epithelial cell line. *J Clin Invest.* 1985; 76: 1837–1842. PMID: [2997291](#)
43. Wasserman SI, Barrett KE, Huott PA, Beuerlein G, Kagnoff MF, et al. Immune-related intestinal Cl⁻ secretion. I. Effect of histamine on the T84 cell line. *Am J Physiol.* 1988; 254: C53–C62. PMID: [3337221](#)
44. Nichols JM, Maiellaro I, Abi-Jaoude J, Curci S, Hofer AM. “Store-operated” cAMP signaling contributes to Ca²⁺-activated Cl⁻ secretion in T84 colonic cells. *Am J Physiol.* 2015; 309: G670–G679.
45. Nakashima R, Kamata Y, Nishikawa Y. Effects of Escherichia coli heat-stable enterotoxin and guanylin on the barrier integrity of intestinal epithelial T84 cells. *Vet Immunol Immunopathol.* 2013; 152: 78–81. doi: [10.1016/j.vetimm.2012.09.026](#) PMID: [23078906](#)