

CORRECTION

Correction: Association between Body Mass Index and Prognosis of Colorectal Cancer: A Meta-Analysis of Prospective Cohort Studies

The *PLOS ONE* Staff

[Fig 2](#), “Relative Risks for the Association between Pre-diagnosis BMI and Colorectal Cancer-Specific and All-cause Mortality,” has been truncated. Please view [Fig 2](#) here. The publisher apologizes for the error.

[Fig 3](#), “Relative Risks for the Association between Post-diagnosis BMI and Colorectal Cancer-specific and All-cause Mortality,” has been truncated. Please view [Fig 3](#) here. The publisher apologizes for the error.




OPEN ACCESS

Citation: The *PLOS ONE* Staff (2016) Correction: Association between Body Mass Index and Prognosis of Colorectal Cancer: A Meta-Analysis of Prospective Cohort Studies. *PLoS ONE* 11(1): e0147456. doi:10.1371/journal.pone.0147456

Published: January 14, 2016


Copyright: © 2016 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

CRC-specific mortality: Underweight vs. Normal weight




Test for heterogeneity:
(Q = 2.64, P = 0.27, $I^2 = 24.30\%$)

All-cause Mortality: Underweight vs. Normal weight




Test for heterogeneity:
(Q = 0.58, P = 0.75, $I^2 = 0.00\%$)

CRC-specific mortality: Overweight vs. Normal weight




Test for heterogeneity:
(Q = 6.84, P = 0.34, $I^2 = 12.32\%$)

All-cause mortality: Overweight vs. Normal weight




Test for heterogeneity:
(Q = 5.07, P = 0.41, $I^2 = 1.39\%$)

CRC-specific mortality: Obesity vs. Normal weight



Test for heterogeneity:
(Q = 2.68, P = 0.00, $I^2 = 0.00\%$)

All-cause mortality: Obesity vs. Normal weight




Test for heterogeneity:
(Q = 4.86, P = 0.30, $I^2 = 17.77\%$)

Fig 2. Relative Risks for the Association between Pre-diagnosis BMI and Colorectal Cancer-Specific and All-cause Mortality. Association between pre-diagnosis BMI and colorectal cancer-specific mortality and all-cause mortality.


doi:10.1371/journal.pone.0147456.g001

CRC-specific mortality: Overweight vs. Normal weight




Test for heterogeneity:
 $(Q = 5.09, P = 0.17, I^2 = 41.06\%)$

All-cause mortality: Overweight vs. Normal weight




CRC-specific mortality: Obesity vs. Normal weight




Test for heterogeneity:
 $(Q = 5.14, P = 0.16, I^2 = 41.68\%)$

All-cause mortality: Obesity vs. Normal weight



All-cause Mortality: Underweight vs. Normal weight



Test for heterogeneity:
 $(Q = 15.42, P = 0.08, I^2 = 41.62\%)$

All-cause mortality: Obesity (BMI ≥ 35) vs. Normal weight




Fig 3. Relative Risks for the Association between Post-diagnosis BMI and Colorectal Cancer-specific and All-cause Mortality. Association between post-diagnosis BMI and colorectal cancer-specific mortality and all-cause mortality.

doi:10.1371/journal.pone.0147456.g002

Reference

1. Lee J, Meyerhardt JA, Giovannucci E, Jeon JY (2015) Association between Body Mass Index and Prognosis of Colorectal Cancer: A Meta-Analysis of Prospective Cohort Studies. PLoS ONE 10(3): e0120706. doi: [10.1371/journal.pone.0120706](https://doi.org/10.1371/journal.pone.0120706) PMID: [25811460](https://pubmed.ncbi.nlm.nih.gov/25811460/)