

RESEARCH ARTICLE

# Biochar, Bentonite and Zeolite Supplemented Feeding of Layer Chickens Alters Intestinal Microbiota and Reduces *Campylobacter* Load

Tanka P. Prasai<sup>1</sup>, Kerry B. Walsh<sup>1</sup>, Surya P. Bhattarai<sup>1</sup>, David J. Midmore<sup>1</sup>, Thi T. H. Van<sup>2</sup>, Robert J. Moore<sup>2,3</sup>, Dragana Stanley<sup>1\*</sup>

**1** Central Queensland University, Institute for Future Farming Systems, Rockhampton, Queensland, Australia, **2** RMIT University, School of Applied Sciences and Health Innovations Research Institute (HIRI), Bundoora, Victoria, Australia, **3** Department of Microbiology, Monash University, Clayton, Victoria, Australia

\* [D.Stanley@cqu.edu.au](mailto:D.Stanley@cqu.edu.au)



## Abstract

A range of feed supplements, including antibiotics, have been commonly used in poultry production to improve health and productivity. Alternative methods are needed to suppress pathogen loads and maintain productivity. As an alternative to antibiotics use, we investigated the ability of biochar, bentonite and zeolite as separate 4% feed additives, to selectively remove pathogens without reducing microbial richness and diversity in the gut. Neither biochar, bentonite nor zeolite made any significant alterations to the overall richness and diversity of intestinal bacterial community. However, reduction of some bacterial species, including some potential pathogens was detected. The microbiota of bentonite fed animals were lacking all members of the order Campylobacterales. Specifically, the following operational taxonomic units (OTUs) were absent: an OTU 100% identical to *Campylobacter jejuni*; an OTU 99% identical to *Helicobacter pullorum*; multiple *Gallibacterium anatis* (>97%) related OTUs; *Bacteroides dorei* (99%) and *Clostridium aldenense* (95%) related OTUs. Biochar and zeolite treatments had similar but milder effects compared to bentonite. Zeolite amended feed was also associated with significant reduction in the phylum Proteobacteria. All three additives showed potential for the control of major poultry zoonotic pathogens.

## OPEN ACCESS

**Citation:** Prasai TP, Walsh KB, Bhattarai SP, Midmore DJ, Van TTH, Moore RJ, et al. (2016) Biochar, Bentonite and Zeolite Supplemented Feeding of Layer Chickens Alters Intestinal Microbiota and Reduces *Campylobacter* Load. PLoS ONE 11(4): e0154061. doi:10.1371/journal.pone.0154061

**Editor:** Gunnar Loh, Max Rubner-Institut, GERMANY

**Received:** January 28, 2016

**Accepted:** April 7, 2016

**Published:** April 26, 2016

**Copyright:** © 2016 Prasai et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** The data are on the MG-RAST server under project ID 4693702.3.

**Funding:** This work was university funded.

**Competing Interests:** The authors have declared that no competing interests exist.

## Introduction

The use of antibiotic growth promoters as feed additives to suppress the pathogenic bacteria in the gut has been common in commercial poultry production, however it is banned in Europe [1] because of concerns for the consequences it could have on human health in terms of the selection of antibiotic resistant microbiota and for the presence of residual antibiotics in poultry products [2]. Alternatives to antibiotic growth promoters are required in order to maintain bird health and deliver the productivity improvements that were sometimes associated with their use.

Laying hens are in great need of antibiotic-free pathogen control given antibiotics can not be used due to residue carry over to eggs. For example, in Queensland, Spotty Liver is emerging as a disease of concern. This disease is caused by *Campylobacter* species [3] and is currently controlled by antibiotics. Layers colonisation with human pathogens such as *Salmonella* and *Campylobacter* is an important issue that the industry is grappling with, and for which new solutions are required. Additionally antibiotic-free pathogen control is needed in organic poultry production. There are many alternative products under investigation. Among them, biochar, bentonite and zeolite are interesting candidates for selective pathogen control as there is mounting evidence that they are safe and beneficial products [4–6]. Properties of these three natural products are outlined below.

Biochar is a carbon rich product produced from the incomplete combustion of biomass in the absence of oxygen through a process termed pyrolysis [7]. Biochar, and in particular biochar bokashi, is used as a feed supplement in Japan and China, with claims for improved digestion and feed conversion ratio [4]. More complete biochar pyrolysis results in production of charcoal. Diet supplementation with charcoal has also been reported to result in increased live weight gain and higher feed conversion ratio (FCR) for commercial meat chickens and ducks [8, 9]. Several mechanisms have been suggested for the benefit of biochar or charcoal in animal diets, including toxin binding, improved digestion and retention of nitrogen. A possible mechanism for the improved FCR associated with biochar supplements could lie with a change in gastro intestinal tract (GIT) microbiota.

Bentonite is a clay mineral with strong colloidal properties and the ability to rapidly absorb many times its volume of water. Clays are often incorporated in animal diets as a stabilizer, lubricant or agglomerant to improve feed manufacture [10]. Nutrient digestibility and enzymatic activity of gastrointestinal secretions has been improved by addition of clay to broilers and pig feedstuffs [11–14]. Bentonite has been used effectively as a feed additive in poultry rations, with the swelling of bentonite causing a reduction in the rate of feed transit through the digestive tract, permitting time for more effective utilisation [15]. Addition of sodium bentonite was effective in ameliorating the negative effect of aflatoxins in poultry diet [16]. The toxin is prevented from being absorbed by the digestive tract and the bound aflatoxin is then excreted [6]. The supplementation of poultry rations with a Cu-montmorillonite clay has been reported to result in reduced total viable counts of *Escherichia coli* and *Clostridium* in the small intestine and caecum of chicks [17].

Clinoptilolite is a common form of natural zeolite. Zeolites are crystalline, hydrated aluminosilicates of alkali and alkaline earth cations. Zeolites have cation exchange properties and are capable of trapping molecules within their pores [18]. For example, the porosity, particle and crystal size of the zeolitic material and its degree of aggregation determine the rate of access of ingesta fluids during passage through the GIT [5]. Average daily live body weight gain and feed conversions in broilers have been improved with dietary inclusion of zeolites [5, 19]. Zeolite feed amendment has also been reported to increase egg production [20] and have positive effects on egg weight and internal egg quality [5, 20]. Papaioannou *et al.* [21] reported zeolite feed amendment to be associated with a reduction in the rate of passage of feed through the digestive system, and an associated reduction in feed intake [22] resulting in better FCR. However, factors including the type of zeolite, its purity, physiochemical properties, and the supplementation level used in the diets may impact the performance effect.

Chemically modified natural zeolites have been associated with bactericidal effects on pathogenic organisms in the guts of birds. A reduction in mortality of broiler chickens and reduced viable counts of *Salmonella enteritidis* and *Escherichia coli* in the proximal and distal gut were associated with inclusion of zeolite in feed, [23]. Zeolite can be modified chemically with organic cations resulting in increased hydrophobicity of the mineral surface, increasing its

adsorptive capacity to certain molecules, and resulting in increased bactericidal effects against *Escherichia coli* and its toxins [24, 25].

The mechanisms of action of biochar, bentonite and zeolite are likely to be multifactorial. In this study we investigate their effects on the gut microbiota of birds fed with these natural products. As noted above, some studies have reported changes in the carriage of a few different bacterial species in the face of these additives based on culturing of selected pathogens. Our goal was to undertake a more comprehensive analysis of the potential suppression of pathogenic microbiota induced by the additives. High throughput DNA sequencing using the 16S ribosomal RNA gene as a phylogenetic marker is a culture free method that allows analysis of all of the complex bacterial populations that make up the gut microbiota rather than only few capable of growing on selected microbiological media. The use of these technologies over the last decade has revealed the high complexity of microbiotas from many different ecological niches including the gastrointestinal tracts (GIT) of animals and humans. The evolving understanding of the many roles and biological functions that are affected by the GIT microbiota has led to the consideration of the GIT microbiota as the largest organ of the body, contributing 10 times more cells and 100 times more gene products to the host than host's own cells. Research on chickens has shown that some commercial practices, such as the addition of antibiotics that reduce microbial diversity in the gut [26], and sanitary hatching practices that remove the influence of maternal microbiota from the hatchlings [27], lead to the formation of distorted microbial communities that may be poorly adapted to digest complex avian diets. To achieve the goal of a healthy and productive chicken GIT, free of major zoonotic and chicken pathogens, without disturbing the natural beneficial intestinal bacteria, alternative treatments and management practices are needed. These must retain the richness and diversity of the chicken GIT microbiota while selectively reducing pathogens that are potentially deleterious to the chicken or consumers of chicken products.

## Materials and Methods

### Animal trial and sample collection

All procedures involving animals were approved by the Animal Ethics Committee of the Central Queensland University (Approval number A 12/06-283). The trial was conducted in a screened shed environment with temperature variation from 22.5 to 36.5°C on the Central Queensland University Rockhampton campus, from May 2014 to December 2014. Eighty Bond Brown Layer (BBL) 17 week old pullets were obtained from Bond Enterprises P/L, Grantham, Qld, Australia. This poultry breed is a brown egg commercial layer developed by Bond Enterprises, marketed as possessing the traits of 95% peak egg production and a feed conversion ratio (FCR) of 2.1–2.3 kg feed/kg of eggs.

The pullets were allowed one week to adjust to the new environment before introducing new diet treatments. The regular commercial layer ration (Blue Ribbon Stocks Feed, Rockhampton) consisted of 90.35% dry matter, metabolisable energy 11.75 MJ/kg, crude protein 17.5% and calcium 4.2% by dry weight, supplied as a crumbed mix. A professional animal nutritionist ([www.wadeagriculture.com.au/](http://www.wadeagriculture.com.au/)) undertook the feed formulation to maintain energy value, protein content and amino acid composition across all treatments. (Table A in [S1 File](#)) Control and amended feeds were all manufactured using the same process, being pelleted first and then crumbled.

Each treatment was applied to five birds in each of four pens (ie. 20 birds per diet treatment). Treatments involved the unsupplemented commercial layer diet (control group) and this ration amended with biochar (BC), bentonite (ZT) and zeolite (ZT) at 4% w/w, with adjustment to maintain the same feed value (calcium, protein, essential amino acids and

metabolisable energy levels) to each group with  $n = 20$  birds. Additives were blended into the ration by Blue Ribbon Stocks Feed, Rockhampton Qld, using a grinder, mixer and feed pelletiser. Biochar was sourced from Pacific Pyrolysis (Sydney), being woody green waste subjected to pyrolysis at 550°C. The biochar contained 76.1% C, 3.16% H, 0.29% N and 0.03% S, and effective cation exchange capacity (ECEC) of 29.7 cmol+/kg. Bentonite was sourced from JNJ Resources P/L (Brisbane, Qld) who market bentonite as a binder for pelletising feed (typically used at up to 2% w/w) and as a feed additive to bind toxins and for alleviation of urea poisoning (through its cation exchange capacity). The material has a claimed ammonia exchange capacity of 76 meq/100g DW, total composition of 67% SiO<sub>2</sub> and 22% w/w Al<sub>2</sub>O<sub>3</sub> (Table B in [S1 File](#)). Zeolite was sourced from Castle Mountain Zeolite (Quirindi, NSW). The material is marketed for use with poultry to increase nitrogen use efficiency and for binding of dietary toxins (<http://www.cmzeolites.com.au/animals/poultry>). The material has a claimed ammonia exchange capacity of 156 meq/100g DW, a total composition of 72% SiO<sub>2</sub> and 12% w/w Al<sub>2</sub>O<sub>3</sub>, and a zeolite composition of 85% clinoptilolite and 15% mordenite. An X ray diffraction analysis suggested a composition of 17% plagioclase feldspar, 20% quartz, 5% pyroxene and 56% w/w of a zeolite likely to be clinoptilolite.

The treatment was continued for 23 weeks until the birds were 41 weeks of age. Birds were housed in a commercial layer caging system (each cage being 60×60×50 cm in height, width and depth, respectively). Pullets were randomly assigned to the cages, with five birds in each unit. Four pens with five birds each were used for each treatment, within a randomised block layout. Thus, a total of 20 birds were used for each treatment.

Water was supplied via two 'on-demand' nipples per cage. Feed of known weight was supplied daily, with unused material weighed. Birds were weighed individually every 15 days. Eggs were collected daily, counted and weighed.

## Microbial sampling and DNA preparation

Cloacal samples were taken using a sterile swab at 37 weeks of bird age. DNA samples from each group were selected for 16S rRNA gene amplicon generation and sequencing. Removing samples with low sequence number resulted in  $n = 12-15$  per treatment for the microbiota analysis.

Total DNA was isolated using Bioline ISOLATE Faecal DNA Kit (#BIO-52038) according to the manufacturer's instructions. DNA amplification was performed using Q5 DNA polymerase (New England Biolabs). Sequencing was completed on an Illumina MiSeq system (2 x 300 bp) using the dual-indexing, variable spacer, method detailed by Fadrosh *et al.* [28]. Sequencing outputs were analysed in Qiime version 1.9.1 software [29] using Qiime default parameters except for split library demultiplexing where only sequences with Phred quality threshold higher than 20 were retained for analysis (default is 3). OTUs were picked using the Uclust algorithm [30] and inspected for chimeric sequences using Pintail [31]. Taxonomy was assigned using blast against GreenGenes database [32]. Additional taxonomic assignment was done using a command line version of blastn [33] against the 16S Microbial database. The complete dataset for this experiment is publically available on the MG-RAST server under project ID 4693702.3.

The analysis was performed using data rarefied to 1850 sequences per sample removing samples with lower coverage. Statistical comparisons of the microbiota composition between the different treatment groups were performed using ANOVA on square root transformed data and Tukey Honest Significant Differences (HSD) test was performed for individual group to group comparisons in R software (<https://cran.r-project.org/>). Alpha diversity comparisons between birds of different treatment groups were calculated using a two-sample nonparametric

t-test and up to  $10^4$  Monte Carlo permutations. Beta diversity statistics was based on Adonis and up to  $10^6$  permutations. Some data was visualised using Calypso (<http://bioinfo.qimr.edu.au/>).

## Results and Discussion

### Bird performance


There were no significant differences in bird weight between the four groups either before ( $p = 0.284$ ) or after ( $p = 0.905$ ) the period of diet supplementation (Fig A in [S1 File](#)). Egg production was higher in the groups with the additives, but differences between feed additive types were not statistically significant. Although there was no significant difference in number of eggs ( $p = 0.053$ ), all three treatments had higher egg production than control. Average egg weight was significantly ( $p = 0.03$ ) different between the treatments and higher in all 3 treatments than in control. Control group had highest feed intake ( $p = 0.001$ ). FCR was significantly better ( $p = 0.007$ ) in all 3 treatments, with 2.12, 2.2 and 2.17 for BC, BT and ZT respectively, compared with control that required 2.4 kg feed/kg egg (Table C in [S1 File](#)).

### Microbiota response to additives

The most highly represented phyla within the cloacal microbiota of the laying hens were Actinobacteria (41%), Firmicutes (37%), Proteobacteria (13%) and Bacteroidetes (4%), while phyla present in low abundance ( $<0.01\%$ ) included Fusobacteria, Tenericutes, Verrucomicrobia and TM7, with traces of Thermi, Elusimicrobia, Deferribacteres and Gemmatimonadetes.


Feed amendment with biochar, bentonite or zeolite did not result in altered microbial community richness and diversity compared to the normal unsupplemented feed group. The indices inspected included richness and evenness index (Fig B in [S1 File](#)), Shannon, Simpson, Chao1, dominance, observed species and Fisher's alpha. The lack of influence on richness and diversity was evident at all phylogenetic levels.

There were no major community shifts caused by addition of BC, BT or ZT as shown by either unweighted ( $p = 0.2019$ ) or weighted ( $p = 0.4817$ ) UniFrac. A redundancy analysis (RDA) ordination plot ([Fig 1](#)) showed slight differentiation ( $p = 0.143$ , 1999 permutations)



**Fig 1. OTU level redundancy analysis (RDA) plot comparing chicken cloacal samples of birds fed control diet (CTRL) and groups with feed supplemented with biochar (BC), bentonite (BT) and zeolite (ZT).** Although the ordination plot shows some separation of control and ZT groups and strong overlap of BC and BT, the first and second ordination axes represent only 2% of the variability in the data set and the separation of the groups was not statistically significant ( $p = 0.143$ , 1999 permutations).

doi:10.1371/journal.pone.0154061.g001




**Fig 2. Influence of biochar (BC), bentonite (BT) and zeolite (ZT) feed supplementation on abundance of phylum Proteobacteria in chicken cloaca.** Phylum Proteobacteria was significantly different between the three additives and control. The reduction of Proteobacteria in additive groups was due to significant alterations in two of its classes: Epsilonproteobacteria ( $p = 0.0179$ ) and Gammaproteobacteria ( $p = 0.0191$ ) (top right panel). The Epsilonproteobacteria was comprised of only two genera—*Campylobacter* and *Helicobacter* (bottom row), each represented with only one species. *Campylobacter* OTU269490 was 100% identical to *Campylobacter jejuni* subsp. *jejuni* NCTC 11168 = ATCC 700819 strain, while *Helicobacter* OTU574168, (not significantly altered,  $p = 0.4758$ ), was identified as 99% identical *H. pullorum*. *Campylobacter jejuni* was reduced from mean of relative abundance 0.0013, equalling 1.3%, in control diet fed birds to mean of 0.02% in BC, completely absent in BT and down to 0.016% in ZT. The bars represent standard error for  $n > 12$ .

doi:10.1371/journal.pone.0154061.g002

between the control and supplemented groups. However, individual phylotypes at different phylogenetic levels did respond differentially to BC, BT and ZT.

The only phylum significantly affected by the feed additives was Proteobacteria. The prevalence of this phylum was reduced ( $p = 0.015$ ) in all three additive groups compared to the control group (Fig 2). A Tukey HSD test showed that only ZT ( $p = 0.015$ ) and BC ( $p = 0.0445$ ) were statistically significantly different to the control group, while the BT group ( $p = 0.0512$ ) was just above the 0.05  $p$ -value cut-off. The reduction of Proteobacteria was due to significant alterations in abundance of the classes Epsilonproteobacteria ( $p = 0.0179$ ) and Gammaproteobacteria ( $p = 0.0191$ ) (Fig 2). The Epsilonproteobacteria was comprised of one order, Campylobacterales ( $p = 0.0179$ ), which was comprised of only two OTUs (OTU269490 and OTU574168) belonging to *Campylobacter* and *Helicobacter* genera, respectively (Fig 2). *Salmonella* sp. were detected in only few birds with <5 sequences in each bird and were removed during the filtering steps.




**Fig 3. Bacterial families significantly ( $p < 0.05$ ) differed between birds fed control diet (Ctrl) and groups with feed supplemented with biochar (BC), bentonite (BT) and zeolite (ZT).** The families altered were members of order Actinomycetales (marked with “Act” above the bar chart) or phylum Proteobacteria’s Alphaproteobacteria, Betaproteobacteria, Epsilonproteobacteria or Gammaproteobacteria (also marked above the bar chart). Families induced in BT Bradyrhizobiaceae, Phyllobacteriaceae, are plant-associated bacteria, Sphingomonadaceae is a candidate for bioremediation and Oxalobacteraceae are known as nitrogen fixing. The bars represent standard error for  $n > 12$ .

doi:10.1371/journal.pone.0154061.g003

OTU269490 ( $p = 0.0362$ ), was identified, using blastn against the 16S Microbial database, as 100% identical to *Campylobacter jejuni* subsp. *jejuni* NCTC 11168 = ATCC 700819 strain across the 405nt length of the amplified OTU sequence. Although the BC, BT and ZT groups all had lower *Campylobacter* numbers compared to the control group, only the BT group was statistically significant in that reduction (Tukey HSD,  $p = 0.0244$ ).

OTU574168 (not significantly altered,  $p = 0.4758$ ) was identified as 99% identical to *Helicobacter pullorum*. (Fig 2). Despite *Helicobacter* being undetected in the BT group there was no statistically significant difference ( $p = 0.4069$ ) to the control group due to high variation in control group and not all birds having *Helicobacter* present.

Families (Fig 3) and genera (Fig 4) significantly altered were either members of order Actinomycetales or phylum Proteobacteria. There were 65 OTUs that were differentiated in abundance between the treatment groups and control group. Blast taxonomic assignments and bar charts showing the distribution between the groups in each of the significantly altered OTUs are shown in Table D and Fig C in S1 File. Only 17 of the 65 significantly altered OTUs showed




**Fig 4. Bacterial genera significantly ( $p < 0.05$ ) differed between birds fed control diet (Ctrl) and groups with feed supplemented with biochar (BC), bentonite (BT) and zeolite (ZT).** Members of order Actinomycetales are marked with "Act" above the bars chart, while other marking indicates genera belonging to Betaproteobacteria (Beta), Epsilonproteobacteria (Eps) or Gammaproteobacteria (Gamma). The bars represent standard error for  $n > 12$ .

doi:10.1371/journal.pone.0154061.g004

sequence similarity to entries in the 16S Microbial database with higher than 95% similarity. Out of the differentially abundant phylotypes with sequence similarity  $> 95\%$  there were a number of OTUs that represented known potential pathogens that were either reduced (six OTUs, Fig 5A) or increased (two OTUs, Fig 5B) by BC, BT and/or ZT.

The uses of biochar, bentonite and zeolite as feed additives potentially have multiple benefits for the chickens. Here we present the first sequencing-based, culture-independent assessment of their effects on the gut microbiota. Most of the classic antimicrobial growth promoters have a broad spectra of action. Thus they reduce richness and diversity of the intestinal microbiota, which is a risk factor for dysbiosis and other gastrointestinal complications [34]. We demonstrated that biochar, bentonite and zeolite can be used to selectively reduce the abundance of some major poultry zoonotic pathogens without reducing chicken microbiota diversity or introducing major microbiota shifts. This is highly relevant for the health of the animals and indeed to human health.

Biochar, bentonite and zeolite treatment groups all had lower levels of an OTU identified as 100% identical to *Campylobacter jejuni*. Although the biochar and zeolite groups did not meet the statistical criteria of significance due to the high degree of variation in carriage in the control group, the reduction in the bentonite group was complete and statistically significant. The



**Fig 5. OTUs significantly ( $p < 0.05$ ) differed between the groups with high sequence alignment identity (>95%) with known pathogenic strains.** Out of 65 OTUs significantly altered between the treatment groups (Fig C in S1 File), the majority could not be provisionally assigned to species or genus level using 97% or 95% similarity cut-off. Among the phylotypes with sequence similarity >95%, some candidates showed high sequence alignment (blast against 16S microbial database) to known pathogens. Panel A shows potentially pathogenic OTUs reduced in additives while panel B shows potentially pathogenic OTUs increased in additive groups. Y axes indicate % of abundance in each group. The bars represent standard error for  $n > 12$ .

doi:10.1371/journal.pone.0154061.g005

Campylobacteriaceae family includes many human and animal pathogenic species; the most significant *C. jejuni*, is recognized as a leading cause of foodborne infections. The European Food Safety Authority (EFSA) has estimated that within the European Union there are 9 million cases of *Campylobacter* food poisoning annually, with related cost of EUR 2.4 billion per year, 20–30% of which cases are attributed to contaminated chicken meat [35].

Shifts and dysbiosis in intestinal microbiota are reported as the first steps in successful colonization by *C. jejuni* [36, 37] and possible disease onset. The bacterium is a common member of the intestinal microbiota in wildlife and agricultural animals, where it usually exists asymptotically. These animals thus represent a reservoir of this major zoonotic human disease.

Additionally, we detected the ability of bentonite to reduce a *Helicobacter pullorum*-related OTU. This species, which was completely absent from the gut of all birds on diet supplemented with bentonite, aligned with 99% similarity to *H. pullorum*. This enterohepatic helicobacter species is found in the gut of healthy chickens as well as in liver and intestine of hens with vibriotic-like liver lesions [38] and in human chronic liver disease [39]. It is also associated with human gastroenteritis, inflammatory bowel disease and Chron’s disease [40–42]. The ability to

reduce both *Helicobacter* and *Campylobacter*, if reproducible under varied feeding and environmental conditions, would be a compelling reason to include bentonite in poultry feed formulations.

Biochar and zeolite also affected the carriage of these potentially pathogenic species, although not as significantly as bentonite. Zeolite was found to significantly reduce the levels of the phylum Proteobacteria. High numbers of Proteobacteria are considered a sign of bad intestinal health and are associated with gastrointestinal health conditions such as chronic dysbiosis [43] and inflammatory bowel disease [44].

All three additives reduced the abundance of four OTUs identified with high sequence identity (>97%) to *Gallibacterium anatis*, another major poultry pathogen [45]. *G. anatis* has been associated with a range of pathological lesions mostly in breeding chicken, including peritonitis, oophoritis, septicemia, follicle degeneration, salpingitis and respiratory tract infections [46]. Together with *Pasteurella multocida*, *G. anatis* causes fowl cholera-like clinical manifestations and lesions. Although *P. multocida* was not identified at the species level, the family Pasteurellaceae was abundant (higher than 20% in control, Fig 3) and was significantly reduced in all three additive groups, most significantly with bentonite (Fig 3).

*Bacteroides dorei* (99% identity) and *Clostridium aldenense* (95% identity) were present in control birds only, as low abundance taxa, and were absent from microbiota in all three additive groups. *Clostridium aldenense* is another pathogen involved in multiple clinical presentations including bacteremia [47, 48]. If further investigated and confirmed, the ability of additives to remove *B. dorei* would be of interest for human health. For example, *B. dorei* was found to dominate gut microbiome prior to onset of autoimmunity in children at high risk for type 1 diabetes [49, 50].

*Enterococcus cecorum* and *Corynebacterium amycolatum* are potential pathogens that were increased in some additive groups. *E. cecorum* was increased in the biochar group and *C. amycolatum* in the zeolite group. *E. cecorum* is an animal pathogen [51] as well as opportunistic human pathogen [52–54]. In chickens it is a major cause of outbreaks of arthritis and osteomyelitis worldwide [55]. *C. amycolatum* has been occasionally associated with infective endocarditis [56]. There were no affected OTUs similar to known pathogens with >95% sequence similarity that were significantly increased in bentonite.

## Conclusions

The data presented here indicate that zeolite, biochar, and in particular bentonite may be viable alternatives to AGPs in poultry and other agricultural industries that could be further developed to help control pathogen load in domestic animals without significantly changing the overall complexity of gut microbiota. Of special interest is the association of bentonite with decreased *Campylobacter* and *Helicobacter* genera. This is especially promising since bentonite has been used and proven as safe through occasional use in poultry feeds [57, 58] for purposes other than pathogen control, mostly to improve feed manufacture [10] and for reduction of the feed passage rate through the chicken gut [15].

## Supporting Information

**S1 File.**  
(PDF)

## Acknowledgments

We wish to thank Jason Bell, of CQU High Performance Computing, for continuous help and support. We also thank Ngare Wilkinson, Giselle Weegenaar and Ingrid Christiansen for their

help with histology and molecular work, and Mr Garry Smalls of Smalls Poultry, Mount Morgan for his support and contribution to building of the caging units.

## Author Contributions

Conceived and designed the experiments: KW SB DM RM DS. Performed the experiments: TP DS TV RM. Analyzed the data: DS. Contributed reagents/materials/analysis tools: KW RM DS. Wrote the paper: TP KW SB DM TV RM DS.

## References

1. Huyghebaert G, Ducatelle R, Van Immerseel F. An update on alternatives to antimicrobial growth promoters for broilers. *The Veterinary Journal*. 2011; 187(2):182–8. doi: [10.1016/j.tvjl.2010.03.003](https://doi.org/10.1016/j.tvjl.2010.03.003) PMID: [20382054](https://pubmed.ncbi.nlm.nih.gov/20382054/)
2. Marshall BM, Levy SB. Food animals and antimicrobials: impacts on human health. *Clinical Microbiology Reviews*. 2011; 24(4):718–33. doi: [10.1128/CMR.00002-11](https://doi.org/10.1128/CMR.00002-11) PMID: [21976606](https://pubmed.ncbi.nlm.nih.gov/21976606/)
3. Crawshaw TR, Chanter JI, Young SC, Cawthraw S, Whatmore AM, Koylass MS, et al. Isolation of a novel thermophilic *Campylobacter* from cases of spotty liver disease in laying hens and experimental reproduction of infection and microscopic pathology. *Vet Microbiol*. 2015; 179(3–4):315–21. doi: [10.1016/j.vetmic.2015.06.008](https://doi.org/10.1016/j.vetmic.2015.06.008) PMID: [26130518](https://pubmed.ncbi.nlm.nih.gov/26130518/).
4. Gerlach H, Schmidt H-P. Biochar in poultry farming. *Ithaka Journal*. 2012; 2012:262–4.
5. Papaioannou D, Katsoulos PD, Panousis N, Karatzias H. The role of natural and synthetic zeolites as feed additives on the prevention and/or the treatment of certain farm animal diseases: A review. *Microporous and Mesoporous Materials*. 2005; 84(1–3):161–70. doi: [10.1016/j.micromeso.2005.05.030](https://doi.org/10.1016/j.micromeso.2005.05.030)
6. Pasha T, Farooq M, Khattak F, Jabbar M, Khan A. Effectiveness of sodium bentonite and two commercial products as aflatoxin absorbents in diets for broiler chickens. *Animal Feed Science and Technology*. 2007; 132(1–2):103–10. doi: <http://dx.doi.org/10.1016/j.anifeedsci.2006.03.014>.
7. Kutlu HR, Ünsal I, Görgülü M. Effects of providing dietary wood (oak) charcoal to broiler chicks and laying hens. *Animal feed science and technology*. 2001; 90(3–4):213–26. doi: [10.1016/S0377-8401\(01\)00205-X](https://doi.org/10.1016/S0377-8401(01)00205-X)
8. Kana J, Tegui A, Tchoumboue J. Effect of dietary plant charcoal from *Canarium schweinfurthii* Engl. and maize cob on aflatoxin B1 toxicosis in broiler chickens. *Adv Anim Biosci*. 2010; 1:462–3.
9. Ruttanavut J, Yamauchi K, Goto H, Erikawa T. Effects of Dietary Bamboo Charcoal Powder Including Vinegar Liquid on Growth Performance and Histological Intestinal Change in Aigamo Ducks. *International Journal of Poultry Science*. 2009; 8(3):229–36. doi: [10.3923/ijps.2009.229.236](https://doi.org/10.3923/ijps.2009.229.236)
10. Angulo E, Brufau J, Esteve-Garcia E. Effect of sepiolite on pellet durability in feeds differing in fat and fibre content. *Animal Feed Science and Technology*. 1995; 53(3):233–41.
11. Cabezas M, Salvador D, Sinisterra J. Stabilization-activation of pancreatic enzymes adsorbed on to a sepiolite clay. *Journal of chemical technology and biotechnology*. 1991; 52(2):265–74.
12. Parisini P, Martelli G, Sardi L, Escribano F. Protein and energy retention in pigs fed diets containing sepiolite. *Animal Feed Science and Technology*. 1999; 79(1):155–62.
13. Ouhida I, Perez J, Piedrafita J, Gasa J. The effects of sepiolite in broiler chicken diets of high, medium and low viscosity. Productive performance and nutritive value. *Animal Feed Science and Technology*. 2000; 85(3):183–94.
14. Alzueta C, Ortiz L, Rebole A, Rodríguez M, Centeno C, Treviño J. Effects of removal of mucilage and enzyme or sepiolite supplement on the nutrient digestibility and metabolizable energy of a diet containing linseed in broiler chickens. *Animal Feed Science and Technology*. 2002; 97(3):169–81.
15. Damiri H, Chaji M, Bojarpour M, Eslami M, Mamoei M. The effect of sodium bentonites on economic value of broiler chickens diet. *Journal of Animal and Veterinary Advances*. 2010; 9:2668–26670.
16. Moghaddam HN, Jahanian R, Najafabadi HJ, Madaeni M. Influence of dietary zeolite supplementation on the performance and egg quality of laying hens fed varying levels of calcium and nonphytate phosphorus. *Journal of Biological Sciences*. 2008; 8(2):328–34.
17. Xia M, Hu C, Xu Z. Effects of copper-bearing montmorillonite on growth performance, digestive enzyme activities, and intestinal microflora and morphology of male broilers. *Poultry science*. 2004; 83(11):1868–75. PMID: [15554064](https://pubmed.ncbi.nlm.nih.gov/15554064/)
18. Eleroğlu H, Yalçın H, Yıldırım A. Dietary effects of Ca-zeolite supplementation on some blood and tibial bone characteristics of broilers. *South African Journal of Animal Science*. 2011; 41(4):319–30.

19. Fethiere R, Miles R, Harms R. The utilization of sodium in sodium zeolite A by broilers. *Poultry Science*. 1994; 73(1):118–21. PMID: [8165157](#)
20. Olver M. Effect of feeding clinoptilolite (zeolite) on the performance of three strains of laying hens. *British Poultry Science*. 1997; 38(2):220–2. PMID: [9158901](#)
21. Papaioannou D, Kyriakis S, Papasteriadis A, Roubies N, Yannakopoulos A, Alexopoulos C. Effect of in-feed inclusion of a natural zeolite (clinoptilolite) on certain vitamin, macro and trace element concentrations in the blood, liver and kidney tissues of sows. *Research in veterinary science*. 2002; 72(1):61–8. PMID: [12002639](#)
22. Fethiere R, Miles R, Harms R. Influence of synthetic sodium aluminosilicate on laying hens fed different phosphorus levels. *Poultry science*. 1990; 69(12):2195–8. PMID: [1964737](#)
23. Olver M. effect of feeding clinoptilolite (zeolite) to laying hens. *South African Journal of Animal Science = Suid-Afrikaanse tydskrif vir veekunde*. 1983.
24. Uchida T, Maru N, Furuhashi M, Fujino A, Muramoto S, Ishibashi A, et al. Anti-bacterial zeolite balloon catheter and its potential for urinary tract infection control. *Hinyokika kyo Acta Urologica Japonica*. 1992; 38(8):973–8. PMID: [1329451](#)
25. Daković A, Tomašević-Čanović M, Dondur V, Rottinghaus GE, Medaković V, Zarić S. Adsorption of mycotoxins by organozeolites. *Colloids and Surfaces B: biointerfaces*. 2005; 46(1):20–5. PMID: [16198090](#)
26. Lin J, Hunkapiller AA, Layton AC, Chang YJ, Robbins KR. Response of intestinal microbiota to antibiotic growth promoters in chickens. *Foodborne Pathog Dis*. 2013; 10(4):331–7. doi: [10.1089/fpd.2012.1348](#) PMID: [23461609](#).
27. Stanley D, Geier MS, Hughes RJ, Denman SE, Moore RJ. Highly variable microbiota development in the chicken gastrointestinal tract. *PLoS One*. 2013; 8(12):e84290. doi: [10.1371/journal.pone.0084290](#) PMID: [24391931](#); PubMed Central PMCID: PMC3877270.
28. Fadrosch DW, Ma B, Gajer P, Sengamalay N, Ott S, Brotman RM, et al. An improved dual-indexing approach for multiplexed 16S rRNA gene sequencing on the Illumina MiSeq platform. *Microbiome*. 2014; 2(1):6. doi: [10.1186/2049-2618-2-6](#) PMID: [24558975](#); PubMed Central PMCID: PMC3940169.
29. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods*. 2010; 7(5):335–6. Epub 2010/04/13. [pii] doi: [10.1038/nmeth.f.303](#) PMID: [20383131](#).
30. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*. 2010; 26(19):2460–1. Epub 2010/08/17. [pii] doi: [10.1093/bioinformatics/btq461](#) PMID: [20709691](#).
31. Ashelford KE, Chuzhanova NA, Fry JC, Jones AJ, Weightman AJ. At least 1 in 20 16S rRNA sequence records currently held in public repositories is estimated to contain substantial anomalies. *Appl Environ Microbiol*. 2005; 71(12):7724–36. Epub 2005/12/08. [pii] doi: [10.1128/AEM.71.12.7724-7736.2005](#) PMID: [16332745](#); PubMed Central PMCID: PMC1317345.
32. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol*. 2006; 72(7):5069–72. Epub 2006/07/06. [pii] doi: [10.1128/AEM.03006-05](#) PMID: [16820507](#); PubMed Central PMCID: PMC1489311.
33. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*. 1997; 25(17):3389–402. Epub 1997/09/01. [pii]. PMID: [9254694](#); PubMed Central PMCID: PMC146917.
34. Scher JU, Ubeda C, Artacho A, Attur M, Isaac S, Reddy SM, et al. Decreased bacterial diversity characterizes the altered gut microbiota in patients with psoriatic arthritis, resembling dysbiosis in inflammatory bowel disease. *Arthritis Rheumatol*. 2015; 67(1):128–39. doi: [10.1002/art.38892](#) PMID: [25319745](#); PubMed Central PMCID: PMC4280348.
35. EFSA. European Food safety Authority 2015. Available from: <http://www.efsa.europa.eu/en/topics/topic/campylobacter>.
36. Haag LM, Fischer A, Otto B, Plickert R, Kuhl AA, Gobel UB, et al. Intestinal microbiota shifts towards elevated commensal *Escherichia coli* loads abrogate colonization resistance against *Campylobacter jejuni* in mice. *PLoS One*. 2012; 7(5):e35988. doi: [10.1371/journal.pone.0035988](#) PMID: [22563475](#); PubMed Central PMCID: PMC3341396.
37. Lone AG, Selinger LB, Uwiera RR, Xu Y, Inglis GD. *Campylobacter jejuni* colonization is associated with a dysbiosis in the cecal microbiota of mice in the absence of prominent inflammation. *PLoS One*. 2013; 8(9):e75325. doi: [10.1371/journal.pone.0075325](#) PMID: [24066174](#); PubMed Central PMCID: PMC3774657.
38. Stanley J, Linton D, Burnens AP, Dewhirst FE, On SL, Porter A, et al. *Helicobacter pullorum* sp. nov.-genotype and phenotype of a new species isolated from poultry and from human patients with

- gastroenteritis. *Microbiology*. 1994; 140 (Pt 12):3441–9. doi: [10.1099/13500872-140-12-3441](https://doi.org/10.1099/13500872-140-12-3441) PMID: [7533595](https://pubmed.ncbi.nlm.nih.gov/7533595/).
39. Veijola L, Nilsson I, Halme L, Al-Soud WA, Makinen J, Ljungh A, et al. Detection of *Helicobacter* species in chronic liver disease and chronic inflammatory bowel disease. *Ann Med*. 2007; 39(7):554–60. doi: [10.1080/07853890701545714](https://doi.org/10.1080/07853890701545714) PMID: [17852032](https://pubmed.ncbi.nlm.nih.gov/17852032/).
  40. Hansen R, Thomson JM, Fox JG, El-Omar EM, Hold GL. Could *Helicobacter* organisms cause inflammatory bowel disease? *FEMS Immunol Med Microbiol*. 2011; 61(1):1–14. doi: [10.1111/j.1574-695X.2010.00744.x](https://doi.org/10.1111/j.1574-695X.2010.00744.x) PMID: [20955468](https://pubmed.ncbi.nlm.nih.gov/20955468/).
  41. Laharie D, Asencio C, Asselineau J, Bulois P, Bourreille A, Moreau J, et al. Association between entero-hepatic *Helicobacter* species and Crohn's disease: a prospective cross-sectional study. *Aliment Pharmacol Ther*. 2009; 30(3):283–93. doi: [10.1111/j.1365-2036.2009.04034.x](https://doi.org/10.1111/j.1365-2036.2009.04034.x) PMID: [19438427](https://pubmed.ncbi.nlm.nih.gov/19438427/).
  42. Steinbrueckner B, Haerter G, Pelz K, Weiner S, Rump JA, Deissler W, et al. Isolation of *Helicobacter pullorum* from patients with enteritis. *Scand J Infect Dis*. 1997; 29(3):315–8. PMID: [9255900](https://pubmed.ncbi.nlm.nih.gov/9255900/).
  43. Shin NR, Whon TW, Bae JW. Proteobacteria: microbial signature of dysbiosis in gut microbiota. *Trends Biotechnol*. 2015; 33(9):496–503. doi: [10.1016/j.tibtech.2015.06.011](https://doi.org/10.1016/j.tibtech.2015.06.011) PMID: [26210164](https://pubmed.ncbi.nlm.nih.gov/26210164/).
  44. Mukhopadhyay I, Hansen R, El-Omar EM, Hold GL. IBD-what role do Proteobacteria play? *Nature reviews Gastroenterology & hepatology*. 2012; 9(4):219–30. doi: [10.1038/nrgastro.2012.14](https://doi.org/10.1038/nrgastro.2012.14) PMID: [22349170](https://pubmed.ncbi.nlm.nih.gov/22349170/).
  45. Paudel S, Liebhart D, Hess M, Hess C. Pathogenesis of *Gallibacterium anatis* in a natural infection model fulfils Koch's postulates: 1. Folliculitis and drop in egg production are the predominant effects in specific pathogen free layers. *Avian Pathol*. 2014; 43(5):443–9. doi: [10.1080/03079457.2014.955782](https://doi.org/10.1080/03079457.2014.955782) PMID: [25144260](https://pubmed.ncbi.nlm.nih.gov/25144260/).
  46. Neubauer C, De Souza-Pilz M, Bojesen AM, Bisgaard M, Hess M. Tissue distribution of haemolytic *Gallibacterium anatis* isolates in laying birds with reproductive disorders. *Avian Pathol*. 2009; 38(1):1–7. doi: [10.1080/03079450802577848](https://doi.org/10.1080/03079450802577848) PMID: [19089694](https://pubmed.ncbi.nlm.nih.gov/19089694/).
  47. Daimon Y, Tanaka K, Watanabe K. [Epidemiological study on cigar-shaped clostridia isolated in a local Japanese general hospital]. *Kansenshogaku Zasshi*. 2008; 82(3):205–12. PMID: [18546850](https://pubmed.ncbi.nlm.nih.gov/18546850/).
  48. Williams OM, Brazier J, Peraino V, Goldstein EJ. A review of three cases of *Clostridium aldenense* bacteremia. *Anaerobe*. 2010; 16(5):475–7. doi: [10.1016/j.anaerobe.2010.08.004](https://doi.org/10.1016/j.anaerobe.2010.08.004) PMID: [20800690](https://pubmed.ncbi.nlm.nih.gov/20800690/).
  49. Davis-Richardson AG, Ardissonne AN, Dias R, Simell V, Leonard MT, Kemppainen KM, et al. *Bacteroides dorei* dominates gut microbiome prior to autoimmunity in Finnish children at high risk for type 1 diabetes. *Frontiers in microbiology*. 2014; 5:678. doi: [10.3389/fmicb.2014.00678](https://doi.org/10.3389/fmicb.2014.00678) PMID: [25540641](https://pubmed.ncbi.nlm.nih.gov/25540641/); PubMed Central PMCID: [PMC4261809](https://pubmed.ncbi.nlm.nih.gov/PMC4261809/).
  50. Davis-Richardson AG, Triplett EW. A model for the role of gut bacteria in the development of autoimmunity for type 1 diabetes. *Diabetologia*. 2015; 58(7):1386–93. doi: [10.1007/s00125-015-3614-8](https://doi.org/10.1007/s00125-015-3614-8) PMID: [25957231](https://pubmed.ncbi.nlm.nih.gov/25957231/); PubMed Central PMCID: [PMC4473028](https://pubmed.ncbi.nlm.nih.gov/PMC4473028/).
  51. Boerlin P, Nicholson V, Brash M, Slavic D, Boyen F, Sanei B, et al. Diversity of *Enterococcus cecorum* from chickens. *Vet Microbiol*. 2012; 157(3–4):405–11. doi: [10.1016/j.vetmic.2012.01.001](https://doi.org/10.1016/j.vetmic.2012.01.001) PMID: [22266160](https://pubmed.ncbi.nlm.nih.gov/22266160/).
  52. De Baere T, Claeys G, Verschraegen G, Devriese LA, Baele M, Van Vlem B, et al. Continuous ambulatory peritoneal dialysis peritonitis due to *Enterococcus cecorum*. *J Clin Microbiol*. 2000; 38(9):3511–2. PMID: [10970419](https://pubmed.ncbi.nlm.nih.gov/10970419/); PubMed Central PMCID: [PMC87422](https://pubmed.ncbi.nlm.nih.gov/PMC87422/).
  53. Delaunay E, Abat C, Rolain JM. *Enterococcus cecorum* human infection, France. *New microbes and new infections*. 2015; 7:50–1. doi: [10.1016/j.nmni.2015.06.004](https://doi.org/10.1016/j.nmni.2015.06.004) PMID: [26199733](https://pubmed.ncbi.nlm.nih.gov/26199733/); PubMed Central PMCID: [PMC4506978](https://pubmed.ncbi.nlm.nih.gov/PMC4506978/).
  54. Greub G, Devriese LA, Pot B, Dominguez J, Bille J. *Enterococcus cecorum* septicemia in a malnourished adult patient. *Eur J Clin Microbiol Infect Dis*. 1997; 16(8):594–8. PMID: [9323472](https://pubmed.ncbi.nlm.nih.gov/9323472/).
  55. Stalker MJ, Brash ML, Weisz A, Ouckama RM, Slavic D. Arthritis and osteomyelitis associated with *Enterococcus cecorum* infection in broiler and broiler breeder chickens in Ontario, Canada. *J Vet Diagn Invest*. 2010; 22(4):643–5. PMID: [20622242](https://pubmed.ncbi.nlm.nih.gov/20622242/).
  56. Belmares J, Detterline S, Pak JB, Parada JP. *Corynebacterium endocarditis* species-specific risk factors and outcomes. *BMC Infect Dis*. 2007; 7:4. doi: [10.1186/1471-2334-7-4](https://doi.org/10.1186/1471-2334-7-4) PMID: [17284316](https://pubmed.ncbi.nlm.nih.gov/17284316/); PubMed Central PMCID: [PMC1804271](https://pubmed.ncbi.nlm.nih.gov/PMC1804271/).
  57. Prvulovic D, Kojic D, Grubor-Lajsic G, Kosarcic S. The effects of dietary inclusion of hydrated aluminosilicate on performance and biochemical parameters of broiler chickens. *Turkish Journal of Veterinary and Animal Sciences*. 2008; 32(3):183–9.
  58. Safaeikatouli M, Jafariahangari Y, Baharlouei A. Effects of dietary inclusion of sodium bentonite on biochemical characteristics of blood serum in broiler chickens. *International Journal of Agriculture and Biology*. 2010; 12:877–80.