

RESEARCH ARTICLE

BRD7 Acts as a Tumor Suppressor Gene in Lung Adenocarcinoma

Yushun Gao, Bing Wang, Shugeng Gao*

Department of thoracic surgical oncology, cancer institute (hospital), Chinese academy of medical sciences, Peking union medical college, Beijing, 100021, China

* yushun1gao@163.com

Abstract

Lung cancer is one of the most malignant tumors and the leading cause of cancer-related deaths worldwide. Among lung cancers, 40% are diagnosed as adenocarcinoma. Bromodomain containing 7 (BRD7) is a member of bromodomain-containing protein family. It was proved to be downregulated in various cancers. However, the role of BRD7 in lung adenocarcinoma is still unknown. Western blot and qRT-PCR was performed to measure the BRD7 expression in lung adenocarcinoma tissues and cells. CCK8 and migration assay was done to detect the functional role of BRD7 in lung adenocarcinoma. In this study, we showed that the expression of BRD7 was downregulated in lung adenocarcinoma tissues and cells. The lower of BRD7 levels in patients with lung adenocarcinoma was associated with shortened disease-free survival. Furthermore, overexpression of BRD7 inhibited lung adenocarcinoma cell proliferation and migration. Inhibition of BRD7 expression promoted cell proliferation and migration by activating ERK phosphorylation. Overexpression of BRD7 inhibited cyclin D and myc expression. Our findings are consistent with a tumor suppressor role for BRD7 in lung adenocarcinoma tumorigenesis.



OPEN ACCESS

Citation: Gao Y, Wang B, Gao S (2016) BRD7 Acts as a Tumor Suppressor Gene in Lung Adenocarcinoma. PLoS ONE 11(8): e0156701. doi:10.1371/journal.pone.0156701

Editor: Srikumar Chellappan, H. Lee Moffitt Cancer Center & Research Institute, UNITED STATES

Received: February 19, 2016

Accepted: May 18, 2016

Published: August 31, 2016

Copyright: © 2016 Gao et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The authors received no specific funding for this work.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

Lung cancer, one of the most malignant tumors, was the leading cause of cancer-related deaths worldwide, with about 226,000 new cases in 2012 in the United States[1–5]. Among lung cancers, 40% are diagnosed as adenocarcinoma and the 5-year survival rate for lung adenocarcinoma patients is only 5–20% at later stages[6–9]. Despite recent advances in the molecular mechanisms and surgical and chemotherapeutic interventions, the prognosis of lung adenocarcinoma has not improved significantly[10, 11]. Therefore, it is an urgent to identify an efficient and novel predictive marker for lung adenocarcinoma.

Bromodomain containing 7 (BRD7), also known as NAG4, BP75 or CELTIX1, is a member of bromodomain-containing proteins family[12–14]. Recent studies demonstrated that BRD7 was mostly located in the nucleus and regulated chromatin remodeling[15–17]. More importantly, increasing evidences showed that BRD7 were downregulated in many cancers such as epithelial ovarian carcinoma, breast cancer, nasopharyngeal cancer and colorectal carcinoma [18–21]. For example, Hu et al. showed that BRD7 acted as a tumor suppressor in

osteosarcoma, and knockdown of BRD7 increased colony formation, tumor growth and cell proliferation of osteosarcoma[22]. Park et al. demonstrated that overexpression of BRD7 inhibited ovarian cancer cells invasion, apoptosis and viability[18]. However, the role of BRD7 in lung adenocarcinoma is still unknown.

In our study, we demonstrated that the expression of BRD7 was downregulated in lung adenocarcinoma tissues and cells. Moreover, lower of BRD7 levels in patients with lung adenocarcinoma was associated with shortened disease-free survival. Overexpression of BRD7 inhibited lung adenocarcinoma cell proliferation and migration. Our findings demonstrated that BRD7 played a tumor suppressor role in lung adenocarcinoma tumorigenesis.

Materials and Methods

Tissue specimens and cell lines and cell transfection

Thirty frozen tissues from lung adenocarcinoma patients were obtained in our hospital between 2010 and 2012. All tissues were immediately snapped frozen at -80°C . All protocols were approved by the Ethics Committee of the cancer institute(hospital). Written informed consent from each patient was collected. Human lung adenocarcinoma cell lines (H1299, H23, SPC-A1 and A549) were obtained from the Cell Resource Center of Institute of Chinese Academy of Medical Sciences (Beijing, China), and keep in RPMI 1640 medium. BRD7 plasmid or mock vector was transfected by using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) following to the manufacturer's instruction.

Cell proliferation and migration assay

CCK8 (Dojindo; Kumamoto, Japan) was performed to measure cell proliferation. Cells were cultured in 10% CCK-8 diluted in normal medium until visual color conversion occurred. Proliferation rates were detected at 0, 24 and 48 hours after transfection. For migration assay, a wound-healing assay was performed. An artificial wound was performed by using a 200- μl pipette tip. To measure migrated cells, pictures were taken at 0, 24 and 48 h.

Western blot

Western blot was performed using standard methods. Protein samples were loaded on 12% SDS gels and then transferred to membranes (Millipore, Danvers, MA, USA). After blocked with 5% nonfat milk, membranes were incubated for 2 hours with the following antibodies: BRD7 and GAPDH (Abcam, Cambridge, MA, USA). Proteins were measured using the enhanced chemiluminescence system (GE Healthcare).

RNA extraction and qRT-PCR

Total RNA was extracted from the tissues and cells using Trizol reagent (Invitrogen, Calsbad, CA, USA). qRT-PCR was performed to detect the BRD7 expression following previous methods on the iQ5 Real-Time PCR Detection System (Bio-Rad, California, USA). The sequences of the specific primers are shown: BRD7 sense: CTGGAGATGCCGAAGCACAC, anti-sense: TGGGATCCACAGGATGGAGA; GAPDH sense: AGCCACATCGCTCAGACAC, anti-sense: GCCCAATACGACCAAATCC.

Statistical analysis

Statistics was done by one-way ANOVA for comparing continuous variables of more than two groups, respectively. The data were shown as the mean \pm SD (standard deviation). $P\leq 0.05$ were regarded as being statistically significant.

Result

The expression level of BRD7 was downregulated in lung adenocarcinoma cell lines

The protein expression of BRD7 was downregulated in lung adenocarcinoma cell lines and lung adenocarcinoma tissue compared to the adjacent no-tumor tissue (Fig 1A and 1B). Meanwhile, the mRNA expression of BRD7 was also lower in four lung adenocarcinoma cell lines and one lung adenocarcinoma tissues compared to one adjacent no-tumor tissues (Fig 1C).

The expression level of BRD7 was downregulated in lung adenocarcinoma tissues

The mRNA expression of BRD7 was lower in lung adenocarcinoma tissues compared to adjacent no-tumor tissues (Fig 2A). The expression of BRD7 was downregulated in 22 cases (22/30, 73%) in comparison with the adjacent tissues (Fig 2B). Furthermore, tissues from lymph node metastases expressed lower levels of BRD7 compared to primary lung adenocarcinoma tissues and the adjacent normal tissue (Fig 2C). When correlated to disease outcome, lower of BRD7 levels in patients with lung adenocarcinoma was associated with shortened disease-free survival (hazards ratio = 0.36, Fig 3A). The protein level of BRD7 was also downregulated in lung adenocarcinoma tissues compared to adjacent no-tumor tissues (Fig 3B).

Overexpression of BRD7 inhibited lung adenocarcinoma cell proliferation and migration

Western blot and real-time PCR analysis showed that BRD7 vector promoted the expression of BRD7 (Fig 4A and 4B). Overexpression of BRD7 inhibited lung adenocarcinoma cell line A549 proliferation and migration (Fig 4C, 4D and 4E).

Inhibition of BRD7 promoted ERK phosphorylation in lung adenocarcinoma

Western blot and real-time PCR analysis showed that BRD7 siRNA inhibited the expression of BRD7 (Fig 5A and 5B). Inhibition of BRD7 promoted the ERK phosphorylation in the A549




Fig 1. The expression level of BRD7 was downregulated in lung adenocarcinoma cell lines. (A) The protein expression level of BRD7 in four lung adenocarcinoma cell lines (H1299, H23, SPC-A1 and A549) and one lung adenocarcinoma tissues and adjacent no-tumor tissue was measured by Western blot. (B) The signal in each lane was quantified using ImageJ software and the ratio of BRD7 to GAPDH was determined. (C) The mRNA expression level of BRD7 in four lung adenocarcinoma cell lines (H1299, H23, SPC-A1 and A549) and one lung adenocarcinoma tissues and adjacent no-tumor tissue was measured by qRT-PCR.

doi:10.1371/journal.pone.0156701.g001




Fig 2. The expression level of BRD7 was downregulated in lung adenocarcinoma tissues. (A) Relative BRD7 mRNA expression levels in lung adenocarcinoma tissues and their corresponding adjacent normal tissues. (B) qRT-PCR analysis of BRD7 expression in 30 pair's lung adenocarcinoma tissues and their corresponding adjacent normal tissues. (C) Tissues from lymph node metastases expressed lower levels of BRD7 compared to primary lung adenocarcinoma tissues and the adjacent normal tissue.

doi:10.1371/journal.pone.0156701.g002

cells (Fig 5C). Knockdown of BRD7 expression increased the A549 cells proliferation, which could be blocked by U0126 (a MEK inhibitor) (Fig 5D). Knockdown of BRD7 expression increased the A549 cells migration, which could be blocked by U0126 (Fig 5E).

Overexpression of BRD7 inhibited Cyclin D and Myc expression

qRT-PCR analysis demonstrated that overexpression of BRD7 suppressed the cyclin D expression in the A549 cells (Fig 6A). BRD7 overexpression inhibited the myc expression in the A549 cells. Moreover, knockdown of BRD7 expression increased cyclin D expression in the A549 cells, which could be blocked by U0126 (Fig 6C). In addition, knockdown of BRD7 expression increased myc expression in the A549 cells, which could be blocked by U0126 (Fig 6D).

Discussion

As one of the most common cancers, lung adenocarcinoma was the leading cause of cancer death in the recent years. The most important prognostic factor for lung adenocarcinoma is the clinical stages [23, 24]. Unfortunately, even at the same clinical pathological stage, outcomes of patients vary considerably [25, 26]. This phenomenon is frequently observed in lung adenocarcinoma patients with clinical stage III and IV [27, 28]. Thereby, improving prognostic markers for clinical use is urgently needed.

In our study, we demonstrated that the expression of BRD7 was downregulated in lung adenocarcinoma tissues and cells. Moreover, the lower of BRD7 levels in patients with lung




Fig 3. The protein expression level of BRD7 was downregulated in lung adenocarcinoma tissues. (A) Lower of BRD7 levels in patients with lung adenocarcinoma was associated with shortened disease-free survival (hazards ratio = 0.36). (B) The protein expression level of BRD7 was measured in lung adenocarcinoma tissues and their corresponding adjacent normal tissues using Western blot.

doi:10.1371/journal.pone.0156701.g003




Fig 4. Overexpression of BRD7 inhibited lung adenocarcinoma cell proliferation and migration. (A) The protein expression level of BRD7 was measured by using Western blot. (B) The mRNA expression level of BRD7 was measured by using qRT-PCR. (C) Overexpression of BRD7 inhibited lung adenocarcinoma cell line A549 proliferation. (D) Overexpression of BRD7 inhibited lung adenocarcinoma cell line A549 migration. (E) Relative ratio of wound closure per field is shown. * $p < 0.05$ and *** $p < 0.001$.

doi:10.1371/journal.pone.0156701.g004

adenocarcinoma was associated with shortened disease-free survival. BRD7 was also downregulated in lung adenocarcinoma tissues compared to adjacent no-tumor tissues. Furthermore, overexpression of BRD7 inhibited lung adenocarcinoma cell proliferation and migration while inhibition of BRD7 expression promoted cell proliferation and migration by activating ERK




Fig 5. Inhibition of BRD7 promoted ERK phosphorylation in lung adenocarcinoma. (A) The protein expression level of BRD7 was measured by using Western blot. (B) The mRNA expression level of BRD7 was measured by using qRT-PCR. (C) Inhibition of BRD7 promoted the ERK phosphorylation in the A549 cells. (D) CCK8 analysis was used to measure the A549 cell proliferation. (E) Overexpression of BRD7 suppressed the ERK phosphorylation in the A549 cells. (F) Wound-healing assay was performed to measure the A549 cell migration. (G) Relative ratio of wound closure per field is shown. * $p < 0.05$ and *** $p < 0.001$.

doi:10.1371/journal.pone.0156701.g005




Fig 6. Overexpression of BRD7 inhibited Cyclin D and Myc expression. (A) Overexpression of BRD7 suppressed the Cyclin D mRNA expression in the A549 cells. (B) Overexpression of BRD7 suppressed the Myc mRNA expression in the A549 cells. (C) The mRNA expression level of Cyclin D was measured by using qRT-PCR in the A549 cells. (D) The mRNA expression level of Myc was measured by using qRT-PCR in the A549 cells.

doi:10.1371/journal.pone.0156701.g006

phosphorylation. Overexpression of BRD7 inhibited cyclin D and myc expression. Our findings demonstrated that BRD7 played a tumor suppressor role in lung adenocarcinoma tumorigenesis.

BRD7 is a member of the bromodomain family and is encoded in a locus on chromosome 16q12 [16, 29–31]. Recent studies have demonstrated that BRD7 acts as a tumor suppressor role in various cancers such as breast cancer, nasopharyngeal carcinoma, prostate cancer and osteosarcoma [19, 20, 22, 32]. For example, Hu et al showed BRD7 was downregulated in osteosarcoma, and knockdown of BRD7 increased colony formation, tumor growth and cell proliferation of osteosarcoma. [22]. BRD7 expression was correlated with patients' survival time in osteosarcoma tissue. Park et al. demonstrated that overexpression of BRD7 inhibited ovarian cancer cells invasion, apoptosis and viability [18]. Drost et al. showed that BRD7 inhibited tumorigenicity by acting as a p53 cofactor in breast cancer [29]. However, the expression of BRD7 in lung adenocarcinoma is still unknown. In our study, the protein and mRNA expression of BRD7 was downregulated in four lung adenocarcinoma cell lines (H1299, H23, SPC-A1 and A549) and one lung adenocarcinoma tissues compared to one adjacent no-tumor tissues. Tissues from lymph node metastases expressed lower levels of BRD7 compared to primary lung adenocarcinoma tissues and the adjacent normal tissue. When correlated to disease outcome, lower of BRD7 levels was associated with shortened disease-free survival in patients with lung adenocarcinoma.

We next studied the functional role of BRD7 in lung adenocarcinoma. BRD7 overexpression repressed lung adenocarcinoma cell proliferation and migration while knockdown of BRD7

expression promoted cell proliferation and migration. Previous study showed that BRD7 inhibited G1–S progression by transcriptionally regulating important molecules involved in ras/MEK/ERK and Rb/E2F pathways[33]. In this study, we also showed that inhibition of BRD7 could promote the ERK phosphorylation in the A549 cells. Moreover, knockdown of BRD7 expression increased the A549 cells proliferation and invasion, which could be blocked by U0126, a MEK inhibitor. These results demonstrated that downregulation of BRD7 promoted lung adenocarcinoma proliferation and migration through induction of ERK phosphorylation in the lung adenocarcinoma.

In conclusion, our results gave important clues for the functions of BRD7, indicating that BRD7 may present a promising candidate tumor suppressor gene in lung adenocarcinoma. Although the exact mechanism of BRD7 remains to be known, BRD7 may act as a potent target for therapeutic strategies for patients with lung adenocarcinoma.

Acknowledgments

We thank Bing Wang (Department of thoracic surgical oncology, cancer institute, China) for his assistance.

Author Contributions

Conceived and designed the experiments: YG BW SG.

Performed the experiments: YG BW SG.

Analyzed the data: YG BW SG.

Contributed reagents/materials/analysis tools: YG BW SG.

Wrote the paper: YG BW SG.

References

1. Tejero R, Navarro A, Campayo M, Vinolas N, Marrades RM, Cordeiro A, et al. miR-141 and miR-200c as markers of overall survival in early stage non-small cell lung cancer adenocarcinoma. *PLoS one*. 2014; 9(7):e101899. Epub 2014/07/09. doi: [10.1371/journal.pone.0101899](https://doi.org/10.1371/journal.pone.0101899) PMID: [25003366](https://pubmed.ncbi.nlm.nih.gov/25003366/); PubMed Central PMCID: PMC4087018.
2. Yang WB, Chen PH, Hsu Ts, Fu TF, Su WC, Liaw H, et al. Sp1-mediated microRNA-182 expression regulates lung cancer progression. *Oncotarget*. 2014; 5(3):740–53. Epub 2014/02/13. PMID: [24519909](https://pubmed.ncbi.nlm.nih.gov/24519909/); PubMed Central PMCID: PMC3996653.
3. Feng J, Xu L, Ni S, Gu J, Zhu H, Wang H, et al. Involvement of FoxQ1 in NSCLC through regulating EMT and increasing chemosensitivity. *Oncotarget*. 2014; 5(20):9689–702. Epub 2014/10/31. PMID: [25356753](https://pubmed.ncbi.nlm.nih.gov/25356753/); PubMed Central PMCID: PMC4259430.
4. Zhang ZY, Fu SL, Xu SQ, Zhou X, Liu XS, Xu YJ, et al. By downregulating Ku80, hsa-miR-526b suppresses non-small cell lung cancer. *Oncotarget*. 2015; 6(3):1462–77. Epub 2015/01/19. PMID: [25596743](https://pubmed.ncbi.nlm.nih.gov/25596743/).
5. Sun Y, Ai X, Shen S, Lu S. NF-kappaB-mediated miR-124 suppresses metastasis of non-small-cell lung cancer by targeting MYO10. *Oncotarget*. 2015. Epub 2015/03/10. PMID: [25749519](https://pubmed.ncbi.nlm.nih.gov/25749519/).
6. Song X, Shi K, Zhou SJ, Yu DP, Liu Z, Han Y. Clinicopathological significance and a potential drug target of RARBeta in non-small-cell lung carcinoma: a meta-analysis and a systematic review. *Drug design, development and therapy*. 2016; 10:1345–54. Epub 2016/04/23. doi: [10.2147/DDDT.S96766](https://doi.org/10.2147/DDDT.S96766) PMID: [27103788](https://pubmed.ncbi.nlm.nih.gov/27103788/); PubMed Central PMCID: PMC4827914.
7. Pal S, Amin PJ, Sainis KB, Shankar BS. Potential Role of TRAIL in Metastasis of Mutant KRAS Expressing Lung Adenocarcinoma. *Cancer microenvironment: official journal of the International Cancer Microenvironment Society*. 2016. Epub 2016/04/24. doi: [10.1007/s12307-016-0184-3](https://doi.org/10.1007/s12307-016-0184-3) PMID: [27106232](https://pubmed.ncbi.nlm.nih.gov/27106232/).

8. Yarchoan M, Lim M, Brahmer JR, Ettinger D. Oligometastatic Adenocarcinoma of the Lung: A Therapeutic Opportunity for Long-Term Survival. *Cureus*. 2015; 7(12):e409. Epub 2016/01/30. doi: [10.7759/cureus.409](https://doi.org/10.7759/cureus.409) PMID: [26824009](https://pubmed.ncbi.nlm.nih.gov/26824009/); PubMed Central PMCID: PMC4725613.
9. Xu P, Zhao M, Liu Z, Liu Y, Chen Y, Luo R, et al. Elevated nuclear CCND1 expression confers an unfavorable prognosis for early stage lung adenocarcinoma patients. *International journal of clinical and experimental pathology*. 2015; 8(12):15887–94. Epub 2016/02/18. PMID: [26884860](https://pubmed.ncbi.nlm.nih.gov/26884860/); PubMed Central PMCID: PMC4730073.
10. Maquilan G, Grover S, Xanthopoulos E, Evans TL, Aggarwal C, Langer CJ, et al. Analysis of the Relationship Between Response to Chemotherapy and Response to Radiation Therapy in Patients With Non-Small Cell Lung Cancer Receiving Sequential Treatment. *American journal of clinical oncology*. 2016. Epub 2016/04/22. doi: [10.1097/COC.0000000000000288](https://doi.org/10.1097/COC.0000000000000288) PMID: [27100960](https://pubmed.ncbi.nlm.nih.gov/27100960/).
11. Li D, He S. Pemetrexed and cyclophosphamide combination therapy for the treatment of non-small cell lung cancer. *International journal of clinical and experimental pathology*. 2015; 8(11):14693–700. Epub 2016/01/30. PMID: [26823793](https://pubmed.ncbi.nlm.nih.gov/26823793/); PubMed Central PMCID: PMC4713579.
12. Park YA, Lee JW, Choi JJ, Jeon HK, Cho Y, Choi C, et al. The interactions between MicroRNA-200c and BRD7 in endometrial carcinoma. *Gynecologic oncology*. 2012; 124(1):125–33. Epub 2011/10/22. doi: [10.1016/j.ygyno.2011.09.026](https://doi.org/10.1016/j.ygyno.2011.09.026) PMID: [22015043](https://pubmed.ncbi.nlm.nih.gov/22015043/).
13. Park SW, Herrema H, Salazar M, Cakir I, Cabi S, Basibuyuk Sahin F, et al. BRD7 regulates XBP1s' activity and glucose homeostasis through its interaction with the regulatory subunits of PI3K. *Cell metabolism*. 2014; 20(1):73–84. Epub 2014/05/20. doi: [10.1016/j.cmet.2014.04.006](https://doi.org/10.1016/j.cmet.2014.04.006) PMID: [24836559](https://pubmed.ncbi.nlm.nih.gov/24836559/); PubMed Central PMCID: PMC4079724.
14. Chiu YH, Lee JY, Cantley LC. BRD7, a tumor suppressor, interacts with p85alpha and regulates PI3K activity. *Molecular cell*. 2014; 54(1):193–202. Epub 2014/03/25. doi: [10.1016/j.molcel.2014.02.016](https://doi.org/10.1016/j.molcel.2014.02.016) PMID: [24657164](https://pubmed.ncbi.nlm.nih.gov/24657164/); PubMed Central PMCID: PMC4004185.
15. Liu H, Zhou M, Luo X, Zhang L, Niu Z, Peng C, et al. Transcriptional regulation of BRD7 expression by Sp1 and c-Myc. *BMC molecular biology*. 2008; 9:111. Epub 2008/12/30. doi: [10.1186/1471-2199-9-111](https://doi.org/10.1186/1471-2199-9-111) PMID: [19111069](https://pubmed.ncbi.nlm.nih.gov/19111069/); PubMed Central PMCID: PMC2632634.
16. Kaeser MD, Aslanian A, Dong MQ, Yates JR 3rd, Emerson. BRD7, a novel PBAF-specific SWI/SNF subunit, is required for target gene activation and repression in embryonic stem cells. *The Journal of biological chemistry*. 2008; 283(47):32254–63. Epub 2008/09/24. doi: [10.1074/jbc.M806061200](https://doi.org/10.1074/jbc.M806061200) PMID: [18809673](https://pubmed.ncbi.nlm.nih.gov/18809673/); PubMed Central PMCID: PMC2583284.
17. Liu H, Li X, Niu Z, Zhang L, Zhou M, Huang H, et al. Preparation of polyclonal antibody specific for BRD7 and detection of its expression pattern in the human fetus. *The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society*. 2008; 56(6):531–8. Epub 2007/12/12. doi: [10.1369/jhc.7A7340.2007](https://doi.org/10.1369/jhc.7A7340.2007) PMID: [18071067](https://pubmed.ncbi.nlm.nih.gov/18071067/); PubMed Central PMCID: PMC2386764.
18. Park YA, Lee JW, Kim HS, Lee YY, Kim TJ, Choi CH, et al. Tumor suppressive effects of bromodomain-containing protein 7 (BRD7) in epithelial ovarian carcinoma. *Clinical cancer research: an official journal of the American Association for Cancer Research*. 2014; 20(3):565–75. Epub 2013/11/08. doi: [10.1158/1078-0432.CCR-13-1271](https://doi.org/10.1158/1078-0432.CCR-13-1271) PMID: [24198243](https://pubmed.ncbi.nlm.nih.gov/24198243/).
19. Pern F, Bogdanova N, Schurmann P, Lin M, Ay A, Langer F, et al. Mutation analysis of BRCA1, BRCA2, PALB2 and BRD7 in a hospital-based series of German patients with triple-negative breast cancer. *PloS one*. 2012; 7(10):e47993. Epub 2012/10/31. doi: [10.1371/journal.pone.0047993](https://doi.org/10.1371/journal.pone.0047993) PMID: [23110154](https://pubmed.ncbi.nlm.nih.gov/23110154/); PubMed Central PMCID: PMC3480465.
20. Liu H, Zhang L, Niu Z, Zhou M, Peng C, Li X, et al. Promoter methylation inhibits BRD7 expression in human nasopharyngeal carcinoma cells. *BMC cancer*. 2008; 8:253. Epub 2008/09/10. doi: [10.1186/1471-2407-8-253](https://doi.org/10.1186/1471-2407-8-253) PMID: [18778484](https://pubmed.ncbi.nlm.nih.gov/18778484/); PubMed Central PMCID: PMC2543047.
21. Wu WJ, Hu KS, Chen DL, Zeng ZL, Luo HY, Wang F, et al. Prognostic relevance of BRD7 expression in colorectal carcinoma. *European journal of clinical investigation*. 2013; 43(2):131–40. Epub 2012/12/12. doi: [10.1111/eci.12024](https://doi.org/10.1111/eci.12024) PMID: [23215825](https://pubmed.ncbi.nlm.nih.gov/23215825/).
22. Hu K, Liao D, Wu W, Han AJ, Shi HJ, Wang F, et al. Targeting the anaphase-promoting complex/cyclosome (APC/C)- bromodomain containing 7 (BRD7) pathway for human osteosarcoma. *Oncotarget*. 2014; 5(10):3088–100. Epub 2013/03/21. PMID: [24840027](https://pubmed.ncbi.nlm.nih.gov/24840027/); PubMed Central PMCID: PMC4102794.
23. Akagi I, Okayama H, Schetter AJ, Robles AI, Kohno T, Bowman ED, et al. Combination of protein coding and noncoding gene expression as a robust prognostic classifier in stage I lung adenocarcinoma. *Cancer research*. 2013; 73(13):3821–32. Epub 2013/05/04. doi: [10.1158/0008-5472.CAN-13-0031](https://doi.org/10.1158/0008-5472.CAN-13-0031) PMID: [23639940](https://pubmed.ncbi.nlm.nih.gov/23639940/).
24. Ho CS, Yap SH, Phuah NH, In LL, Hasima N. MicroRNAs associated with tumour migration, invasion and angiogenic properties in A549 and SK-Lu1 human lung adenocarcinoma cells. *Lung Cancer*. 2014; 83(2):154–62. Epub 2013/12/24. doi: [10.1016/j.lungcan.2013.11.024](https://doi.org/10.1016/j.lungcan.2013.11.024) PMID: [24360396](https://pubmed.ncbi.nlm.nih.gov/24360396/).

25. Yan G, Yao R, Tang D, Qiu T, Shen Y, Jiao W, et al. Prognostic significance of microRNA expression in completely resected lung adenocarcinoma and the associated response to erlotinib. *Med Oncol*. 2014; 31(10):203. Epub 2014/09/07. doi: [10.1007/s12032-014-0203-5](https://doi.org/10.1007/s12032-014-0203-5) PMID: [25192889](https://pubmed.ncbi.nlm.nih.gov/25192889/).
26. Ge X, Zheng L, Huang M, Wang Y, Bi F. MicroRNA expression profiles associated with acquired gefitinib-resistance in human lung adenocarcinoma cells. *Molecular medicine reports*. 2015; 11(1):333–40. Epub 2014/10/24. doi: [10.3892/mmr.2014.2757](https://doi.org/10.3892/mmr.2014.2757) PMID: [25339453](https://pubmed.ncbi.nlm.nih.gov/25339453/).
27. Schliekelman MJ, Taguchi A, Zhu J, Dai X, Rodriguez J, Celiktas M, et al. Molecular portraits of epithelial, mesenchymal, and hybrid States in lung adenocarcinoma and their relevance to survival. *Cancer research*. 2015; 75(9):1789–800. Epub 2015/03/07. doi: [10.1158/0008-5472.CAN-14-2535](https://doi.org/10.1158/0008-5472.CAN-14-2535) PMID: [25744723](https://pubmed.ncbi.nlm.nih.gov/25744723/).
28. Robles AI, Arai E, Mathe EA, Okayama H, Schetter AJ, Brown D, et al. An Integrated Prognostic Classifier for Stage I Lung Adenocarcinoma Based on mRNA, microRNA, and DNA Methylation Biomarkers. *Journal of thoracic oncology: official publication of the International Association for the Study of Lung Cancer*. 2015; 10(7):1037–48. Epub 2015/07/03. doi: [10.1097/JTO.0000000000000560](https://doi.org/10.1097/JTO.0000000000000560) PMID: [26134223](https://pubmed.ncbi.nlm.nih.gov/26134223/).
29. Drost J, Mantovani F, Tocco F, Elkon R, Comel A, Holstege H, et al. BRD7 is a candidate tumour suppressor gene required for p53 function. *Nature cell biology*. 2010; 12(4):380–9. Epub 2010/03/17. doi: [10.1038/ncb2038](https://doi.org/10.1038/ncb2038) PMID: [20228809](https://pubmed.ncbi.nlm.nih.gov/20228809/).
30. Van de Vijver P, Scheer L, van Beijnum J, Griffioen A, Hackeng TM. Application of an omonasteine ligation strategy for the total chemical synthesis of the BRD7 bromodomain. *Chem Commun (Camb)*. 2012; 48(75):9403–5. Epub 2012/08/15. doi: [10.1039/c2cc34956f](https://doi.org/10.1039/c2cc34956f) PMID: [22889936](https://pubmed.ncbi.nlm.nih.gov/22889936/).
31. Sun H, Liu J, Zhang J, Shen W, Huang H, Xu C, et al. Solution structure of BRD7 bromodomain and its interaction with acetylated peptides from histone H3 and H4. *Biochemical and biophysical research communications*. 2007; 358(2):435–41. Epub 2007/05/15. doi: [10.1016/j.bbrc.2007.04.139](https://doi.org/10.1016/j.bbrc.2007.04.139) PMID: [17498659](https://pubmed.ncbi.nlm.nih.gov/17498659/).
32. Kikuchi M, Okumura F, Tsukiyama T, Watanabe M, Miyajima N, Tanaka J, et al. TRIM24 mediates ligand-dependent activation of androgen receptor and is repressed by a bromodomain-containing protein, BRD7, in prostate cancer cells. *Biochimica et biophysica acta*. 2009; 1793(12):1828–36. Epub 2009/11/17. doi: [10.1016/j.bbamcr.2009.11.001](https://doi.org/10.1016/j.bbamcr.2009.11.001) PMID: [19909775](https://pubmed.ncbi.nlm.nih.gov/19909775/).
33. Zhou J, Ma J, Zhang BC, Li XL, Shen SR, Zhu SG, et al. BRD7, a novel bromodomain gene, inhibits G1-S progression by transcriptionally regulating some important molecules involved in ras/MEK/ERK and Rb/E2F pathways. *Journal of cellular physiology*. 2004; 200(1):89–98. Epub 2004/05/12. doi: [10.1002/jcp.20013](https://doi.org/10.1002/jcp.20013) PMID: [15137061](https://pubmed.ncbi.nlm.nih.gov/15137061/).