

RESEARCH ARTICLE

Cloning and Expression of β -Defensin from Soiny Mullet (*Liza haematocheila*), with Insights of its Antibacterial Mechanism

Zhitao Qi^{1,2}, Wei Xu¹, Fancui Meng³, Qihuan Zhang¹, Chenglung Chen^{4*}, Rong Shao^{1*}

1 Key Laboratory of Biochemistry and Biotechnology of Marine Wetland of Jiangsu Province, Yancheng Institute of Technology, Yancheng, 224051, Jiangsu, China, **2** Key Laboratory of Aquaculture and Ecology of Coastal Pool in Jiangsu Province, Yancheng Institute of Technology, Yancheng, 224051, Jiangsu, China, **3** Tianjin Institute of Pharmaceutical Research, Tianjin, 300193, China, **4** Department of Chemistry, National Sun Yat-Sen University, Kaohsiung, 80424, Taiwan, ROC

* chen1@mail.nsysu.edu.tw (CC); sr@163.com (RS)



OPEN ACCESS

Citation: Qi Z, Xu W, Meng F, Zhang Q, Chen C, Shao R (2016) Cloning and Expression of β -Defensin from Soiny Mullet (*Liza haematocheila*), with Insights of its Antibacterial Mechanism. PLoS ONE 11(6): e0157544. doi:10.1371/journal.pone.0157544

Editor: Zhan Yin, Chinese Academy of Sciences, CHINA

Received: January 18, 2016

Accepted: June 1, 2016

Published: June 20, 2016

Copyright: © 2016 Qi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information file.

Funding: This work was supported by the Major Projects of Natural Science Research in Universities and Colleges in Jiangsu Province (Grant no. 15KJA240001) and partially by the National Natural Science Foundation of China (Grant no. 31302221 and 31272666) and Jiangsu Province (Grant no. BK2011418 and BK20151297). ZQ was financially supported by the "Qinglan" project of Jiangsu province.

Abstract

Beta-defensins are important part of innate immunity of fish, which are the first defense line against invading pathogens. In this study, the β -defensin (Lh β -defensin) gene was cloned from spleen tissue of soiny mullet (*Liza haematocheila*). Lh β -defensin cDNA was 747 bp in length, encoding 63 amino acids. Sequence alignment revealed that Lh β -defensin contained six conserved cysteine residues and shared 97.5% sequence identities with grouper (*Epinephelus coioides*) β -defensin. Realtime PCR revealed that Lh β -defensin was highest expressed in the immune related organs, such as spleen, kidney and gut of healthy fish. Following *Streptococcus dysgalactiae* infection, Lh β -defensin was up-regulated in immune related organs, e.g. 17.6-fold in spleen and 10.87-fold in gut at 24 h post infection (hpi). Lh β -defensin possessed a monomeric structure of a three-stranded anti-parallel β -sheet and an α -helix stabilized by three disulfide bonds formed by Cys30-Cys58, Cys36-Cys52, and Cys40-Cys59. In addition to the experimental work, computer simulation was also carried out to determine the possible conformation of β -defensin and its interaction with palmitoyl-oleoylphosphatidylglycerol (POPG), a model of bacteria membrane. The Lh β -defensin was found to form dimeric structure stabilized by the van der Waals contacts of Leu35 and Cys37 in two anti-parallel β 1-strands and the cation- π interaction between Tyr32 and Arg54 respectively in the two β 1-strands. The most important interactions between β -defensin and membrane are the electrostatic interactions between Arg residues in β -defensin and head group of POPG bilayer as well as hydrogen bond interactions between them. Our results were useful for further understanding the potential mechanism of antimicrobial property of fish β -defensins.

Competing Interests: The authors have declared that no competing interests exist.

Introduction

Antimicrobial peptides (AMPs) are important components of the innate immunity acting as the first defense line against invading pathogens. The defensins are cysteine-rich cationic antimicrobial peptides which are divided into three distinct groups, α -, β - and θ -defensins, based on the different disulfide bond [1,2]. α -defensins were isolated from mammalian neutrophils and macrophages which are stabilized by a mode of disulfide bonds of C1-C6, C2-C4 and C3-C5. θ -defensins are the macrocyclic product of a head-to-tail of two truncated α -defensins which are only found in leukocytes and bone marrows of Old World monkeys [2, 3]. β -defensins are the largest group of defensins family and are the most characteristic defensins. β -defensins are comprised of 35 to 50 amino acids with a core of three anti-parallel β -strands stabilized by a mode of disulfide bond of C1-C5, C2-C4 and C3-C6 [4].

β -defensins had been characterized in a number of fish species, such as zebrafish (*Danio rerio*), fugu (*Takifugu rubripes*), tetraodon (*Tetraodon nigroviridis*) [5], blunt snout bream (*Megalobrama amblycephala*) [6] and Nile tilapia (*Oreochromis niloticus*) [7]. Fish β -defensins exhibited similar functions as mammalian β -defensins, including antimicrobial activity against bacterial [8, 9] and viruses [10], chemotactic attractants of immune cells [11] and modulator of immune response [12, 13].

Many studies had been carried out to elucidate the mechanism of action of various mammalian AMPs. The positively charged surface of defensins played a key role in their actions, which enabled them to initially attach to the bacterial membrane, destabilize and disrupt the cell membrane and form pores on membrane resulting in the leakage of the cell contents [14–16]. However, the diversity of sequences and structures of AMPs revealed that these molecules might act in diverse ways [16]. And so far, there were less studies on the mechanism of the fish AMPs. Thus, the present study described the successful cloning and characterization of β -defensin from soiny mullet (*Liza haematocheila*), an economically important aquaculture mugilid species in China and other Asian countries [17]. Since it is very difficult to determine experimentally the structure of Lh β -defensin, computational method was adopted to probe into monomeric and dimeric structure of Lh β -defensin by using protein docking and followed by molecular dynamics (MD) simulation. The interaction of Lh β -defensin dimer and bacterial membrane (POPG lipid membrane) was also investigated base on the MD simulation.

Materials and Methods

This study was carried out in strict accordance with the approval of the Academic committee of Yancheng Institute of Technology (No. 2015016). The fish were euthanized by using ethyl-3 aminobenzoate methanesulfonate (MS-222, Sigma) according to the Measures for the Administration of Experimental Animals in Jiangsu province (No. 2008[45]). This study did not involve endangered or protected species. The *Streptococcus dysgalactiae*, a kind of gram positive bacterial, was kindly provided Professor Li Aihua from the State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, the Chinese Academy of Sciences.

Fish

Healthy soiny mullet with an average weight of 5.0 ± 0.5 g were purchased from Dafuyuan mill of fishery (Sheyang town, Yancheng city, China) and transported to our laboratory by an oxygen-supplying car. The fish were reared in 120 L plastic aquaria supplied with oxygen and fed with commercial diet twice daily. The acclimation lasted for 2 weeks prior to experiments.

Molecular cloning of Lhβ-defensin gene

Total RNA was extracted from spleen tissue using TRIzol reagent according to manufacturer's instructions. After the quality verification of RNA by electrophoresis and spectrophotometry, RNA was reverse transcribed into first-strand cDNA using First Strand cDNA Synthesis Kit (Fermentas, USA) according to manufacturer's constructions.

Degenerate primers (LhBD-dF1 and LhBD-dR1) were designed based on the conserved regions of teleost β-defensin genes (Table 1). PCR was carried out using degenerate primers with spleen cDNA as template to obtain the partial sequence of Lhβ-defensin. The PCR products were ligated into pMD18-T vector (TaKaRa, Japan) and then transformed into *E. coli* DH5α competent cells. The selected positive clones were used for subsequent sequencing (Sangon Biotechnology Company, Shanghai, China). After the sequence was suggested to be the partial sequence of β-defensin gene by BLASTx analysis, specific primers were designed to obtain the full length cDNA sequence of Lhβ-defensin gene by 5' and 3' rapid amplification of cDNA ends (RACE) PCR method. The primers used in this study were listed in Table 1.

Sequence analysis

The amino acid (aa) sequence was deduced by Translate software in the ExPasy website (<http://www.expasy.org>). The alignment of amino acid sequences was carried out using ClustalO software (<http://www.ebi.ac.uk/tools/msa/clustalo/>), and decorated with BoxShade software (http://www.ch.embnet.org/software/BOX_form.html). The amino acids sequence identities were analyzed using MegAlign software in the DNASTar package. The phylogenetic tree was constructed by MEGA 4.1 software using neighbor-joining algorithm in which the bootstrap was set as 10,000 replicates to measure the confidence of branch topology [18].

Tissue distribution of Lhβ-defensin transcripts

Four tissues including liver, spleen, kidney and gut were sampled from six healthy fish, respectively. RNA extraction with TRIzol reagent, cDNA synthesis and realtime quantitative PCR were carried out as described previously [19]. The expression level of Lhβ-defensin was normalized to that of β-actin. Primers used for real-time quantitative PCR were listed in Table 1.

Table 1. Primers used for gene cloning and expression analysis.

Primer	Sequence (5' to 3')	Application
LhBD-dF1	ATGAAGGGACTGAGCTTGGTTC	RT-PCR
LhBD-dR1	GTGATG (CA) CCAACG (AG) TGTACTCCTG	RT-PCR
LhBD-5out	ACCCACAGGTCCAATACTGCAT	5'RACE PCR
LhBD-5in	AGAACCGTCTGCAGAGTCCTCA	5'RACE PCR
LhBD-3out	ATGCAGTATTGGACCTGTGGGT	3'RACE PCR
LhBD-3in	AGAGGACTCTGCAGACGGTT	3'RACE PCR
LhBD-F	CAGTATTGGACCTGTGGG	Realtime PCR
LhBD-R	CTAAGACCGCACCGCACA	Realtime PCR
Lhactin-F	CAGCCATACTGTGCCCATCT	Realtime PCR
Lhactin-R	TCCTTGATGTCACGCACGAT	Realtime PCR

doi:10.1371/journal.pone.0157544.t001

Expression of Lh β -defensin following *Streptococcus dysgalactiae* infection

The bacterial infection experiments were carried out on the basis of our previous studies [17, 19]. Briefly, twenty fish were randomly divided into two groups including control group in which fish were infected intraperitoneally (i.p) with 100 μ L PBS, and bacterial infection group in which fish were injected with 100 μ L live *S. dysgalactiae*, (2×10^6 CFU). The fish were monitored every two hours and no dead fish was observed during the whole experiment. The tissues including liver, spleen, kidney and gut from nine fish per group were sampled respectively at 24 h post infection (hpi). Real-time PCR was done as described previously and expressed as fold change relative to the time-matched controls [19].

Homology modeling and MD analysis of Lh β -defensin monomer and dimer

The monomeric structure of Lh β -defensin was constructed by homology modeling method using Prime module of Schrödinger software [20] with the available human β -defensin-1 (HBD-1) structure (PDB ID: 1IJV) as template [21]. Afterward, the dimeric structures were generated using SymmDock server [22, 23]. The generated 100 structures were clustered according to RMSD (root mean square deviation) values and four typical dimeric structures were obtained. Then, molecular dynamics simulations of the four typical dimers were performed using GROMACS 5.1 software [24] and AMBER99SB force field [25]. Briefly, the initial dimeric structure was placed in a box with an edge of 1.2 nm and solvated with TIP3P (three-point transferable intermolecular potential) water molecules. Chloride ions were added to neutralize the system. After energy minimization, 100 ps NVT and 100 ps NPT MD simulations were performed with position restrictions on peptides. Finally, MD simulation was carried out for 20 ns with a time step of 1 fs and trajectory was saved every 20 ps for analysis. The V-rescale method was used for temperature coupling with the reference temperature 300 K. The pressure was controlled isotropically using Parrinello-Rahman coupling method. Particle mesh Ewald (PME) method was used to consider the long-range electrostatic interaction. The cut-off distances for electrostatic and van der Waals interactions were both set to be 1.4 nm.

Interaction between Lh β -defensin dimer and POPG bilayer

The palmitoylphosphatidylglycerol (POPG) bilayer was constructed to assemble the bacterial membrane. Both the coordinate and the topology parameters for POPG bilayer were obtained from lipidbook (<http://lipidbook.bioch.ox.ac.uk>) [26]. This lipid bilayer includes 128 POPG molecules and has been tested in previous reported calculations [27]. The calculated order parameters, area per lipid and thickness of this bilayer with the AMBER force field were in excellent agreement with experiments [28, 29]. MD simulation was carried out to investigate the interaction between Lh β -defensin and the membrane. The dimeric structure of Lh β -defensin was placed in a random orientation about 5 nm away from the POPG bilayer center. Then, the system was solvated with roughly 10,000 water molecules and counter ions were added to neutralize the system. MD simulation was performed on this system for 100 ns. The reference temperature was set at 323 K, which is higher than the phase transition temperature of POPG bilayer. All the calculations were carried out using GROMACS package 5.1 [24] with AMBER99SB force field [25].

Results and Discussion

Sequence features of Lhβ-defensin

The cloned Lhβ-defensin cDNA was 747 bp in length (GenBank accession no. KJ872680), including 5'-untranslated region (UTR) of 215 bp, open reading frame (ORF) of 192 bp, and 3'-UTR of 340 bp. There was one polyadenylation signal (AATAAA) upstream of the poly (A) tail (S1 Fig).

The fish β-defensins ranged from 62 to 68 amino acids in length and contained a signal peptide of 18–26 amino acids [30]. In the present study, the ORF of Lhβ-defensin cDNA encoded 63 amino acids, which shared similar length with known fish β-defensin pro-peptide. A signal peptide of 20 amino acids was also predicted from Lhβ-defensin using SignalP software. Thus, the mature peptide of Lhβ-defensin contains 43 amino acids. The molecular weight of mature peptide was 5.18 kD and the isoelectric point (pI) 8.92, indicating that the net cationic feature of Lhβ-defensin. Further sequence alignment revealed that Lhβ-defensin shared 22.5–97.5% sequence identities with teleost β-defensins, among which highest sequence identities was with grouper (*Epinephelus coioides*) β-defensin (97.5%) and lowest with defensin-4 of Japanese flounder (*Paralichthys olivaceus*) (22.5%). Six conserved cysteine residues were found to exist in Lhβ-defensin (Fig 1), which were well conserved in insects, fish, chicken and human β-defensins, and were crucial for structural stability and functions of β-defensin [31].

Phylogenetic tree analysis

A neighbor-joining phylogenetic tree was constructed with the reported β-defensin amino acids sequences of vertebrates using MEGA4.1 software. Results revealed that the phylogenetic




Fig 1. Multiple alignment of vertebrates β-defensins. C1-C6 indicated the six conserved cysteine residues. Black shade indicated identical amino acids and gray shade indicated similar amino acids.

doi:10.1371/journal.pone.0157544.g001

tree was divided into two main clusters. One cluster contained mammalian and birds β -defensins and another one contained the fish β -defensins. The cluster of fish β -defensins was further divided into two sub-clusters, one sub-cluster including all members of fish β -defensin1, and another sub-cluster containing fish β -defensin 2 and 3. The Lh β -defensin was placed well in the sub-cluster of fish β -defensin 2 and 3. Similar results were also observed in the cluster of mammalian β -defensins (Fig 2). β -defensins in fish had expanded and diverged into two separated types. These two separated types of β -defensins may have diverged before the appearance of fish species and arisen from a common ancestor by gene or genome duplication events [5]. Multiple and divergent β -defensins may help animals to better defense against the diverse microbial pathogens in their habitats [32].

Expression pattern of β -defensin in tissues from healthy and *S. dysgalactiae* infected mullets

Mammalian β -defensins were mainly expressed in the epithelia of many organs while fish β -defensins were mainly expressed in the immune and mucosal tissues. However, expression patterns of fish β -defensins were different in various fish species, e.g mandarin fish β -defensin was highly expressed in the spleen, intestine, gill and head-kidney [9], and Wuchang bream β -defensin1 was highly expressed in skin, blood and immune related organs [6], while Nile tilapia β -defensin was highly in skin, spleen, kidney and muscle [7]. In the present study, Lh β -defensin was highest expressed in spleen, followed by the kidney and gut, and lowest in liver (Fig 3A). Thus, it was suggested that fish β -defensins may be species-specific and play a broad role in different fish tissues.

The expression of fish β -defensins was modulated by microbial pathogens infections. Following *Aeromonas sobria* infection, the expression of β -defensin was up-regulated in the skin of Wuchang bream [6]. Rainbow trout β -defensins were also up-regulated in the intestine and gill at 48 h post of *Yersinia ruckeri* infection [33]. In addition, Nile tilapia β -defensin exerted different up-regulated pattern at different time points post *S.agalactiae* infection, e.g. 3.3-fold in skin at 6 hpi, 57.1-fold in muscle at 24 hpi and 1.6-fold in gill at 72 hpi [7]. Similar to previous reports, Lh β -defensin was also up-regulated following *S. dysgalactiae* at 24 hpi, e.g. 17.6-fold in spleen, 10.87-fold in gut and 1.23-fold in liver (Fig 3B). The up-regulated β -defensin may take part in the immune response against invading pathogens.

Monomeric and dimeric structure of Lh β -defensin

The monomer of Lh β -defensin shared a similar structure with human β -defensin2 [34], possessing a three-stranded anti-parallel β -sheet and an α -helix stabilized by three disulfide bonds formed by Cys30-Cys58, Cys36-Cys52, and Cys40-Cys59 (Fig 4A). The mode of disulfide bonds in this monomer was perfectly matched the mode of disulfide bonds of β -defensin family members, which confirmed the gene we cloned was exactly the homologue of β -defensin. The results of Ramachandran angle analysis of this monomer showed that most of residues appeared in the energetic favored and allowed regions, and only three residues occurred in the disallowed region (Fig 4B). This indicated that the constructed structure was reliable.

Further, the dimeric structures of Lh β -defensin were constructed using SymmDock method. Four typical dimers of Lh β -defensin were obtained after clustering by RMSD analysis. Dimers A and B were formed by parallel and anti-parallel β 1- β 1 strands interactions while dimers C and D were created through parallel and anti-parallel β 2- β 2 strands interactions, respectively (S2 Fig). Then, the stability of each dimeric structure was analyzed by equilibrium MD simulation and solvent accessible surface areas analysis using GROMACS 5.0 software. Dimer C had the largest RMSD value among the four dimers and its RMSD fluctuated widely compared




Fig 2. Phylogenetic tree of vertebrates β-defensins. The tree was constructed by MEGA 4.1 software using neighbor-joining method in which the bootstrap was set as 10,000 replicates to measure the confidence of branch topology. The Genbank accession number of sequences used for the tree was listed behind the common name of species. The newly obtained Lhβ-defensin was marked by round black circle in the tree.

doi:10.1371/journal.pone.0157544.g002




Fig 3. Tissues expression of Lhβ-defensin in tissues of healthy and *S. dysgalactiae* infected fish. (A) Expression of Lhβ-defensin in healthy fish was measured by realtime PCR and normalized to β-actin. (B) Expression of Lhβ-defensin in fish infected with *S. dysgalactiae*. Fish in the bacterial infected group were infected intraperitoneally with 2×10^6 CFU live *S. dysgalactiae* and the control group was injected with the same amount of PBS solution. At 24 hpi, the tissues were selected for the gene expression analysis by Realtime PCR method. Asterisks indicate that the expression of Lhβ-defensin in the infected group was significantly up-regulated compared with that of control group (* $P < 0.05$).

doi:10.1371/journal.pone.0157544.g003

with other dimers, revealing that the structure of dimer C deviated seriously from its initial structure and did not reach equilibrated state after 20 ns MD simulation. The overall RMSD of dimer B was about 0.3 nm with moderate fluctuation after 10 ns simulation, suggesting that it was well equilibrated during MD analysis. The RMSD profiles of dimer A and D were similar and their RMSD values ranged in the middle of that of dimers B and C (S3 Fig). Further solvent accessible surface areas revealed that dimer B had the largest buried surface and oscillated at 12 nm², while the buried surface areas of the other three dimers fluctuated at 10 nm². Significant fluctuation was observed for Dimer C, revealed large structure alteration occurred during the simulation (S4 Fig). Thus, it was reasonable to choose the dimer B for further analysis. The interface of dimer B was formed through van der Waals contacts of Leu35 and Cys37 in two




Fig 4. Predicted monomeric structure (A) and Ramachandran plot (B) of Lh β -defensin.

doi:10.1371/journal.pone.0157544.g004




Fig 5. Reduced density gradient isosurface and residue type of Lh β -defensin dimer. (A) Reduced density gradient isosurface map between Arg54 and Tyr32 in dimeric structure of Lh β -defensin created by Multiwfn software [36]. The blue, red and green (or earth green) colors indicate the strong attractive, strong repulsive and van der Waals interaction, respectively. (B) Surface of dimer colored by residue type. Blue are basic residues, red are acidic residues, green are polar residues, and white are non-polar residues.

doi:10.1371/journal.pone.0157544.g005

anti-parallel β 1-strands. The overlap of β 1 strands was stabilized by the cation- π interactions between Tyr32 and Arg54 respectively in the two β 1-strands (Fig 5A and 5B). Tyr56 also participated to create cation- π interactions with Arg54. Cation- π interaction is a non-covalent molecular interaction between the electron-rich π system and a positively charged species. The hydrophobic residues create a large surface of dimer B, including residues Tyr27, Cys30, Gly31 and Tyr 32 in α -helix, Gly34, Leu35 and Cys36 in β 1 strand, and Pro53 in the loop between β 2 and β 3 strands (Fig 5B). The buried hydrophobic surface increased the positively charged surface during formation of dimer B, which play a crucial role in stabilizing protein structure and may enhance the interaction of β -defensin with negatively charged membrane [35].

Insights of antibacterial mechanism of Lh β -defensin

To reveal the potential antibacterial mechanism of Lh β -defensin, the structure order of acyl chains of POPG membrane, one of model of bacterial membrane [37], with or without Lh β -defensin dimer were analyzed [38]. There was a decreasing in the order parameters of atom 10 in the absence of Lh β -defensin dimer, which might be caused by the presence of the double bond in the sn2 chain of POPG. Similar results were also observed in previous studies [16, 27]. Compared with pure POPG, the order parameters of POPG decreased greatly in the presence of Lh β -defensin dimer, especially when the POPG was around the 10 Å of defensin dimer, suggesting that large deformation of lipid structure may occur around the defensin contact regions (Fig 6). This deformation of POPG leads to water permeating inside the hydrophobic region of the bilayer. The initial contact of defensin on the surface of POPG bilayer caused expansion of membrane and hence reducing the packing density of acyl hydrocarbon chains in tail region. This resulted in water translocation in the contact region and destructed the membrane (Fig 7A). Meanwhile, the arginines of Lh β -defensin also played important roles in the interaction between β -defensin and POPG bilayer, which led to strong electrostatic interactions and formed large number of hydrogen bonds with the anionic surface of POPG bilayer (Fig 7B). Thus, we speculated that the electrostatic and hydrogen bond interactions between




Fig 6. Order parameters (S_{CD}) of saturated (sn-1) and unsaturated (sn-2) hydrocarbon chains in POPG computed using the last 20 ns trajectories of stimulation. Sn-1 and sn-2 referred to S_{CD} calculated by POPG in the presence of Lh β -defensin dimer, sn1-local and sn2-local to S_{CD} calculated when POPG was around 10 Å of defensin dimer, while sn1-pure POPG and sn2-pure POPG to S_{CD} calculated by POPG in the absence of Lh β -defensin dimer.

doi:10.1371/journal.pone.0157544.g006




Fig 7. Snapshot showing the water defects and water translocations across the membrane (A); Interactions between arginines of Lh β -defensin and POPG bilayer (B).

doi:10.1371/journal.pone.0157544.g007

Lh β -defensin and membranes played important roles in the microbicidal activity of β -defensin. These results were in line with studies on human β -defensin [34, 39, 40], indicating a conserved antibacterial mechanism may exist both in fish and mammalian β -defensin.

In conclusion, a β -defensin gene was successfully cloned from soiny mullet (*Liza haematocheila*) and its expression patterns in tissues from healthy and *S. dysgalactiae* infected fish were investigated. The Lh β -defensin was highly expressed in the immune-related organs of soiny mullet and up-regulated by *S. dysgalactiae* infection, suggesting its important role in the immune defense against bacterial. Lh β -defensin possessed conserved monomeric structure and formed dimeric structure which was mainly stabilized by the van der Waals contacts of Leu35 and Cys37 in two anti-parallel β 1-strands and the cation- π interaction between Tyr32 and Arg54 respectively in the two β 1-strands. Electrostatic interactions between Arg residues and head group of POPG play important roles in the interaction of β -defensin with membrane. Hydrogen bond interactions played an important role as well. To our best of knowledge, this is the first study to analyze the structure and potential antimicrobial mechanism of fish β -defensin. Our results were useful for further studying the mechanism of antimicrobial property of fish β -defensin.

Supporting Information

S1 Fig. The cDNA and deduced amino acids of Lh β -defensin. The signal peptide was marked by black line. The start codon (ATG) and stop codon (TAG) was boxed. The polyadenylation signal was marked by double black line.

(DOC)

S2 Fig. Final snapshot of four dimers after 20 ns MD simulation.

(DOC)

S3 Fig. RMSD analysis of four dimers of Lh β -defensin. RMSD of dimer A was marked by black line, dimer B by red line, dimer C by blue line and dimer D by green line.

(DOC)

S4 Fig. Buried surface of four dimers of Lh β -defensin. Buried surface area of dimer A was marked by black line, dimer B by red line, dimer C by blue line and dimer D by green line. (DOC)

Acknowledgments

We thank Pro. Li Aihua, the State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, the Chinese Academy of Sciences, for kindly providing *S. dysgalactiae*, and Dr. Zou Jun, Scottish Fish Immunology Research Center, University of Aberdeen for his suggestions on primers design.

Author Contributions

Conceived and designed the experiments: ZQ WX CC RS. Performed the experiments: ZQ WX FM. Analyzed the data: ZQ WX QZ FM CC. Contributed reagents/materials/analysis tools: FM CC. Wrote the paper: ZQ FM CC RS.

References

1. Ganz T (2003) Defensins: antimicrobial peptides of innate immunity. *Nat Rev Immunol.* 3: 828–838.
2. Zhu S, Gao B (2013) Evolutionary origin of beta-defensins. *Dev Comp Immunol.* 39: 79–84. doi: [10.1016/j.dci.2012.02.011](https://doi.org/10.1016/j.dci.2012.02.011) PMID: [22369779](https://pubmed.ncbi.nlm.nih.gov/22369779/)
3. Penberthy WT, Chari S, Cole AL, Cole AM (2011) Retrocyclins and their activity against HIV-1. *Cell Mol Life Sci.* 68: 2231–2242. doi: [10.1007/s00018-011-0715-5](https://doi.org/10.1007/s00018-011-0715-5) PMID: [21553001](https://pubmed.ncbi.nlm.nih.gov/21553001/)
4. Sawai MV, Jia HP, Liu L, Aseyev V, Wienczek JM, McCray PB Jr, et al. (2001) The NMR structure of human β -defensin-2 reveals a novel alpha-helical segment. *Biochemistry.* 40: 3810. PMID: [11300761](https://pubmed.ncbi.nlm.nih.gov/11300761/)
5. Zou J, Mercier C, Koussounadis A, Secombes CJ (2007) Discovery of multiple β -defensin like homologues in teleost. *Mol Immunol.* 44: 638–647. PMID: [16540171](https://pubmed.ncbi.nlm.nih.gov/16540171/)
6. Liang T, Wang DD, Zhang GR, Wei KJ, Wang WM, Zou GW (2013) Molecular cloning and expression analysis of two β -defensin genes in the blunt snout bream (*Megalobrama amblycephala*). *Comp. Biochem. Physiol. B Biochem. Mol Biol.* 166: 91–98. doi: [10.1016/j.cbpb.2013.07.006](https://doi.org/10.1016/j.cbpb.2013.07.006) PMID: [23876385](https://pubmed.ncbi.nlm.nih.gov/23876385/)
7. Dong JJ, Wu F, Ye X, Sun CF, Tian YY, Lu MX, et al. (2015) β -defensin in Nile tilapia (*Oreochromis niloticus*): sequence, tissue expression, and anti-bacterial activity of synthetic peptides. *Gene.* 566: 23–31. doi: [10.1016/j.gene.2015.04.025](https://doi.org/10.1016/j.gene.2015.04.025) PMID: [25871516](https://pubmed.ncbi.nlm.nih.gov/25871516/)
8. Nam BH, Moon JY, Kim YO, Kong HJ, Kim WJ, Lee SJ, et al. (2010) Multiple beta-defensin isoforms identified in early developmental stages of the teleost *Paralichthys olivaceus*. *Fish Shellfish Immunol.* 28: 267–274. doi: [10.1016/j.fsi.2009.11.004](https://doi.org/10.1016/j.fsi.2009.11.004) PMID: [19900559](https://pubmed.ncbi.nlm.nih.gov/19900559/)
9. Wang G, Li J, Zou P, Xie H, Huang B, Nie P, et al. (2012) Expression pattern, promoter activity and bactericidal property of beta-defensin from the mandarin fish *Siniperca chuatsi*. *Fish Shellfish Immunol.* 33: 522–531. doi: [10.1016/j.fsi.2012.06.003](https://doi.org/10.1016/j.fsi.2012.06.003) PMID: [22705342](https://pubmed.ncbi.nlm.nih.gov/22705342/)
10. Guo M, Wei J, Huang X, Huang Y, Qin Q (2012) Antiviral effects of beta-defensin derived from orange-spotted grouper (*Epinephelus coioides*). *Fish Shellfish Immunol.* 32: 828–838. doi: [10.1016/j.fsi.2012.02.005](https://doi.org/10.1016/j.fsi.2012.02.005) PMID: [22343108](https://pubmed.ncbi.nlm.nih.gov/22343108/)
11. Cuesta A, Meseguer J, Esteban MA (2011) Molecular and functional characterization of the gilthead seabream beta-defensin demonstrate its chemotactic and antimicrobial activity. *Mol Immunol.* 48: 1432–1438. doi: [10.1016/j.molimm.2011.03.022](https://doi.org/10.1016/j.molimm.2011.03.022) PMID: [21497909](https://pubmed.ncbi.nlm.nih.gov/21497909/)
12. Garcia-Valtanen P, Martinez-Lopez A, Ortega-Villaizan M, Perez L, Coll JM, Estepa A (2014) In addition to its antiviral and immunomodulatory properties, the zebrafish beta-defensin 2 (zFBD2) is a potent viral DNA vaccine molecular adjuvant. *Antiviral Res.* 101: 136–147. doi: [10.1016/j.antiviral.2013.11.009](https://doi.org/10.1016/j.antiviral.2013.11.009) PMID: [24286781](https://pubmed.ncbi.nlm.nih.gov/24286781/)
13. Katzenback BA (2015) Antimicrobial peptides as mediators of innate immunity in teleost. *Biology.* 4: 607–639. doi: [10.3390/biology4040607](https://doi.org/10.3390/biology4040607) PMID: [26426065](https://pubmed.ncbi.nlm.nih.gov/26426065/)
14. Wimley WC (2010) Describing the mechanism of antimicrobial peptide action with the interfacial activity model. *ACS Chem Biol.* 5: 905–917. doi: [10.1021/cb1001558](https://doi.org/10.1021/cb1001558) PMID: [20698568](https://pubmed.ncbi.nlm.nih.gov/20698568/)
15. Rivas L, Luque-Ortega JR, Fernandez-Reyes M, Andreu D (2010) Membrane-active peptides as anti-infectious agents. *J Appl Biomed.* 8: 159–167.

16. Li JG, Liu SP, Lakshminarayanan R, Bai Y, Pervushin K, Verma C, et al. (2013) Molecular simulations suggest how a branched antimicrobial peptide perturbs a bacterial membrane and enhances permeability. *Biochim Biophys Acta.* 1828: 1112–1121. doi: [10.1016/j.bbmem.2012.12.015](https://doi.org/10.1016/j.bbmem.2012.12.015) PMID: [23274275](https://pubmed.ncbi.nlm.nih.gov/23274275/)
17. Qi ZT, Wu P, Zhang QH, Wei YC, Wang ZS, Qiu M, et al. (2016) Transcriptome analysis of soiny mullet (*Liza haematocheila*) spleen in response to *Streptococcus dysgalactiae*. *Fish Shellfish Immunol.* 49: 194–204. doi: [10.1016/j.fsi.2015.12.029](https://doi.org/10.1016/j.fsi.2015.12.029) PMID: [26707943](https://pubmed.ncbi.nlm.nih.gov/26707943/)
18. Kumar S, Dudley J, Nei M, Tamura K (2008) MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Brief Bioinform.* 9: 299–306. doi: [10.1093/bib/bbn017](https://doi.org/10.1093/bib/bbn017) PMID: [18417537](https://pubmed.ncbi.nlm.nih.gov/18417537/)
19. Qi ZT, Zhang QH, Wang ZS, Zhao WH, Chen SN, Gao Q (2015) Molecular cloning, expression analysis and functional characterization of interleukin-22 in soiny mullet, *Liza haematocheila*. *Mol Immunol.* 63: 245–252. doi: [10.1016/j.molimm.2014.07.006](https://doi.org/10.1016/j.molimm.2014.07.006) PMID: [25108846](https://pubmed.ncbi.nlm.nih.gov/25108846/)
20. Prime, version 2.1, Schrödinger, LLC, New York, NY, 2009.
21. Hoover DM, Lubkowski CJ (2001) The structure of human β-defensin-1: new insights into structural properties of β-defensins. *J Biol Chem.* 276: 39021. PMID: [11486002](https://pubmed.ncbi.nlm.nih.gov/11486002/)
22. Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ (2005) PatchDock and SymmDock: servers for rigid and symmetric docking. *Nucleic Acids Res.* 33: W363–W367. PMID: [15980490](https://pubmed.ncbi.nlm.nih.gov/15980490/)
23. Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ (2005) Geometry based flexible and symmetric protein docking. *Proteins.* 60: 224–231. PMID: [15981269](https://pubmed.ncbi.nlm.nih.gov/15981269/)
24. Hess B, Kutzner C, van der Spoel D, Lindahl E (2008) GROMACS 4: Algorithms for highly efficient, load-balanced, and scalable molecular simulation. *J Chem Theory Comput.* 4: 435–447. doi: [10.1021/ct700301q](https://doi.org/10.1021/ct700301q) PMID: [26620784](https://pubmed.ncbi.nlm.nih.gov/26620784/)
25. Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C, et al. (2006) Comparison of multiple Amber force fields and development of improved protein backbone parameters. *Proteins.* 65: 712–725. PMID: [16981200](https://pubmed.ncbi.nlm.nih.gov/16981200/)
26. Domański J, Stansfeld P, Sansom MSP, Beckstein O (2010) Lipidbook: A public repository for force field parameters used in membrane simulations. *J Membrane Biol.* 236: 255–258.
27. Zhao W, Róg T, Gurtovenko AA, Vattulainen I, Karttunen M (2008) Role of phosphatidylglycerols in the stability of bacterial membranes. *Biochimie.* 90: 930–938. doi: [10.1016/j.biochi.2008.02.025](https://doi.org/10.1016/j.biochi.2008.02.025) PMID: [18373983](https://pubmed.ncbi.nlm.nih.gov/18373983/)
28. Jämbeck JP, Lyubartsev AP (2012) Derivation and systematic validation of a refined all-atom force field for phosphatidylcholine lipids. *J Phys Chem B.* 116:3164–3179. doi: [10.1021/jp212503e](https://doi.org/10.1021/jp212503e) PMID: [22352995](https://pubmed.ncbi.nlm.nih.gov/22352995/)
29. Jämbeck JP, Lyubartsev AP (2012) An extension and further validation of an all-atomistic force field for biological membranes, *J Chem Theory Comput.* 8: 2938–2948. doi: [10.1021/ct300342n](https://doi.org/10.1021/ct300342n) PMID: [26592132](https://pubmed.ncbi.nlm.nih.gov/26592132/)
30. Chaturvedi P, Dhanik M, Pande A (2015) Molecular characterization and in silico analysis of defensin from *Tor putitora* (Hamilton). *Probiotics Antimicro Prot.* 7: 207–215.
31. Kluver E, Adermann K, Schulz A (2006) Synthesis and structure-activity relationship of β-defensins, multi-functional peptides of the immune system. *J Pept Sci.* 12: 243–257. PMID: [16491514](https://pubmed.ncbi.nlm.nih.gov/16491514/)
32. Tu J, Li D, Li Q, Zhang L, Zhu Q, Gaur U, et al. (2015) Molecular evolutionary analysis β-defensin peptides in vertebrates. *Evol Bioinform Online.* 11:105–114. doi: [10.4137/EBO.S25580](https://doi.org/10.4137/EBO.S25580) PMID: [26056425](https://pubmed.ncbi.nlm.nih.gov/26056425/)
33. Casadei E, Wang TH, Zou J, González-Vecino JL, Wadsworth S, Secombes CJ (2009) Characterization of three novel β-defensin antimicrobial peptides in rainbow trout (*Oncorhynchus mykiss*). *Mol Immunol.* 46: 3358–3366. doi: [10.1016/j.molimm.2009.07.018](https://doi.org/10.1016/j.molimm.2009.07.018) PMID: [19709750](https://pubmed.ncbi.nlm.nih.gov/19709750/)
34. Hoover DM, Rajashankar KR, Blumenthal R, Puri A, Oppenheim JJ, Chertov O, et al. (2000) The structure of human β-defensin shows evidence of higher order oligomerization. *J Biol Chem.* 275:32911–32918. PMID: [10906336](https://pubmed.ncbi.nlm.nih.gov/10906336/)
35. Slutsky MM, Marsh ENG (2004) Cation-π interactions studied in a model coiled-coil peptide. *Protein Sci.* 13: 2244–2251. PMID: [15238639](https://pubmed.ncbi.nlm.nih.gov/15238639/)
36. Lu T, Chen FW (2012) Multiwfn: a multifunctional wave function analyzer, *J Comp Chem.* 35:580–592.
37. Zhao W, Róg T, Gurtovenko AA, Vattulainen I, Karttunen M (2007) Atomic-scale structure and electrostatics of anionic palmitoyloleoylphosphatidylglycerol lipid bilayers with Na⁺ counterions. *Biophys J.* 92: 1114–1124. PMID: [17114222](https://pubmed.ncbi.nlm.nih.gov/17114222/)
38. Ngo AT, Jakubek ZJ, Lu Z, Joós B, Morris CE, Johnston LJ (2014) Membrane order parameters for interdigitated lipid bilayers measured via polarized total-internal-reflection fluorescence microscopy. *Biochim Biophys Acta.* 1838: 2861–2869.

39. Fujii G, Selsted MM, Eisenberg D (1993) Defensins promote fusion and lysis of negatively charged membranes. *Protein Sci.* 2:1301–1312. PMID: [8401215](#)
40. Sudheendra US, Dhople V, Datta A, Kar RK, Shelburne CE, Bhunia A, et al (2015) Membrane disruptive antimicrobial activities of human-defensin-3 analogs. *Eur J Med Chem.* 91: 91–99. doi: [10.1016/j.ejmech.2014.08.021](#) PMID: [25112689](#)