

CORRECTION

## Correction: Modeling Occupancy of Hosts by Mistletoe Seeds after Accounting for Imperfect Detectability

Rodrigo F. Fadini, Renato Cintra

The images for Figs 2 and 3 are incorrectly switched. The image that appears as Fig 2 should be Fig 3, and the image that appears for Fig 3 should be Fig 2. The figure captions appear in the correct order. Please see the corrected Figs 2 and 3 here.





Citation: Fadini RF, Cintra R (2016) Correction: Modeling Occupancy of Hosts by Mistletoe Seeds after Accounting for Imperfect Detectability. PLoS ONE 11(9): e0163008. doi:10.1371/journal.pone.0163008

Published: September 16, 2016

Copyright: © 2016 Fadini, Cintra. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.






Fig 2. Occupancy and detection probability of mistletoe seeds of Psittacanthus plagiophyllus deposited on the host Anacardium occidentale according to proximity to infected hosts host size (host crown diameter). Central markers represent means, and lines represent 95% confidence intervals. Both graphs were traced with estimates from the model  $\Psi$  (size+neighborhood), p (size+neighborhood).

doi:10.1371/journal.pone.0163008.g001




Fig 3. Comparison of occupancy estimates of seeds of *Psittacanthus plagiophyllus* between two models: one using naïve estimates fitted with a logistic regression (logit (p) = 0.251 + 0.18 (host crown) - 1.64 (neighbor)), and other using occupancy estimates accounted for detectability (first model of Table 1).

doi:10.1371/journal.pone.0163008.g002



## Reference

 Fadini RF, Cintra R (2015) Modeling Occupancy of Hosts by Mistletoe Seeds after Accounting for Imperfect Detectability. PLoS ONE 10(5): e0127004. doi:10.1371/journal.pone.0127004 PMID: 25973754