

CORRECTION

Correction: Molecular Paleoclimate Reconstructions over the Last 9 ka from a Peat Sequence in South China

Xinxin Wang, Xianyu Huang, Dirk Sachse, Weihua Ding, Jiantao Xue

In Fig 4, the curve of hopanoids from a. Dajiuhu peat was plotted incorrectly. The y-axis scale of oxygen isotope values from c. Dongge should shift up by .5, so that the lower value is 6.5 and the upper value is 9.5. Please see the corrected Fig 4 here.



Citation: Wang X, Huang X, Sachse D, Ding W, Xue J (2016) Correction: Molecular Paleoclimate Reconstructions over the Last 9 ka from a Peat Sequence in South China. PLoS ONE 11(9): e0163650. doi:10.1371/journal.pone.0163650

Published: September 21, 2016

Copyright: © 2016 Wang et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 4. Comparisons of records. (a) Mass accumulation rate of hopanoids in the Dajiuhu peatland [26]. (b) BNA15 record from the Dajiuhu peatland [24], (c) δ 18Ocarb record from Dongge Cave [30], (d) δ DC31 record from the Hongyuan peat sequence [19], (e) ratios of total herb pollen over total tree pollen from the Daping sedimentary sequence [29], (f) δ 13C C29 record (this study) and (g) δ DC29 record (this study).

doi:10.1371/journal.pone.0163650.g001

Reference

1. Wang X, Huang X, Sachse D, Ding W, Xue J (2016) Molecular Paleoclimate Reconstructions over the Last 9 ka from a Peat Sequence in South China. PLoS ONE 11(8): e0160934. doi: 10.1371/journal. pone.0160934 PMID: 27505008