

RESEARCH ARTICLE

# German validation of Quality of Life after Brain Injury (QOLIBRI) assessment and associated factors


Nicole von Steinbüchel<sup>1☯‡</sup>, Ruben G. L. Real<sup>1☯‡\*</sup>, Nadine Sasse<sup>1</sup>, Lindsay Wilson<sup>2</sup>, Christiane Otto<sup>1,3</sup>, Ryan Mullins<sup>1</sup>, Robert Behr<sup>4</sup>, Wolfgang Deinsberger<sup>5</sup>, Ramon Martinez-Olivera<sup>6</sup>, Wolfgang Puschendorf<sup>7</sup>, Werner Petereit<sup>8</sup>, Veit Rohde<sup>9</sup>, Holger Schmidt<sup>1,10</sup>, Stephan Sehmisch<sup>11</sup>, Klaus Michael Stürmer<sup>11</sup>, Klaus von Wild<sup>12</sup>, Henning Gibbons<sup>1,13</sup>

**1** Institute of Medical Psychology and Medical Sociology, University Medical Center Goettingen, Goettingen, Germany, **2** Department of Psychology, University of Stirling, Stirling, United Kingdom, **3** Department of Child and Adolescent Psychiatry, Psychotherapy and Psychosomatics, University Medical Center Hamburg Eppendorf, Hamburg, Germany, **4** Department of Neurosurgery, Clinical Center Fulda, Fulda, Germany, **5** Department of Neurosurgery, Clinical Center Kassel, Kassel, Germany, **6** Department of Neurosurgery & Neurotraumatology at Bergmannsheil University Hospital Bochum, Bochum, Germany, **7** Westend Neurological Clinic, Bad Wildungen, Germany, **8** Department of Neurosurgery, Clinical Center Bernburg, Bernburg, Germany, **9** Department of Neurosurgery, University Medical Center Goettingen, Goettingen, Germany, **10** Department of Neurology, University Medical Center Goettingen, Goettingen, Germany, **11** Trauma surgery, plastic and reconstructive surgery, University Medical Center Goettingen, Goettingen, Germany, **12** KvW Neuroscience Consulting, Muenster, Germany, **13** Department of Psychology, University of Bonn, Bonn, Germany

☯ These authors contributed equally to this work.

‡ These authors are co-first authors on this work.

\* [ruben.real@med.uni-goettingen.de](mailto:ruben.real@med.uni-goettingen.de)



**OPEN ACCESS**

**Citation:** von Steinbüchel N, Real RGL, Sasse N, Wilson L, Otto C, Mullins R, et al. (2017) German validation of Quality of Life after Brain Injury (QOLIBRI) assessment and associated factors. *PLoS ONE* 12(5): e0176668. <https://doi.org/10.1371/journal.pone.0176668>

**Editor:** Rita Formisano, Santa Lucia Foundation, ITALY

**Received:** July 25, 2016

**Accepted:** April 12, 2017

**Published:** May 24, 2017

**Copyright:** © 2017 von Steinbüchel et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This project was funded by the ZNS – Hannelore Kohl Foundation (project number 2008014; [www.hannelore-kohl-stiftung.de](http://www.hannelore-kohl-stiftung.de)), the German Research Foundation, and the Open Access Publication Funds of the Göttingen University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Abstract

The consequences of traumatic brain injury (TBI) for health-related quality of life (HRQoL) are still poorly understood, and no TBI-specific instrument has hitherto been available. This paper describes in detail the psychometrics and validity of the German version of an internationally developed, self-rated HRQoL tool after TBI—the QOLIBRI (Quality of Life after Brain Injury). Factors associated with HRQoL, such as the impact of cognitive status and awareness, are specifically reported. One-hundred seventy-two participants after TBI were recruited from the records of acute clinics, most of whom having a Glasgow Coma Scale (GCS) 24-hour worst score and a Glasgow Outcome Scale (GOSE) score. Participants had severe (24%), moderate (11%) and mild (56%) injuries as assessed on the GCS, 3 months to 15 years post-injury. The QOLIBRI uses 37 items to measure “satisfaction” in the areas of “Cognition”, “Self”, “Daily Life and Autonomy”, and “Social Relationships”, and “feeling bothered” by “Emotions” and “Physical Problems”. The scales meet standard psychometric criteria ( $\alpha = .84$  to  $.96$ ; intra-class correlation—ICC =  $.72$  to  $.91$ ). ICCs (0.68 to 0.90) and  $\alpha$ s (.83 to  $.96$ ) were also good in a subgroup of participants with lower cognitive performance. The six-subscale structure of the international sample was reproduced for the German version using confirmatory factor analyses and Rasch analysis. Scale validity was supported by systematic relationships observed between the QOLIBRI and the GOSE, Patient Competency Rating Scale for Neurorehabilitation (PCRS-NR), Hospital Anxiety and Depression Scale

**Competing interests:** The authors have declared that no competing interests exist.

(HADS), Profile of Mood States (POMS), Short Form 36 (SF-36), and Satisfaction with Life Scale (SWLS). The German QOLIBRI contains novel information not provided by other currently available measures and has good psychometric criteria. It is potentially useful for clinicians and researchers, in post-acute and rehabilitation studies, on a group and individual level.

## Introduction

Health care and rehabilitation programmes following traumatic brain injury (TBI) aim to restore a person to everyday life with as high a quality of life (HRQoL) as possible. Accomplishing this poses remarkable challenges given that TBI can result in lifelong physical, cognitive, emotional, and behavioural impairments, as well as significant restrictions in social participation. Outcomes after TBI have traditionally been assessed using functional indicators, such as disability [1,2], health status [3–6], return to work or productivity [7–9], or psychosocial and social functioning [10–12]. While these outcomes are certainly linked to health-related quality of life (HRQoL) after TBI, they do not incorporate the specific subjective perspective of wellbeing and functioning, which are important aspects of HRQoL [13,14]. However, lately wellbeing and HRQoL have become important outcome variables after TBI: even patients after mild TBI (mTBI) show significantly reduced general wellbeing in a prospective study with a longitudinal design [15]. Very little literature exists on the assessment of wellbeing or HRQoL in German TBI samples. German studies have used the Aachen Life Quality Inventory [16] or the SF-36 [5,17], both of which are instruments for assessing generic HRQoL. However, when so assessed even individuals after mTBI display significantly reduced general wellbeing [18].

QoL is defined by the World Health Organization QOL Group [19] (p. 153) as “... an individual’s perception of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectations, standards, and concerns. ...”. HRQoL refers to the specific effects of health conditions on a person’s wellbeing and functioning. HRQoL measures are increasingly regarded as essential to elicit a more complete assessment of treatment effects and providing information concerning the patient’s subjective experience via self-reporting. Such patient-reported outcomes (PRO) therefore provide a measure of the subjective outcome and, furthermore, attempt to avoid significant clinician bias in outcome studies. Generic HRQoL instruments and measures of subjective health, such as the SF-36 [20] or the Sickness Impact Profile (SIP, [21]), do not cover specific areas that are relevant to TBI, for example, cognition or self-esteem, as well as future prospects. A disease-specific HRQoL instrument may be more sensitive to the particular consequences of TBI and more relevant to the patient. In response to the need for a disease-specific measure of HRQoL after TBI, an international multidisciplinary group was formed to develop the QOLIBRI descriptive system [14,22,23]. The instrument was constructed in a consensual fashion, in several languages simultaneously, and validated in two cross-sectional studies. The second international study included six language versions of the QOLIBRI (Dutch, English, Finnish, French, German, and Italian) and a total of 921 persons with TBI [22,23].

The purpose of this study is to examine the psychometric properties of the German language version of the QOLIBRI. Some information concerning the reliability and validity of different language versions has already been reported in the context of the international study [22,23], but there has been a lack of a detailed examination of individual language versions to show that their psychometric properties are appropriate for use.

As well as studying reliability and aspects of construct validity, we were particularly interested in whether confirmatory factor analysis in a single-language version, in this case German, would support the QOLIBRI's descriptive system. The second aim of this study is to extend the validation of the QOLIBRI by examining relationships with measures that were not part of the international studies, namely measures of disability, mood states, and general wellbeing and satisfaction. Additionally, the generally little investigated issue of the influence of impaired cognition on self-reported HRQoL after brain injury is examined here in more detail.

## Materials and methods

### Participants

A total of 172 individuals (118 male) after TBI participated in the study. Patients were recruited from hospitals in the German communities of Goettingen, Fulda and Kassel ( $n = 166$ ). Additionally, six individuals from a rehabilitation centre took part. The *inclusion criteria* were: ICD-10 diagnosis of TBI, 3 months to 15 years post-injury, aged 15 years or more at time of injury, outpatient status, aged 17 to 68 years at time of interview, and able and willing to give written informed consent. *Exclusion criteria* included a Glasgow Outcome Score-Extended (GOSE; [24]) score less than 3 (vegetative state); spinal cord injury; significant current or pre-injury psychiatric history; ongoing severe addiction; inability to understand, cooperate and answer; and having a terminal illness. In contrast to the two international developmental and validation studies reported elsewhere [22,23], a Glasgow Coma Scale (GCS) score [25] was not required for inclusion in the present study, however a GCS score was obtained for 90% of participants. The study was approved by the local ethical review board of the University Medical Center Göttingen (No. 26308) and conformed to the Declaration of Helsinki [26].

### Measures

The QOLIBRI (Quality of Life after Brain Injury) questionnaire with its 37 items generates a profile of HRQoL in the six domains: "Cognition" (7 items), "Self" (7 items), "Daily Life and Autonomy" (7 items), "Social Relationships" (6 items), "Emotions" (5 items), and "Physical Problems" (5 items), as well as a total score. The first four scales contain items requiring "satisfaction" judgements, whereas the last two scales ask for judgements about how "bothered" the person feels. Patient responses are assessed via a five-point Likert scale from "not at all" to "very".

To measure wellbeing in terms of generic HRQoL, the Satisfaction with Life Scale (SWLS; [27]) and the Short Form-36 (SF-36, [20]) were administered. In its usual form, the SWLS has seven-point Likert response scales. In the present study, five-point scales were used in order to simplify the scale for patients with cognitive deficits ("I disagree completely" to "I agree completely"). SF-36 Physical (PCS) and Mental (MCS) component summary scores were calculated using US normative data. The global functional status was assessed using the Glasgow Outcome Scale-Extended (GOSE). Interviewers were trained to administer the GOSE based on the manual [24]. Disabilities and competencies were investigated by means of the Patient Competency Rating Scale for Neurorehabilitation (PCRS-NR) [28], providing a total score, as well as three subscale scores (Emotional Functioning, Interpersonal Functioning, Cognitive Functioning).

The Hospital Anxiety and Depression Scale (HADS; [29]) served as a measure of emotional distress and symptoms of the participants. Depression and anxiety were categorised using conventional cut-offs (i.e. 0–7 represented the normal range, 8–10 mild, 11–14 moderate, and 15

or above severe depression or anxiety) [29]. Mood was screened in this study using a short version of the Profile of Mood States (POMS, [30]). This instrument has been validated in the German language and consists of 35 items on four subscales (Depression/Dejection, Vigour/Activity, Fatigue/Inertia, and Anger/Hostility).

The Telephone Interview for Cognitive Status (TICS; [31]) was administered to screen the cognitive status of the participant. A cut-off of 32/33 was used to define two groups, having low cognitive performance and normal performance.

Socio-demographic data, including age, gender, education, employment, profession, living situation, autonomy, leisure interests and activities, aspects of social life, alcohol and nicotine consumption, as well as participation in rehabilitation, were assessed using a questionnaire that was completed by the participants [32].

Participants' current health conditions were self-reported on a health symptoms list of 28 health conditions (adapted from the WHO-QoL project; [32]). For severity of injury, the 24-hour worst Glasgow Coma Scale (GCS; [25]) score was used, a standard index for severity classification, whereby scores of 3 to 8 indicate severe injury, 9 to 12 moderate injury, and 13 to 15 indicate mild injury. Comorbid health problems, such as epilepsy, hemiparesis, vision and hearing problems, extra-cerebral injuries, communication, attention and memory dysfunction, executive function, and affective and behavioural disorders, as well as participation in rehabilitation and medications, were also noted.

## Data collection

The GOSE and TICS were administered as a telephone interview. During this interview, a clinical checklist was also completed that covered the inclusion and exclusion criteria for the study as well as comorbid health conditions. The 24-hour worst GCS score was obtained from the patients' medical records. These records also provided information concerning major brain lesions and the date of injury. HRQoL was assessed via self-rated questionnaires.

Two methods were used for data collection: a) patients received and returned the questionnaire by post and were interviewed by telephone ( $N = 168$ ) or b) patients visited the clinic to complete the questionnaire and were interviewed in a face-to-face meeting ( $N = 4$ ). The QOLIBRI was assessed at two points in time, i.e. test and retest, with an average interval of two weeks.

## Statistical analyses

We followed the approach used in the international sample, as described in detail elsewhere [22,23]. Data were analysed using PASW 18 software. Item scores on the "bothered" QOLIBRI scales "Emotions" and "Physical Problems" were recoded to match scores on the "satisfaction" scales. Missing values were prorated if less than 33% of the answers per scale were missing, and mean percentage scores were calculated for the six QOLIBRI subscales and the QOLIBRI total.

The analyses included descriptive and confirmatory analyses. The psychometric analyses focused on I) factorial validity of the QOLIBRI in the German sample, II) reliability and III) construct validity, and IV) partial correlations.

1. Classical as well as modern test-theoretical approaches were applied. Rasch analysis, Principal Components Analysis (PCA), and Confirmatory Factor Analysis (CFA) using methods based on Structural Equation Modelling (SEM) served to investigate the factorial validity of the QOLIBRI. Rasch analyses were conducted using the Winsteps 3.66 implementation of a partial credit model [33]. Item fit to a Rasch model was examined using 'infit' and 'outfit' statistics: values between 0.70 and 1.30 are regarded as acceptable for rating scales. CFA was

carried out using Amos 18. Likewise, Promax rotation in PCA was used, and a second-order CFA model was specified, including a higher order factor besides the six latent variables representing the six QOLIBRI scales. To describe how well our model fits the data, recommended fit indices were chosen which are relatively independent of sample size [34]. The Root Mean Square Error of Approximation (RMSEA; [35]) served to describe overall fit, while the incremental Comparative Fit Index (CFI; [36]) compared our model to a baseline model. In addition, the Standardized Root Mean Square Residual (SRMR; [37]) focuses on residuals unexplained by the specified model. The following “rules of thumb” were used for the fit indices: an RMSEA  $\leq 0.05$  indicates a close fit, while a value  $\leq 0.08$  represents an acceptable fit [38]. For the CFI, a value  $\geq 0.97$  stands for a close fit, but a CFI  $\geq 0.95$  represents an acceptable fit [34]. Finally, an SRMR  $\leq 0.05$  indicates a close fit and values  $\leq 0.1$  represent acceptable one [39].

2. The internal consistency of the QOLIBRI was analysed by calculating Cronbach’s  $\alpha$  for each scale using data from the first and second time points. Cronbach’s  $\alpha \geq 0.70$  indicates acceptable reliability for measures used in group comparisons [40], while measures applied to individuals in clinical settings should reach values  $\geq 0.90$  [41]. To investigate the test-retest reliability, we calculated the intra-class correlation coefficient (ICC). ICCs between 0.40 and 0.75 are usually interpreted as fair to good, while ICCs  $> 0.75$  indicate excellent test-retest reliability [42]. To analyse the reliability of the QOLIBRI in relation to the cognitive status of participants, Cronbach’s  $\alpha$  and ICC were calculated for subgroups of patients defined by the TICS using a recommended cut-off (i.e. a sum score  $< 33$  indicates a low cognitive status, while a score  $> 32$  represents a normal cognitive status; [43]).
3. The analysis involved a mixture of continuous and ordinal measures, and some skewness was present in the majority of variables. Unless otherwise noted, analyses were carried out on ranked variables [44], and Spearman correlations were used to examine relationships between HRQoL and other variables.
4. Partial correlations were used to assess whether the QOLIBRI total score contained information not included in other measures.

## Results

### Sample characteristics

The sample consisted of 172 patients with TBI who met the inclusion criteria and gave their informed consent to participate in this study; they completed the QOLIBRI questionnaire and were interviewed by telephone for the GOSE and TICS. In most cases, a telephone interview was conducted, followed by the collection of self-reported data by post (98%). The response rate for the QOLIBRI retest was 76% (i.e. all study participants were asked to take part in the retest, and 130 individuals responded). Table 1 provides a description of the sample.

More than half of the participants were in the mild range on the GCS and more than half of the sample achieved a good recovery on the GOSE. In addition, more than 70% of the patients were living independently; while on the HADS 73% were classified as below the threshold for anxiety and 72% below the threshold for depression. Moreover, 45% of the sample had a low cognitive status on the TICS.

### Factorial validity

Less than 5% of the responses on QOLIBRI items were missing at initial testing (e.g. satisfaction with sexual life; 4.1%). However, 10% of the patients did not answer the QOLIBRI item

**Table 1. Characteristics of the sample (N = 172).**

|  | <b>Group</b>        | <b>Frequency (rate)</b> |
|--|---------------------|-------------------------|
| <b>Sex</b>   | Male                | 118 (69%)               |
|  | Female              | 54 (31%)                |
| <b>Age</b>   | 17 to 30 years      | 34 (20%)                |
|  | 31 to 44 years      | 35 (20%)                |
|  | 45 to 68 years      | 103 (60%)               |
| <b>Employment status</b>                               | Full-time           | 67 (39%)                |
| <b>Relationship status</b>                             | Single              | 34 (20%)                |
|  | Partnered           | 120 (70%)               |
|  | Past partnered      | 17 (10%)                |
| <b>Living arrangements</b>                             | Independent         | 123 (72%)               |
|  | Supported           | 48 (28%)                |
| <b>Years since injury</b>                              | <1 year             | 20 (12%)                |
|  | 1 to <2 years       | 24 (14%)                |
|  | 2 to < 4 years      | 53 (31%)                |
|  | 4 to 15 years       | 75 (44%)                |
| <b>Major lesion</b>                                    | None                | 37 (22%)                |
|  | Focal               | 113 (66%)               |
|  | Diffuse             | 6 (4%)                  |
| <b>Glasgow Coma Scale (worst score, first 24 hrs.)</b> | Severe (3–8)        | 40 (24%)                |
|  | Moderate (9–12)     | 18 (11%)                |
|  | Mild (13–15)        | 97 (56%)                |
| <b>Glasgow Outcome Scale—Extended score</b>            | Severe disability   | 20 (12%)                |
|  | Moderate disability | 60 (35%)                |
|  | Good recovery       | 92 (54%)                |
| <b>Number of comorbid health conditions</b>            | 0 to 3              | 71 (41%)                |
|  | 4 to 6              | 40 (23%)                |
|  | 7 or more           | 61 (36%)                |
| <b>HADS<sup>1</sup> - anxiety</b>                      | Normal (0–7)        | 125 (73%)               |
|  | Mild (8–10)         | 19 (11%)                |
|  | Moderate (11–14)    | 19 (11%)                |
|  | Severe (15–21)      | 9 (5%)                  |
| <b>HADS—depression</b>                                 | Normal (0–7)        | 123 (72%)               |
|  | Mild (8–10)         | 19 (11%)                |
|  | Moderate (11–14)    | 25 (15%)                |
|  | Severe (15–21)      | 5 (3%)                  |
| <b>TICS<sup>2</sup> - cognitive status</b>             | Low                 | 77 (45%)                |
|  | Normal              | 95 (55%)                |

<sup>1</sup> Hospital Anxiety and Depression Scale;

<sup>2</sup> Telephone Interview for Cognitive Status.

<https://doi.org/10.1371/journal.pone.0176668.t001>

concerning “participation in work or education”. This can be explained by the fact that in general this question does not apply to persons who have retired (12 out of 17 of these did not respond). After prorating, the QOLIBRI data collected at the first time point and at the retest were complete for each patient. However, we detected a significant deviation from a normal distribution for the QOLIBRI items (i.e. many items were skewed). Therefore, the ranked items were used in the analysis.

**Table 2. Principal components analyses of QOLIBRI items.** Loadings < 0.25 are suppressed.

| QOLIBRI scale                    | Item                      | 1 <sup>st</sup> component | h <sup>2</sup> | Factor 1     | Factor 2     | Factor 3     | Factor 4     | Factor 5     | Factor 6     |
|----------------------------------|---------------------------|---------------------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Cognition</b>                 | Concentrate               | 0.719                     | 0.693          |              | <b>0.685</b> |              |              | 0.253        |              |
|                                  | Express yourself          | 0.574                     | 0.601          |              | <b>0.791</b> |              |              |              |              |
|                                  | Remember                  | 0.565                     | 0.662          |              | <b>0.945</b> |              |              |              |              |
|                                  | Plan and problem solve    | 0.667                     | 0.684          |              | <b>0.801</b> |              |              |              |              |
|                                  | Decisions                 | 0.723                     | 0.630          | 0.400        | <b>0.451</b> |              |              |              |              |
|                                  | Find way                  | 0.712                     | 0.635          |              | <b>0.614</b> |              |              |              |              |
|                                  | Speed of thinking         | 0.730                     | 0.691          |              | <b>0.723</b> |              |              |              |              |
| <b>Self</b>                      | Energy                    | 0.753                     | 0.624          |              |              |              |              |              | <b>0.438</b> |
|                                  | Motivation                | 0.736                     | 0.589          | <b>0.299</b> |              |              |              |              | 0.256        |
|                                  | Self-esteem               | 0.748                     | 0.718          |              |              |              |              |              | <b>0.656</b> |
|                                  | Way you look              | 0.582                     | 0.547          |              |              |              |              |              | <b>0.683</b> |
|                                  | Achievements              | 0.702                     | 0.639          |              |              |              |              |              | <b>0.621</b> |
|                                  | Self-perception           | 0.755                     | 0.707          |              |              |              |              |              | <b>0.640</b> |
|                                  | Own future                | 0.715                     | 0.564          |              |              |              |              |              | <b>0.374</b> |
| <b>Daily life &amp; autonomy</b> | Independence              | 0.721                     | 0.723          | <b>0.809</b> |              |              |              |              |              |
|                                  | Get out & about           | 0.705                     | 0.707          | <b>0.863</b> |              |              |              |              |              |
|                                  | Domestic activities       | 0.619                     | 0.582          | <b>0.834</b> |              |              |              |              |              |
|                                  | Run personal finances     | 0.621                     | 0.540          | <b>0.688</b> |              |              |              |              |              |
|                                  | Participation work        | 0.744                     | 0.617          | <b>0.546</b> |              |              |              |              |              |
|                                  | Social&leisure activities | 0.737                     | 0.661          | <b>0.601</b> |              |              |              |              |              |
|                                  | In charge of life         | 0.758                     | 0.703          | <b>0.762</b> |              |              |              |              |              |
| <b>Social relation-ships</b>     | Affection towards others  | 0.607                     | 0.700          |              |              | <b>0.780</b> |              |              |              |
|                                  | Family                    | 0.615                     | 0.647          |              |              | <b>0.667</b> |              |              |              |
|                                  | Friends                   | 0.645                     | 0.625          |              | <b>0.369</b> | <b>0.367</b> | 0.333        |              |              |
|                                  | Partner                   | 0.431                     | 0.738          |              |              | <b>0.901</b> |              |              |              |
|                                  | Sex life                  | 0.566                     | 0.571          |              |              | <b>0.695</b> |              |              |              |
|                                  | Attitudes of others       | 0.659                     | 0.581          | 0.257        |              | <b>0.530</b> |              |              |              |
| <b>Emotions</b>                  | Loneliness                | 0.592                     | 0.715          |              |              |              | <b>0.803</b> |              |              |
|                                  | Boredom                   | 0.410                     | 0.604          |              |              |              | <b>0.858</b> |              | -0.289       |
|                                  | Anxiety                   | 0.584                     | 0.635          |              |              |              | <b>0.743</b> |              |              |
|                                  | Depression                | 0.670                     | 0.732          |              |              |              | <b>0.690</b> |              | 0.327        |
|                                  | Anger/aggression          | 0.527                     | 0.501          |              |              |              | <b>0.624</b> |              |              |
| <b>Physical problems</b>         | Slow/clumsiness           | 0.666                     | 0.736          |              |              |              |              | <b>0.709</b> |              |
|                                  | Other injuries            | 0.405                     | 0.607          |              |              |              |              | <b>0.776</b> |              |
|                                  | Pain                      | 0.496                     | 0.552          |              |              |              | 0.320        | <b>0.624</b> |              |
|                                  | See/hear                  | 0.351                     | 0.579          |              |              |              |              | <b>0.708</b> | -0.396       |
|                                  | TBI effects               | 0.635                     | 0.735          |              |              |              |              | <b>0.651</b> |              |

<https://doi.org/10.1371/journal.pone.0176668.t002>

A Rasch analysis of each QOLIBRI subscale in turn indicated an acceptable fit for the majority of items. Minor deviations of infit were observed for two items: “Way you look” (1.34) and “See/hear” (1.36). Deviations of outfit were also found for two items: “Decisions” (1.31) and “See/hear” (1.35). Scale person reliabilities ranged from .81 to .90 for the first four scales, and were lower for the last two scales, “Emotions” (.63) and “Physical Problems” (.69). The interpretation of these reliability statistics is similar to Cronbach’s alpha, and indicates that the last two scales are somewhat weaker than the first four and show poorer discrimination between persons.

Two PCAs were conducted, forcing either one or six components (see [Table 2](#)).

**Table 3. Inter-correlation of QOLIBRI scales.**

| QOLIBRI scales        | Cognition | Self  | Daily life & autonomy | Social relation-ships | Emotions |
|-----------------------|-----------|-------|-----------------------|-----------------------|----------|
| Self                  | 0.72*     |       |                       |                       |          |
| Daily life & autonomy | 0.67*     | 0.72* |                       |                       |          |
| Social relationships  | 0.51*     | 0.69* | 0.57*                 |                       |          |
| Emotions              | 0.49*     | 0.60* | 0.54*                 | 0.62*                 |          |
| Physical problems     | 0.54*     | 0.55* | 0.64*                 | 0.36*                 | 0.47*    |

Pearson correlations shown;  
\* p<0.001

<https://doi.org/10.1371/journal.pone.0176668.t003>

Table 2 shows a one-factor solution and a six-factor solution. In the one-factor solution ( $R^2 = .41$ ) all QOLIBRI items had a satisfactory loading ( $>0.40$ ), except for item “problems in seeing and/or hearing”. In the six-factor solution ( $R^2 = .64$ ), all but two QOLIBRI items had their highest loading on the appropriate scale, the number of cross-loadings ( $>.25$ ) was low, and all factors had an *eigenvalue*  $> 1.0$ . The “Motivation” item had its highest loading on the factor representing “Daily Life and Autonomy” scale, while the loadings for the item “Friends” were nearly identical on the factors corresponding to the “Social Relationships” and “Cognition” scales.

Moderate to high intercorrelations of the six QOLIBRI scales (Table 3) suggest a second-order CFA model, which is in line with the concept of an overall dimension of QoL representing subjective wellbeing. The second-order CFA model was specified (see Fig 1) and tested ( $\chi^2 = 1,007$ ,  $df = 623$ ,  $RMSEA = 0.060$ ,  $SRMR = 0.070$  and  $CFI = 0.897$ ). Our model showed an acceptable overall fit on RMSEA, but the CFI for our model indicated lack of fit. On the other hand, the SRMR showed an acceptable fit of the model with respect to residuals. The loadings of the first-order factors on the second-order factor were high, ranging from .71 to .91. As is common in SEM-based second-order CFA models, measurement errors for each indicator/item were included in the model (i.e. e1 to e37) and a residual term was estimated for each first-order factor (i.e. d1 to d6).

## Reliability

Cronbach’s  $\alpha$ s for the QOLIBRI scales are shown in Table 4.


Similarly, the ICCs indicate a good to excellent test-retest reliability (Table 5). Analyses additionally revealed that the QOLIBRI is reliable for TBI patients with low as well as a normal cognitive status (see Tables 4 and 5; note that these analyses are based on unranked data).

## Construct validity

The only gender difference found was for the “Cognition” scale ( $p = 0.022$ ), where the mean for women was slightly higher (mean = 88.18;  $SD = 43.84$ ) than for men (mean = 85.73;  $SD = 52.33$ ). For age, weak correlations were observed for the “Cognition” ( $r = -0.166$ ;  $p = 0.030$ ), “Daily Life and Autonomy” ( $r = -0.235$ ;  $p = 0.002$ ), “Physical Problems” ( $r = -0.235$ ;  $p = 0.002$ ), and the total QOLIBRI scales ( $r = -0.159$ ;  $p = 0.037$ ), indicating a decline in HRQoL with age. None of the other QOLIBRI scales had significant correlations with age.

Table 6 depicts relationships between the QOLIBRI scales, injury severity and outcome variables. The relationships shown in Table 6 were previously examined in the international study [22,23], and are included here for comparison with the larger group. HRQoL was not significantly correlated to severity of injury on the GCS, but significant positive associations were





**Fig 1. Path diagram of the second-order model of the QOLIBRI.**

<https://doi.org/10.1371/journal.pone.0176668.g001>

found with the GOSE for all QOLIBRI scales. HRQoL was related both to the amount of help needed in five activities of daily living and to the number of comorbid health conditions. TBI-specific HRQoL was significantly and negatively associated with HADS anxiety as well as HADS depression for each QOLIBRI scale.

**Table 4. Internal consistencies at test and retest for QOLIBRI scales for all participants, and for groups with low or normal cognitive status.**

|                                  | All |                      |     |                        | Low TICS <sup>1</sup> |                      |    |                        | High TICS |                      |    |                        |
|----------------------------------|-----|----------------------|-----|------------------------|-----------------------|----------------------|----|------------------------|-----------|----------------------|----|------------------------|
|                                  | n   | Cronbach's α at test | n   | Cronbach's α at retest | n                     | Cronbach's α at test | n  | Cronbach's α at retest | n         | Cronbach's α at test | n  | Cronbach's α at retest |
| <b>Cognition</b>                 | 172 | 0.91                 | 130 | 0.92                   | 77                    | 0.90                 | 64 | 0.90                   | 95        | 0.89                 | 66 | 0.93                   |
| <b>Self</b>                      | 172 | 0.90                 | 130 | 0.91                   | 77                    | 0.90                 | 64 | 0.89                   | 95        | 0.89                 | 66 | 0.91                   |
| <b>Daily life &amp; autonomy</b> | 172 | 0.90                 | 130 | 0.92                   | 77                    | 0.89                 | 64 | 0.91                   | 95        | 0.89                 | 66 | 0.91                   |
| <b>Social relationships</b>      | 172 | 0.85                 | 130 | 0.85                   | 77                    | 0.85                 | 64 | 0.81                   | 95        | 0.86                 | 66 | 0.88                   |
| <b>Emotions</b>                  | 172 | 0.84                 | 129 | 0.86                   | 77                    | 0.83                 | 63 | 0.86                   | 95        | 0.82                 | 66 | 0.80                   |
| <b>Physical problems</b>         | 172 | 0.84                 | 130 | 0.81                   | 77                    | 0.84                 | 64 | 0.79                   | 95        | 0.81                 | 66 | 0.82                   |
| <b>QOLIBRI total</b>             | 172 | 0.96                 | 130 | 0.97                   | 77                    | 0.96                 | 64 | 0.96                   | 95        | 0.96                 | 66 | 0.97                   |

<sup>1</sup> Telephone Interview for Cognitive Status.

<https://doi.org/10.1371/journal.pone.0176668.t004>

**Table 5. Test-retest intra-class correlations for all retested participants, and for groups with low or normal cognitive status.**

|                                  | All |                  |                           |              | Low TICS <sup>1</sup> |      |              |              | High TICS |      |              |              |
|----------------------------------|-----|------------------|---------------------------|--------------|-----------------------|------|--------------|--------------|-----------|------|--------------|--------------|
|                                  | n   | ICC <sup>2</sup> | Lower 95% CI <sup>3</sup> | Upper 95% CI | N                     | ICC  | Lower 95% CI | Upper 95% CI | n         | ICC  | Lower 95% CI | Upper 95% CI |
| <b>Cognition</b>                 | 130 | 0.85             | 0.80                      | 0.89         | 64                    | 0.84 | 0.75         | 0.90         | 66        | 0.83 | 0.74         | 0.89         |
| <b>Self</b>                      | 130 | 0.87             | 0.83                      | 0.91         | 64                    | 0.85 | 0.76         | 0.91         | 66        | 0.88 | 0.82         | 0.93         |
| <b>Daily life &amp; autonomy</b> | 130 | 0.87             | 0.82                      | 0.90         | 64                    | 0.87 | 0.79         | 0.92         | 66        | 0.85 | 0.77         | 0.91         |
| <b>Social relationships</b>      | 130 | 0.77             | 0.69                      | 0.83         | 64                    | 0.72 | 0.58         | 0.82         | 66        | 0.82 | 0.72         | 0.88         |
| <b>Emotions</b>                  | 129 | 0.72             | 0.63                      | 0.80         | 63                    | 0.68 | 0.52         | 0.79         | 66        | 0.72 | 0.58         | 0.82         |
| <b>Physical problems</b>         | 130 | 0.91             | 0.88                      | 0.94         | 64                    | 0.90 | 0.84         | 0.94         | 66        | 0.92 | 0.87         | 0.95         |
| <b>QOLIBRI total</b>             | 130 | 0.91             | 0.87                      | 0.94         | 64                    | 0.89 | 0.83         | 0.93         | 66        | 0.91 | 0.86         | 0.94         |

<sup>1</sup> Telephone Interview for Cognitive Status;

<sup>2</sup> intra-class correlation;

<sup>3</sup> confidence interval.

<https://doi.org/10.1371/journal.pone.0176668.t005>

**Table 6. Relationships between QOLIBRI scales and GCS, GOSE, HADS, and SF-36.**

|                                   | Cognition | Self     | Daily life & autonomy | Social relationships | Emotions | Physical problems | QOLIBRI total |
|-----------------------------------|-----------|----------|-----------------------|----------------------|----------|-------------------|---------------|
| <b>GCS</b>                        | 0.12      | 0.04     | 0.12                  | 0.01                 | <-0.01   | 0.14              | 0.08          |
| <b>GOSE</b>                       | 0.45***   | 0.38***  | 0.52***               | 0.17*                | 0.26**   | 0.61***           | 0.50***       |
| <b>Help needed</b>                | -0.41***  | -0.42*** | -0.63***              | -0.30***             | -0.36*** | -0.51***          | -0.54***      |
| <b>Comorbid health conditions</b> | -0.46***  | -0.52*** | -0.50***              | -0.34***             | -0.50*** | -0.67***          | -0.61***      |
| <b>HADS anxiety</b>               | -0.60***  | -0.66*** | -0.54***              | -0.43***             | -0.69*** | -0.51***          | -0.71***      |
| <b>HADS depression</b>            | -0.59***  | -0.78*** | -0.67***              | -0.67***             | -0.63*** | -0.49***          | -0.79***      |
| <b>SF-36 PCS</b>                  | 0.44***   | 0.42***  | 0.55***               | 0.17*                | 0.28***  | 0.74***           | 0.54***       |
| <b>SF-36 MCS</b>                  | 0.46***   | 0.64***  | 0.48***               | 0.65***              | 0.70***  | 0.37***           | 0.66***       |


\* p<0.05,

\*\* p<0.01,

\*\*\* p<0.001

GCS = Glasgow Coma Scale, GOSE = Glasgow Outcome Scale-Extended, HADS = Hospital Anxiety and Depression Scale PCS = Physical Component Summary; MCS = Mental Component Summary.

<https://doi.org/10.1371/journal.pone.0176668.t006>




**Fig 2. QOLIBRI scale percentage scores (means and 95% confidence intervals) for GOSE groups.**

<https://doi.org/10.1371/journal.pone.0176668.g002>

Positive correlations were found between SF-36 scales and each of the QOLIBRI scales (Table 6), indicating that a higher HRQoL is related to a better subjective health status in the Physical as well as Mental Component Scores. The SF-36 PCS was strongly correlated with the QOLIBRI “Physical Problems” scale, while the association with the “Social Relationships” scale was rather low. On the other hand, the SF-36 MCS was most closely associated with the QOLIBRI “Emotions” scale, while the relationship with “Physical Problems” was weaker.

Figs 2 to 4 present QOLIBRI percentage scores for groups of patients categorized by the GOSE and the HADS. In the present study, a small number of patients suffered from severe anxiety and an even smaller number from severe depression, as assessed on the HADS (i.e. < 10% in both cases). Therefore, these categories were collapsed for both variables, assigning moderate and severe cases to one group each for the anxiety and depression scales. Group-specific percentage scores on the QOLIBRI scales underline the validity of the QOLIBRI.

These results indicate that persons with a severe disability rate their HRQoL significantly worse on all dimensions in comparison to those with good recovery. Also, those with moderate disability experience a significantly worse HRQoL in all scales except for the ‘Social’ and ‘Emotions’ scales.




**Fig 3. QOLIBRI scale percentage scores (means and 95% confidence intervals) for HADS anxiety groups.**

<https://doi.org/10.1371/journal.pone.0176668.g003>

Table 7 shows the relationship between the QOLIBRI scales and measures that were included for the German sample and not described in studies already published. Cognition was assessed using both the TICS and clinician ratings. Significant correlations were found between the TICS and all QOLIBRI scales except the “Social Relationships” scale (see Table 7). The correlations with the TICS indicate that cognitive status is associated with HRQoL particularly on the “Cognition” and “Daily Life and Autonomy” scales. Clinician ratings of cognitive problems are also related to patient-reported HRQoL (Table 7). The ratings of attention and memory dysfunction generally appear to show the strongest relationships with QOLIBRI scales, particularly with the “Cognition” scale, the “Physical Problems” scale, and the QOLIBRI total.

In this German sample, the relationship between self-rated competency/disability and HRQoL after TBI was investigated using the patient version of the PCRS-NR. Robust correlations are seen between the PCRS-NR and all QOLIBRI scales, indicating a strong link between HRQoL and self-rated competence in the three domains measured by this instrument (Table 7). Strong relationships are observed between “Emotional Functioning” and the “Emotions” scale, and between “Cognitive Functioning” and the “Cognition” scale. More generally, strong correlations (all  $r > 0.49$ ) are present between the PCRS-NR total and the individual QOLIBRI scales. The relationship between the PCRS-NR total score and the QOLIBRI total score is particularly strong ( $r = 0.75$ ).



**Fig 4. QOLIBRI scale percentage scores (means and 95% confidence intervals) for HADS depression groups.**

<https://doi.org/10.1371/journal.pone.0176668.g004>

The POMS was included as an assessment of mood states, in addition to the HADS scales. As might be expected, there is a strong correlation between all POMS scales and the QOLIBRI. The strongest relationships between specific scales are between the POMS Depression/dejection scale and the QOLIBRI “Self”, “Emotions”, and total scales.

Correlations between the QOLIBRI scales and the SWLS sum score are shown in Table 7. Significant correlations for all QOLIBRI scales indicate that TBI-specific HRQoL and general wellbeing are closely related. The SWLS shows a particularly strong correlation with the QOLIBRI total score ( $r = 0.74$ ), and the question arises whether there is any difference in the constructs that are being measured.

We studied the overlap between the QOLIBRI total score and the SWLS in more detail using partial correlations. The correlation between the GOSE total score and the QOLIBRI total scale, controlling for the SWLS, remained highly significant ( $r = 0.35$ ;  $p < 0.001$ ), whereas the correlation between SWLS and GOSE was no longer significant after controlling for the QOLIBRI total score ( $r = 0.03$ ;  $p = 0.695$ ). These results indicate that the QOLIBRI provides information which would be lost if solely the SWLS was used. Differences between the SWLS and the QOLIBRI are also seen in the partial correlations using the HADS anxiety scale. The correlation between the HADS anxiety scale score and the QOLIBRI total scale,

**Table 7. Relationships between QOLIBRI scales and measures of cognition, disability, emotional status, and wellbeing.**

|                                     | Cognition | Self     | Daily life & autonomy | Social relationships | Emotions | Physical problems | QOLIBRI total |
|-------------------------------------|-----------|----------|-----------------------|----------------------|----------|-------------------|---------------|
| <b>TICS</b>                         | 0.38***   | 0.31***  | 0.43***               | 0.13                 | 0.32***  | 0.33***           | 0.41***       |
| <b>Clinician Ratings:-</b>          |           |          |                       |                      |          |                   |               |
| Communication problems              | 0.36***   | 0.25***  | 0.27***               | 0.25***              | 0.16*    | 0.23**            | 0.33***       |
| Attention dysfunction               | 0.46***   | 0.34***  | 0.29***               | 0.22**               | 0.30**   | 0.41***           | 0.42***       |
| Memory dysfunction                  | 0.38***   | 0.24**   | 0.31***               | 0.12                 | 0.17*    | 0.39***           | 0.33***       |
| Executive function disorders        | 0.28***   | 0.20**   | 0.23**                | 0.12                 | 0.23**   | 0.22**            | 0.29***       |
| Affective and behavioural Disorders | 0.22**    | 0.20**   | 0.24**                | 0.16*                | 0.25**   | 0.32***           | 0.28***       |
| <b>PCRS:-</b>                       |           |          |                       |                      |          |                   |               |
| Emotional functioning               | 0.55***   | 0.60***  | 0.50***               | 0.53***              | 0.64***  | 0.42***           | 0.66***       |
| Interpersonal functioning           | 0.64***   | .58***   | .50***                | .51***               | .60***   | .43***            | .68***        |
| Cognitive functioning               | 0.65***   | 0.46***  | 0.49***               | 0.30***              | 0.40***  | 0.37***           | 0.56***       |
| PCRS total                          | 0.71***   | 0.65***  | 0.58***               | 0.54***              | 0.67***  | 0.49***           | 0.75***       |
| <b>POMS:-</b>                       |           |          |                       |                      |          |                   |               |
| Depression/ dejection               | -0.54***  | -0.72*** | -0.59***              | -0.58***             | -0.68*** | -0.38***          | -0.72***      |
| Vigour/ activity                    | 0.54***   | 0.59***  | 0.52***               | 0.45***              | 0.49***  | 0.41***           | 0.62***       |
| Fatigue/ inertia                    | -0.49***  | -0.63*** | -0.55***              | -0.55***             | -0.55*** | -0.47***          | -0.65***      |
| Anger/ hostility                    | -0.36***  | -0.50*** | -0.36***              | -0.46***             | -0.52*** | -0.33***          | -0.50***      |
| <b>SWLS</b>                         | 0.52***   | 0.72***  | 0.66***               | 0.64***              | 0.57***  | 0.46***           | 0.74***       |

TICS = Telephone Interview for Cognitive Status; PCRS = Patient Competency Rating Scale; POMS = Profile of Mood States; SWLS = Satisfaction With Life Scale.

\* p<0.05,

\*\* p<0.01,

\*\*\* p<0.001

<https://doi.org/10.1371/journal.pone.0176668.t007>

controlling for the SWLS, was highly significant ( $r = -0.56$ ;  $p < 0.001$ ), but the correlation between the SWLS and the HADS anxiety scale was not significant when controlling for the QOLIBRI total score ( $r = 0.90$ ;  $p = -0.010$ ). However, for the HADS depression scale both partial correlations reached significance ( $r_{\text{SWLS to Depression controlling for QOLIBRI}} = -0.32$ ;  $p < 0.001$ ;  $r_{\text{SWLS to Depression controlling for QOLIBRI}} = -0.55$ ;  $p < 0.001$ ).

## Discussion

Brain injuries are not only associated with bodily symptoms and physical limitations, but also with impairments in cognitive [45,46], social [47,48], and emotional [12,49,50] domains. In addition, many patients experience an altered sense of self [51,52] reduced overall functioning [14], and limited participation [53,54]. However, existing measures of HRQoL often focus on few isolated domains and pay less attention to domains such as cognition and perception of the self [13]. In contrast, the QOLIBRI is a disease-specific instrument that captures multiple aspects of HRQoL after TBI, encompassing physical, psychological (emotional and cognitive), social, self-image, and functional domains [14].

In contrast to the international study, this German-language sample was recruited in emergency and surgical wards of hospitals offering acute care for head injuries. Therefore, there were greater proportions of mild and elderly cases in this sample than in the international study, in which participants were predominantly recruited through rehabilitation centres. The variation in patients and the differences in the degree of severity presents an advantage for

psychometric testing, although the relatively modest number of participants may have a limiting effect on the outcome of some analyses, e.g. the item response theory analyses, factor analyses, and structural equation modelling.

## Properties of the QOLIBRI

For international multi-centre studies, the equivalence of both the content and metrics of the instrument needs to be demonstrated [55–57]. The German translation of the QOLIBRI was based on a standardized procedure, including translation, back-translation, review, and cognitive debriefing [58,59]. Cross-cultural development included the demonstration of comparable metric properties for the whole sample.

**Reliability.** Results indicate that the psychometric properties of the German QOLIBRI are favourable. Internal consistencies and test-retest reliabilities for the individual German QOLIBRI subscales and the total score are good to very good, and they are slightly higher than in the international sample. The QOLIBRI scales are thus appropriate for analyses on a group level and most of them are also sufficiently reliable for assessing individuals. One concern with self-report instruments in TBI is the potential lack of insight that may be experienced by those with cognitive impairment. Experiences with diseases such as dementia suggest that subjective HRQoL judgments can be obtained reliably even in people with substantial cognitive impairments [60,61]. In the German sample, the internal consistency (i.e., Cronbach's  $\alpha$ ) and test-retest reliability for participants with poorer cognitive performance were good to very good (0.79–0.91). The lowest reliability was recorded for the “Physical Problems” scale, an area in which some fluctuation might be expected. These findings indicate that even participants with lower cognitive performance were responding to the questionnaire rationally and consistently. However, we should add that this encouraging result should be considered preliminary as our study did not include a detailed neuropsychological assessment.

When the correlation between neuropsychological functioning and QoL have been investigated in the case of head injuries or other diseases of the central nervous system, the associations have usually been rather weak or absent [62]. It is notable, however, that in this German validation, correlations between the TICS and the QOLIBRI were quite high. One limitation is that the TICS is a rather crude instrument designed for screening cognitive functioning. HRQoL was also related to clinician ratings of cognitive problems, particularly problems of memory and attention. These findings indicate that impaired cognitive status is associated with poor HRQoL, i.e. low QOLIBRI scores. This result is promising and novel in TBI research and suggests a measurable relationship between these two classes of variables, and further research is warranted on this topic.

**Structural validity.** The conceptual model of HRQoL, on which the international QOLIBRI descriptive system was based, suggests a six-dimensional model with four “satisfaction” scales and two “bothered” scales. The German PCA and SEM analyses also support this six-factor structure of the QOLIBRI in a second-order latent variable model. As expected, the factor loadings were a little lower than in the international sample due to the small number of participants. The Rasch analysis also supports the structure of the QOLIBRI. The analysis identified only a small number of items that did not meet relatively conservative criteria for fit to a Rasch scale, and the extent of the deviation in these instances was minor. A cut-off of  $1\pm 0.4$  is sometimes suggested for rating scales [63], and all items met this criterion. The QOLIBRI subscales generally had satisfactory Rasch reliability statistics, but the lower values for the “Emotions” and “Physical Problems” scales suggest that some caution is necessary when using these as individual measures.

**Convergent validity.** Concerning the validity of the German version of the QOLIBRI, the results show the expected pattern of relationships with other scales, thus confirming the validity of the scale. Particular patterns in the findings are consistent with expectations and comparable to, or slightly better than, the international studies (see Table 6): The SF-36 PCS has its highest correlation with the QOLIBRI “Physical Problems Scale” (.74), and a similar pattern is seen for “Comorbid Health Conditions” (.67). The SF-36 MCS correlates most highly with the “Emotions” (.69), and “Self” (.65) scales. The SWLS correlates most highly with “Self” (0.72) and “Daily Life and Autonomy” (0.66) scales. The HADS “Anxiety” scale correlates most strongly with the QOLIBRI “Emotions” scale (0.57), and HADS “Depression” with the “Self” scale (0.63). “Help needed with activities” correlates most with the “Daily Life” (0.63) and “Physical Problems” (0.51) scales. As would be expected, the strongest correlations with the GOSE are for “Physical Problems” (.56) and “Daily Life” (.51).

The SWLS and the QOLIBRI total score were strongly correlated in the sample, indicating substantial overlap in the constructs assessed. Both the SWLS and the QOLIBRI are satisfaction scales, and would thus be expected to overlap. The SWLS has been recommended as a core QoL measure for assessing TBI [64], and the question arises whether the QOLIBRI total adds anything to this well-established measure. Analysis using partial correlations with the GOSE demonstrates that the QOLIBRI contains information that is not captured by the SWLS. The relationships found with the GOSE are consistent with the idea that the SWLS reflects satisfaction with life in general, and the QOLIBRI more specifically tracks satisfaction with areas of life that are affected by a brain injury.

Another notable finding is the impact of disability, even moderate disability, on generic and TBI-specific HRQoL, as reported by the participants. The results of the comparison with the GOSE show that in each domain of the QOLIBRI, severely and moderately disabled TBI participants indicate lower HRQoL than those who had achieved a good level of recovery. It is particularly noteworthy that patients with moderate disabilities reported significantly lower HRQoL than those with a good recovery. It is still not always acknowledged, however, that even those with apparently mild injuries can suffer from moderate disability [65].

In the German as well as the international sample, the overall relationship between the GOSE and the QOLIBRI was only moderate, indicating that people can have a poor outcome on the GOSE and yet have good HRQoL, and vice versa. A new finding is that self-rated competency on the PCRS is quite strongly related to the QOLIBRI [66]. This supports the idea that the QOLIBRI is sensitive to common types of disability caused by head injury. There has been some debate over the relationship between functional outcome and HRQoL, with some authors claiming that the two are only weakly related. Brown and Gordon [67] have studied the relationship between disablement and quality of life, and found that less than 20% of the variance could typically be predicted by measures of disability. In contrast, the correlation between the total scores of the PCRS and the QOLIBRI here suggests that shared variance is over 50%. Brown and Gordon [67] argue that the measures of disablement with greater face validity are better predictors of QoL. In the present study, the person’s own view of their competence in functioning seems to be particularly strongly related to HRQoL.

The GOS/GOSE is currently the most popular outcome measure in acute brain injury trials; however, it is widely acknowledged that this assessment has limitations. It does not address some important domains, including cognition or self-perception [68,69]. As an assessment of functional outcome, it does not capture the subjective experience and self-report by the individual. In the field of brain injury, the interest in HRQoL has been partly encouraged by the failure of clinical trials in the acute stage to demonstrate clinical benefits using functional outcomes as a primary endpoint [70]. The development of this disease-specific HRQoL scale after TBI may therefore provide a good end point for clinical trials. It opens the possibility of



constructing a composite multidimensional outcome assessment that covers both functional and HRQoL outcomes. Such a composite assessment would complement the picture of outcome after brain injury, and potentially provide a more sensitive tool for clinical trials.

Examining predictors of HRQoL, a regression analysis with the German data showed similar findings to those of the international sample, whereby the strongest predictors of QOLIBRI scores are aspects of the patient's current status, and specifically emotional state, comorbid health conditions, and functional outcome, whereas demographic and clinical background factors have little if any influence. Given the average length of time since injury, it is not surprising that GCS played little if any role, and the current findings are in accordance with the work of others [14,45].

Previous reports of an association between QoL and emotional state [12,71–74], and loss of independence [75–77] are highlighted in our study in more detail. This has been underlined by the correlations between HADS anxiety and depression, POMS and self-reported health status symptoms, and the QOLIBRI. The association between depression and HRQoL is a ubiquitous finding, also reported in most other diseases, but it is generally acknowledged that HRQoL goes beyond simply assessment of emotional state [78,79].

## Limitations

The current study also has some limitations. Its intention was to validate the QOLIBRI in the German language, and it was therefore desirable to collect data from a large sample of brain injuries with a wide variety of clinical characteristics. The current cross-sectional design of the study was not designed to allow a straightforward analysis of recovery over time. In addition, given the wide range of intervals after the injury in our sample, our results do not allow inference of QoL after a certain mean interval, but are more indicative of “general” QoL after TBI. Another issue yet to be resolved by future research is the responsiveness of the QOLIBRI to change and susceptibility to response shift. The results show that the German QOLIBRI is sensitive to differences in outcome on the GOSE and other functional and patient characteristics. Moreover, due to the scope of our study it was impossible to investigate the impact of impairment of self-awareness or cognition on generic and TBI-specific HRQoL self-rating in more detail. However, relying on the instruments we used we were able to establish, for the German as well as for the total sample, that level of cognitive function did not affect the reliability of reporting [22,23]. For detailed information on the impact of impaired self-awareness on HRQoL see [66,80].

## Conclusion & outlook

In conclusion, the psychometric properties of the German QOLIBRI suggest that it is a practical and reliable instrument that can be considered for use in studies examining HRQoL after TBI in clinical as well as in research settings. The QOLIBRI offers a profile of HRQoL in domains that are relevant for TBI, as well as a summary score. It provides an assessment of TBI-specific HRQoL that potentially complements functional outcome measures such as the GOS or GOSE and measures of social participation, general wellbeing and subjective health status.

Development of this instrument was encouraged by many people working in post-acute brain injury management, and especially in rehabilitation. Currently the English-language version of the QOLIBRI has been downloaded more than 1000 times by research and rehabilitation centres; project outlines describe the use of the QOLIBRI, for example, to measure the effectiveness of cognitive and functional therapy. Outcome measures assessing TBI-specific HRQoL may contribute to quality control in short- and long-term care, medical decision-making, and rehabilitation planning. In clinical use in the rehabilitation setting, it can help to

identify and set appropriate goals for therapy. The QOLIBRI covers several domains currently missing from TBI-specific item bank initiatives such as PROMIS or NEURO-QOL, and these systems may profit from including items from the QOLIBRI [81].

The final descriptive system of the German QOLIBRI provides both a comprehensive HRQoL profile across six domains of life, and a total index of HRQoL after TBI. When used in evaluation studies, shifts in individual domains will reflect areas of life where gains or losses following intervention or treatment are noticeable. In contrast, where an index is required, the QOLIBRI total score can be used to assess the impact of treatments on HRQoL. In addition, where a very short instrument is needed, we have developed an overall screening measure, the German QOLIBRI-OS, with six items and good psychometric properties covering the most important areas of HRQoL [82].

## Supporting information

**S1 File. Raw data.**  
(POR)

## Acknowledgments

The QOLIBRI Task Force consists of collaborating investigators in the following countries (\*Steering Committee, ‡Methodological Group): Argentina (Armando Basso), Australia (Graeme Hawthorne‡), Austria (Stefan Hoefler‡), Belgium (Christine Croisau, Andrew Maas\*), Brazil (Lucia Braga), China (Wai Poon, Zhang Tong), Denmark (Anne-Lise Christensen), Finland (Jaana Sarajuuri, Sanna Koskinen), France (Philippe Azouvi, Michelle Montreuil, Pierre North, Jean-Luc Truelle\*), Germany (Monika Bullinger\*, Henning Gibbons‡, Holger Mühlen‡, Edmund Neugebauer\*, Tanja Lischetzke‡, Nadine Sasse, Silke Schmidt, Nicole von Steinbüchel\*‡, Klaus von Wild‡\*), Greece (Eva Tazopoulou), Eco Wahjoe- Pramono (Indonesia), Italy (Rita Formisano), Japan (Izumi Kondo), The Netherlands (Wilbert Bakx, Peter Vos), Portugal (Sandra Guerreiro), Russia (Boleslav Lichterman), Taiwan (Wen-Ta Chiu), United Kingdom (Richard Greenwood, Jane Powell\*, Lindsay Wilson‡), United States of America (John DaVanzo, George Zitnay\*).

*This project was funded by ZNS—Hannelore Kohl Foundation (project number 2008014; [www.hannelore-kohl-stiftung.de](http://www.hannelore-kohl-stiftung.de)). We acknowledge support by the German Research Foundation and the Open Access Publication Funds of the Göttingen University.*

## Author Contributions

**Conceptualization:** NvS.

**Data curation:** NvS.

**Formal analysis:** RGLR CO HG.

**Funding acquisition:** NvS.

**Investigation:** NS RB WD RMO W. Puschendorf W. Petereit VR HS SS KMS KvW.

**Methodology:** NvS.

**Project administration:** NvS.

**Resources:** NvS.

**Supervision:** NvS.

**Visualization:** RGLR.

**Writing – original draft:** NvS CO HG.

**Writing – review & editing:** NvS RGLR NS LW RM RB WD RMO W. Puschendorf W. Peter-eit VR HS SS KMS KvW.

## References

- Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet*. 1975; 1: 480–484. [https://doi.org/10.1016/S0140-6736\(75\)92830-5](https://doi.org/10.1016/S0140-6736(75)92830-5) PMID: 46957
- Shukla D, Devi BI, Agrawal A. Outcome measures for traumatic brain injury. *Clin Neurol Neurosurg*. 2011; 113: 435–441. <https://doi.org/10.1016/j.clineuro.2011.02.013> PMID: 21440363
- Corrigan JD, Bogner J. Latent factors in measures of rehabilitation outcomes after traumatic brain injury. *J Head Trauma Rehabil*. 2004; 19: 445–458. PMID: 15602308
- McCarthy ML, Dikmen SS, Langlois JA, Selassie AW, Gu JK, Horner MD. Self-Reported Psychosocial Health Among Adults With Traumatic Brain Injury. *Arch Phys Med Rehabil*. 2006; 87: 953–961. <https://doi.org/10.1016/j.apmr.2006.03.007> PMID: 16813783
- Lippert-Grüner M, Maegele M, Haverkamp H, Klug N, Wedekind C. Health-related quality of life during the first year after severe brain trauma with and without polytrauma. *Brain Inj*. 2007; 21: 451–455. <https://doi.org/10.1080/02699050701343961> PMID: 17522984
- Hawthorne G, Gruen R, Kaye AH. Traumatic brain injury and long-term quality of life: Findings from an Australian study. *J Neurotrauma*. 2009; 26: 1623–1633. <https://doi.org/10.1089/neu.2008-0735> PMID: 19317590
- Klonoff PS, Costa LD, Snow WG. Predictors and indicators of quality of life in patients with Closed-Head injury. *J Clin Exp Neuropsychol*. 1986; 8: 469–485. <https://doi.org/10.1080/01688638608405171> PMID: 3805248
- Shames J, Treger I, Ring H, Giaquinto S. Return to work following traumatic brain injury: Trends and challenges. *Disabil Rehabil*. 2007; 29: 1387–1395. <https://doi.org/10.1080/09638280701315011> PMID: 17729084
- Nakase-Richardson R, Sherer M, Seel RT, Hart T, Hanks R, Arango-Lasprilla JC, et al. Utility of post-traumatic amnesia in predicting 1-year productivity following traumatic brain injury: Comparison of the Russell and Mississippi PTA classification intervals. *J Neurol Neurosurg Psychiatry*. 2011; 82: 494–499. <https://doi.org/10.1136/jnnp.2010.222489> PMID: 21242285
- Powell JH, Beckers K, Greenwood RJ. Measuring progress and outcome in community rehabilitation after brain injury with a new assessment instrument—the BICRO-39 scales. 1998; 79: 1213–1225.
- Arango-Lasprilla JC, Rosenthal M, Deluca J, Komaroff E, Sherer M, Cifu D, et al. Traumatic brain injury and functional outcomes: does minority status matter? *Brain Inj*. 2007; 21: 701–708. <https://doi.org/10.1080/02699050701481597> PMID: 17653944
- Diaz AP, Schwarzbald ML, Thais ME, Hohl A, Bertotti MM, Schmoeller R, et al. Psychiatric Disorders and Health-Related Quality of Life after Severe Traumatic Brain Injury: A Prospective Study. *J Neurotrauma*. 2012; 29: 1029–1037. <https://doi.org/10.1089/neu.2011.2089> PMID: 22111890
- Koskinen S, Hokkinen E-M, Wilson L, Sarajuuri J, von Steinbüchel N, Truelle J-L. Comparison of subjective and objective assessments of outcome after traumatic brain injury using the International Classification of Functioning, Disability and Health (ICF). *Disabil Rehabil*. 2011; 33: 2464–2478. <https://doi.org/10.3109/09638288.2011.574776> PMID: 21534850
- von Steinbüchel N, Richter S, Morawetz C, Riemsma R. Assessment of subjective health and health-related quality of life in persons with acquired or degenerative brain injury. *Curr Opin Neurol*. 2005; 18: 681–691. <https://doi.org/10.1097/01.wco.0000194140.56429.75> PMID: 16280680
- McMahon PJ, Hricik A, Yue JK, Puccio AM, Inoue T, Lingsma HF, et al. Symptomatology and Functional Outcome in Mild Traumatic Brain Injury: Results from the Prospective TRACK-TBI Study. *J Neurotrauma*. 2014; 31: 26–33. <https://doi.org/10.1089/neu.2013.2984> PMID: 23952719
- Hütter BO, Gilsbach JM. Background and first results about methodological characteristics of the Aachen Life Quality Inventory. *Zentralblatt Für Neurochir*. 2001; 62: 37–42. <https://doi.org/10.1055/s-2002-19476> PMID: 11786934
- Ahmadi SA, Meier U, Lemcke J. Detailed long-term outcome analysis after decompressive craniectomy for severe traumatic brain injury. *Brain Inj*. 2010; 24: 1539–1549. <https://doi.org/10.3109/02699052.2010.523049> PMID: 20973624

18. Findler Marianne, Cantor Joshua, Li. The reliability and validity of the SF-36 health survey questionnaire for use with individuals with traumatic brain injury. *Brain Inj.* 2001; 15: 715–723. <https://doi.org/10.1080/02699050010013941> PMID: 11485611
19. WHOQOL group. The World Health Organization Quality of Life Assessment (WHOQOL): development and general psychometric properties. *Soc Sci Med* 1982. 1998; 46: 1569–1585.
20. Ware JE, Sherbourne CD. The MOS 36-Item Short-Form Health Survey (SF-36): I. Conceptual Framework and Item Selection. *Med Care.* 1992; 30: 473–483. PMID: 1593914
21. Bergner M. Development, testing, and use of the Sickness Impact Profile. In: Walker SR, Rosser R, editors. *Quality of life, assessment and application.* Lancaster and Boston: MTP Press; 1988. pp. 79–94.
22. von Steinbüchel N, Wilson L, Gibbons H, Hawthorne G, Höfer S, Schmidt S, et al. Quality of Life after Brain Injury (QOLIBRI): Scale Validity and Correlates of Quality of Life. *J Neurotrauma.* 2010; 27: 1157–1165. <https://doi.org/10.1089/neu.2009.1077> PMID: 20210602
23. von Steinbüchel N, Wilson L, Gibbons H, Hawthorne G, Höfer S, Schmidt S, et al. Quality of Life after Brain Injury (QOLIBRI): Scale Development and Metric Properties. 2010; 27: 1167–1185. <https://doi.org/10.1089/neu.2009.1076> PMID: 20486801
24. Wilson L, Pettigrew LEL, Teasdale G. Structured Interviews for the Glasgow Outcome Scale and the Extended Glasgow Outcome Scale: Guidelines for Their Use. 1998; 15: 573–585. <https://doi.org/10.1089/neu.1998.15.573> PMID: 9726257
25. Teasdale G, Jennett B. Assessment of coma and impaired consciousness: A practical scale. *The Lancet.* 1974; 2: 81–84. [https://doi.org/10.1016/S0140-6736\(74\)91639-0](https://doi.org/10.1016/S0140-6736(74)91639-0)
26. World Medical Association. Declaration of Helsinki. Ethical principles for medical research involving human subjects. 2009;
27. Diener E, Emmons RA, Larsen RJ, Griffin S. The Satisfaction With Life Scale. *J Pers Assess.* 1985; 49: 71–75. [https://doi.org/10.1207/s15327752jpa4901\\_13](https://doi.org/10.1207/s15327752jpa4901_13) PMID: 16367493
28. Borgaro SR, Prigatano GP. Modification of the Patient Competency Rating Scale for use on an acute neurorehabilitation unit: The PCRS-NR. *Brain Inj.* 2003; 17: 847–853. <https://doi.org/10.1080/0269905031000089350> PMID: 12963551
29. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand.* 1983; 67: 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x> PMID: 6880820
30. Albani C, Blaser G, Geyer M, Schmutzer G, Brähler E, Bailer H, et al. Überprüfung der Gütekriterien der deutschen Kurzform des Fragebogens “Profile of Mood States” (POMS) in einer repräsentativen Bevölkerungsstichprobe. *Psychother—Psychosom—Med Psychol.* 2005; 55: 324–330. <https://doi.org/10.1055/s-2004-834727> PMID: 15986282
31. Brandt J, Spencer M, Folstein M. The Telephone Interview for Cognitive Status. *Neuropsychiatry Neuropsychol Behav Neurol.* 1988; 1: 111–117.
32. von Steinbüchel N, Lischetzke T, Gurny M, Eid M. Assessing quality of life in older people: Psychometric properties of the WHOQOL-BREF. *Eur J Ageing.* 2006; 3: 116–122. <https://doi.org/10.1007/s10433-006-0024-2>
33. Masters GN. A rasch model for partial credit scoring. *Psychometrika.* 1982; 47: 149–174. <https://doi.org/10.1007/BF02296272>
34. Schermelleh-Engel K, Moosbrugger H, Müller H. Evaluating the Fit of Structural Equation Models: Tests of Significance and Descriptive Goodness-of-Fit Measures. *Methods Psychol Res Online.* 2003; 8: 23–27.
35. Steiger JH. Structural Model Evaluation and Modification: An Interval Estimation Approach. *Multivar Behav Res.* 1990; 25: 173–180. [https://doi.org/10.1207/s15327906mbr2502\\_4](https://doi.org/10.1207/s15327906mbr2502_4) PMID: 26794479
36. Bentler PM. Comparative fit indexes in structural models. *Psychol Bull.* 1990; 107: 238–246. PMID: 2320703
37. Bentler PM. EQS structural equations program manual [Internet]. Encino, Calif.: Multivariate Software; 1995. <http://www.econ.upf.edu/satorra/CourseSEMVVienna2010/EQSManual.pdf>
38. Browne M, Cudeck R. Alternative ways of assessing model fit. In: Bollen KA, Long SJ, editors. *Testing structural equation.* Newbury Park (Calif.): Sage Publications; 1993. pp. 136–162.
39. Hu L, Bentler P. Evaluating model fit. In: Hoyle RH, editor. *Structural equation modeling.* Thousand Oaks: Sage Publications; 1995. pp. 76–99.
40. Moosbrugger H, Kelava A. *Testtheorie und Fragebogenkonstruktion.* Heidelberg: Springer; 2007.
41. Bland JM, Altman DG. Statistics notes: Cronbach's alpha. *Br Med J.* 1997; 314: 572. <https://doi.org/10.1136/bmj.314.7080.572>
42. Fleiss JL. Analysis of data from multiclinic trials. *Control Clin Trials.* 1986; 7: 267–275. [https://doi.org/10.1016/0197-2456\(86\)90034-6](https://doi.org/10.1016/0197-2456(86)90034-6) PMID: 3802849

43. Brandt J, Folstein MF. Telephone Interview for Cognitive Status: Professional Manual. Lutz, FL.: Psychological Assessment Resources, Inc.; 2003.
44. Conover WJ, Iman RL. Rank Transformations as a Bridge between Parametric and Nonparametric Statistics. *Am Stat*. 1981; 35: 124–129. <https://doi.org/10.1080/00031305.1981.10479327>
45. Dikmen SS, Machamer JE, Powell JM, Temkin NR. Outcome 3 to 5 years after moderate to severe traumatic brain injury. *Arch Phys Med Rehabil*. 2003; 84: 1449–1457. [https://doi.org/10.1016/S0003-9993\(03\)00287-9](https://doi.org/10.1016/S0003-9993(03)00287-9) PMID: 14586911
46. Whiting MD, Baranova AI, Hamm RJ. Cognitive Impairment following Traumatic Brain Injury. In: Levin ED, Buccafusco JJ, editors. *Animal models of cognitive impairment*. Boca Raton: CRC/Taylor & Francis; 2006.
47. Mazaux J-M, Masson F, Levin HS, Alaoui P, Maurette P, Barat M. Long-term neuropsychological outcome and loss of social autonomy after traumatic brain injury. 1997; 78: 1316–1320. [https://doi.org/10.1016/S0003-9993\(97\)90303-8](https://doi.org/10.1016/S0003-9993(97)90303-8)
48. Andelic N, Sigurdardottir S, Schanke A-K, Sandvik L, Sveen U, Roe C. Disability, physical health and mental health 1 year after traumatic brain injury. *Disabil Rehabil*. 2010; 32: 1122–1131. <https://doi.org/10.3109/09638280903410722> PMID: 20113311
49. Hoofien Dan, Gilboa Assaf, Vaki Eli. Traumatic brain injury (TBI) 10–20 years later: a comprehensive outcome study of psychiatric symptomatology, cognitive abilities and psychosocial functioning. *Brain Inj*. 2001; 15: 189–209. <https://doi.org/10.1080/026990501300005659> PMID: 11260769
50. Hart T, Brenner L, Clark AN, Bogner JA, Novack TA, Chervoneva I, et al. Major and Minor Depression After Traumatic Brain Injury. *Arch Phys Med Rehabil*. 2011; 92: 1211–1219. <https://doi.org/10.1016/j.apmr.2011.03.005> PMID: 21807140
51. Prigatano GP. Personality disturbances associated with traumatic brain injury. *J Consult Clin Psychol*. 1992; 60: 360–368. <https://doi.org/10.1037/0022-006X.60.3.360> PMID: 1619090
52. Riley GA, Dennis RK, Powell T. Evaluation of coping resources and self-esteem as moderators of the relationship between threat appraisals and avoidance of activities after traumatic brain injury. *Neuropsychol Rehabil*. 2010; 20: 869–882. <https://doi.org/10.1080/09602011.2010.503041> PMID: 20665337
53. Ponsford J, Olver J, Ponsford M, Nelms R. Long-term adjustment of families following traumatic brain injury where comprehensive rehabilitation has been provided. *Brain Inj*. 2003; 17: 453–468. <https://doi.org/10.1080/0269905031000070143> PMID: 12745702
54. Laxe S, Zasler N, Tschiesner U, López-Blazquez R, Tormos JM, Bernabeu M. ICF use to identify common problems on a TBI neurorehabilitation unit in Spain. *NeuroRehabilitation*. 2011; 29: 99–110. <https://doi.org/10.3233/NRE-2011-0683> PMID: 21876302
55. Bullinger M, Anderson R, Cella D, Aaronson N. Developing and evaluating cross-cultural instruments from minimum requirements to optimal models. *Qual Life Res*. 1993; 2: 451–459. <https://doi.org/10.1007/BF00422219> PMID: 8161979
56. Stucki G, Stucki S, Sangha O. Patienten-zentrierte Evaluation der Krankheitsauswirkungen bei muskuloskelettalen Erkrankungen: Adaptation und Neuentwicklung von Outcome-Instrumenten (Patients-oriented outcome assessment in musculoskeletal disease: cross-cultural adaptation and development of instruments). *Z Für Rheumatol*. 1997; 56: 266–275. <https://doi.org/10.1007/s003930050041>
57. Riemsmas RP, Forbes CA, Glanville JM, Eastwood AJ, Kleijnen J. General health status measures for people with cognitive impairment: Learning disability and acquired brain injury. *Health Technol Assess*. 2001; 5: 1–100. <https://doi.org/10.3310/hta5060>
58. Acquadro C, Jambon B, Ellis D, and Marquis P. Language and translation issues. In: Spilker B, editor. *Quality of life and pharmacoeconomics in clinical trials*. Philadelphia: Lippincott-Raven; 1996. pp. 575–585.
59. Steinbüchel N von, Petersen C, Bullinger M. Study Protocol, Cognitive Debriefing Protocol and Harmonization Guidelines for the QOLIBRI Development in TBI Study. 2002.
60. Novella JL, Jochum C, Ankri J, Morrone I, Jolly D, Blanchard F. Measuring general health status in dementia: Practical and methodological issues in using the SF-36. *Aging Clin Exp Res*. 2001; 13: 362–369. <https://doi.org/10.1007/BF03351504>
61. Wlodarczyk JH, Brodaty H, Hawthorne G. The relationship between quality of life, Mini-Mental State Examination, and the Instrumental Activities of Daily Living in patients with Alzheimer's disease. *Arch Gerontol Geriatr*. 2004; 39: 25–33. <https://doi.org/10.1016/j.archger.2003.12.004> PMID: 15158578
62. von Steinbüchel N. *Gesundheitsbezogene Lebensqualität und mentale Funktionen*. Habilitationsschrift im Fach Psychologie an der Naturwissenschaftlichen Fakultät der Leopold-Franzens-Universität. Innsbruck; 1996.
63. Bond TG, Fox CM. *Applying the Rasch model: Fundamental measurement in the human sciences*. 2nd ed. Mahwah, N.J.: Lawrence Erlbaum Associates Publishers; 2007.

64. Wilde EA, Whiteneck GG, Bogner J, Bushnik T, Cifu DX, Dikmen S, et al. Recommendations for the Use of Common Outcome Measures in Traumatic Brain Injury Research. *Arch Phys Med Rehabil.* 2010; 91: 1650–1660.e17. <https://doi.org/10.1016/j.apmr.2010.06.033> PMID: 21044708
65. Thornhill S, Teasdale GM, Murray GD, McEwen J, Roy CW, Penny KI. Disability in young people and adults one year after head injury: prospective cohort study. *Br Med J Clin Res Ed.* 2000; 320: 1631–1635. <https://doi.org/10.1136/bmj.320.7250.1631>
66. Sasse N, Gibbons H, Wilson L, Martinez-Olivera R, Schmidt H, Hasselhorn M, et al. Self-Awareness and Health-Related Quality of Life After Traumatic Brain Injury. *J Head Trauma Rehabil.* 2013; 28: 464–472. <https://doi.org/10.1097/HTR.0b013e318263977d> PMID: 22935572
67. Brown M, Gordon WA. Empowerment in measurement: “muscle,” “voice,” and subjective quality of life as a gold standard 11No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the author(s) or upon any organization with which the author(s) is/are associated. *Arch Phys Med Rehabil.* 2004; 85: 13–20. <https://doi.org/10.1016/j.apmr.2003.08.110>
68. Berger E, Leven F, Pirente N, Bouillon B, Neugebauer E. Quality of Life after traumatic brain injury: A systematic review of the literature. *Restor Neurol Neurosci.* 1999; 14: 93–102.
69. Dijkers MP. Quality of life after traumatic brain injury: A review of research approaches and findings 11No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the author(s) or upon any organization with which the author(s) is/are associated. *Arch Phys Med Rehabil.* 2004; 85: 21–35. <https://doi.org/10.1016/j.apmr.2003.08.119>
70. Narayan RK, Michel ME, Ansell B, Baethmann A, Biegon A, Bracken MB, et al. Clinical Trials in Head Injury. *J Neurotrauma.* 2002; 19: 503–557. <https://doi.org/10.1089/089771502753754037> PMID: 12042091
71. Koskinen S. Quality of life 10 years after a very severe traumatic brain injury (TBI): The perspective of the injured and the closest relative. *Brain Inj.* 1998; 12: 631–648. <https://doi.org/10.1080/026990598122205> PMID: 9724835
72. Corrigan JD, Bogner JA, Mysiw WJ, Clinchot D, Fugate L. Life satisfaction after traumatic brain injury. *J Head Trauma Rehabil.* 2001; 16: 543–555. PMID: 11732970
73. Underhill AT, Lobello SG, Stroud TP, Terry KS, Devivo MJ, Fine PR. Depression and life satisfaction in patients with traumatic brain injury: A longitudinal study. *Brain Inj.* 2003; 17: 973–982. <https://doi.org/10.1080/0269905031000110418> PMID: 14514448
74. Guillamondegui OD, Montgomery SA, Phibbs FT, McPheeters ML, Alexander PT, Jerome RN, et al. Traumatic Brain Injury and Depression [Internet]. Rockville, MD: Agency for Healthcare Research and Quality; 2011. [www.effectivehealthcare.ahrq.gov/reports/final.cfm](http://www.effectivehealthcare.ahrq.gov/reports/final.cfm)
75. Mailhan L, Azouvi P, Dazard A. Life satisfaction and disability after severe traumatic brain injury. *Brain Inj.* 2005; 19: 227–238. <https://doi.org/10.1080/02699050410001720149> PMID: 15832869
76. Lane AK, Benoit D. Driving, brain injury and assistive technology. *NeuroRehabilitation.* 2011; 28: 221–229. <https://doi.org/10.3233/NRE-2011-0651> PMID: 21558628
77. Bottari C, Gosselin N, Guillemette M, Lamoureux J, Ptitto A. Independence in managing one’s finances after traumatic brain injury. *Brain Inj.* 2011; 25: 1306–1317. <https://doi.org/10.3109/02699052.2011.624570> PMID: 22077536
78. Gladis MM, Gosch EA, Dishuk NM, Crits-Christoph P. Quality of life: Expanding the scope of clinical significance. *J Consult Clin Psychol.* 1999; 67: 320–331. <https://doi.org/10.1037/0022-006X.67.3.320> PMID: 10369052
79. Höfer S, Benzer W, Alber H, Ruttman E, Kopp M, Schussler G, et al. Determinants of Health-Related Quality of Life in Coronary Artery Disease Patients: A Prospective Study Generating a Structural Equation Model. *Psychosomatics.* 2005; 46: 212–223. <https://doi.org/10.1176/appi.psy.46.3.212> PMID: 15883142
80. Formisano R, Longo E, Azicnuda E, Silvestro D, D’Ippolito M, Truelle J-L, et al. Quality of life in persons after traumatic brain injury as self-perceived and as perceived by the caregivers. *Neurol Sci.* 2017; 38: 279–286. <https://doi.org/10.1007/s10072-016-2755-y> PMID: 27826793
81. Carlozzi NE, Tulskey DS, Kisala PA. Traumatic Brain Injury Patient-Reported Outcome Measure: Identification of Health-Related Quality-of-Life Issues Relevant to Individuals With Traumatic Brain Injury. *Arch Phys Med Rehabil.* 2011; 92: 52–60. <https://doi.org/10.1016/j.apmr.2010.12.046> PMID: 21958923
82. von Steinbuechel N, Wilson L, Gibbons H, Muehlan H, Schmidt H, Schmidt S, et al. QOLIBRI overall scale: a brief index of health-related quality of life after traumatic brain injury. *J Neurol Neurosurg Psychiatry.* 2012; 83: 1041–1047. <https://doi.org/10.1136/jnnp-2012-302361> PMID: 22851609