


CORRECTION

Correction: Role of Interleukin-10 on Nasal Polypogenesis in Patients with Chronic Rhinosinusitis with Nasal Polyps

The *PLOS ONE* Staff

The image for Fig 8 is incorrectly duplicated in [Fig 1](#). Please see the correct image for [Fig 1](#) here. The publisher apologizes for the error.



 OPEN ACCESS

Citation: The *PLOS ONE* Staff (2017) Correction: Role of Interleukin-10 on Nasal Polypogenesis in Patients with Chronic Rhinosinusitis with Nasal Polyps. *PLoS ONE* 12(5): e0177755. <https://doi.org/10.1371/journal.pone.0177755>

Published: May 11, 2017

Copyright: © 2017 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 1. Expression of IL-10 among the different groups. A, mRNA expression of IL-10 in the UT tissue or NP tissue among groups. B, representative images of western blot assay and relative expression of IL-10 in the UT or NP among the groups. C, representative images of IL-10 immunohistochemistry and the number of IL-10 positive cells in the tissues from the different groups. CRSsNP, chronic rhinosinusitis without NP; CRSwNP, chronic rhinosinusitis with NP. * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$.

<https://doi.org/10.1371/journal.pone.0177755.g001>

Reference

- Xu J, Han R, Kim DW, Mo J-H, Jin Y, Rha K-S, et al. (2016) Role of Interleukin-10 on Nasal Polypogenesis in Patients with Chronic Rhinosinusitis with Nasal Polyps. PLoS ONE 11(9): e0161013. doi:10.1371/journal.pone.0161013 PMID: 27584662