

RESEARCH ARTICLE

# Egg white hydrolysate shows insulin mimetic and sensitizing effects in 3T3-F442A pre-adipocytes

Forough Jahandideh<sup>1,2</sup>, Subhadeep Chakrabarti<sup>1,2</sup>, Sandra T. Davidge<sup>2,3,4,5</sup>, Jianping Wu<sup>1,2\*</sup>

**1** Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, Alberta, Canada, **2** Cardiovascular Research Centre, University of Alberta, Edmonton, Alberta, Canada, **3** Department of Obstetrics and Gynecology, University of Alberta, Edmonton, Alberta, Canada, **4** Department of Physiology, University of Alberta, Edmonton, Alberta, Canada, **5** Women and Children's Health Research Institute, University of Alberta, Edmonton, Alberta, Canada

\* [jwu3@ualberta.ca](mailto:jwu3@ualberta.ca)



**OPEN ACCESS**

**Citation:** Jahandideh F, Chakrabarti S, Davidge ST, Wu J (2017) Egg white hydrolysate shows insulin mimetic and sensitizing effects in 3T3-F442A pre-adipocytes. PLoS ONE 12(10): e0185653. <https://doi.org/10.1371/journal.pone.0185653>

**Editor:** Nobuyuki Takahashi, Tokyo University of Agriculture, JAPAN

**Received:** May 29, 2017

**Accepted:** September 16, 2017

**Published:** October 3, 2017

**Copyright:** © 2017 Jahandideh et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information file.

**Funding:** This work was supported by grants from Alberta Livestock and Meat Agency (ALMA), Egg Farmers of Canada, Egg Farmers of Alberta, and the Natural Sciences and Engineering Research Council (NSERC) to JW. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Abstract

Insulin resistance and inflammation in adipose tissue is a key mechanism underlying metabolic syndrome, a growing health problem characterized by diabetes, obesity and hypertension. Previous work from our research group has demonstrated the potential of egg white ovotransferrin derived bioactive peptides against hypertension, oxidative stress and inflammation *in vitro* and *in vivo*. Egg white hydrolysate (EWH) has also shown anti-hypertensive effects in spontaneously hypertensive rats. Given the interplay among hypertension, inflammation, oxidative stress and metabolic syndrome, the objective of the study was to test the EWH on differentiation, insulin signaling and inflammatory responses in 3T3-F442A pre-adipocytes. Our study suggested that EWH could promote adipocyte differentiation as shown by increased lipid accumulation, increased release of adiponectin and upregulation of peroxisome proliferator associated receptor gamma (PPAR $\gamma$ ) and CCAAT/enhancer binding protein alpha (C/EBP- $\alpha$ ). In addition to enhanced insulin effects on the upregulation of protein kinase B/Akt phosphorylation, EWH treatment increased extracellular signal regulated kinase 1/2 (ERK1/2) phosphorylation to a level similar to that of insulin, indicating insulin sensitizing and mimetic properties of the EWH. EWH further attenuated cytokine induced inflammatory marker; cyclooxygenase -2 (COX-2) by 48.78%, possibly through the AP-1 pathway by down regulating c-Jun phosphorylation in adipocytes. Given the critical role of adipose in the pathogenesis of insulin resistance and metabolic syndrome, EWH may have potential applications in the prevention and management of metabolic syndrome and its complications.

## Introduction

Metabolic syndrome, a combination of several abnormalities that increase the risk for type II diabetes and atherosclerosis is global health problem of growing concern [1–3]. It consists of

atherogenic dyslipidemia (elevated triglycerides and low high-density lipoproteins), hypertension, glucose intolerance, and proinflammatory states [4].

Hypertension and insulin resistance are the key features of metabolic syndrome. Renin-angiotensin system (RAS), the classical pathway for controlling blood pressure and fluid balance, has also a role in the pathogenesis of metabolic syndrome [5]. In the RAS, angiotensin-converting enzyme (ACE) plays a critical role in the formation of angiotensin II (Ang II), the primary active peptide of this system which increases blood pressure by enhancing vascular constriction. RAS blockade by ACE-inhibitors or angiotensin receptor blockers beneficially affects insulin sensitivity and prevents the development of diabetes [6,7]. Given the role of RAS impairments in the pathogenesis of hypertension and metabolic syndrome, there is much interest in developing novel therapies that can target the common pathologies to hypertension and insulin resistance in more complicated disease conditions [2,8,9].

Insulin is essential for normal metabolic functions of various tissues in the body [10]. Adipose tissue with a central role in lipid and glucose metabolism is a key target of insulin [11]. Insulin promotes differentiation of pre-adipocytes into mature adipocytes; a process accompanied by incorporation of lipid droplets and upregulation of immunomodulatory proteins like peroxisome proliferator-associated receptor gamma (PPAR $\gamma$ ) [12]. Collectively, insulin actions on adipose tissue appear to be beneficial and anti-inflammatory in nature. Under metabolic syndrome, insulin signaling in adipose tissue is perturbed, associated with insulin resistance and chronic inflammation [9,13–16]. As such, there is significant interest in developing therapeutic agents to improve insulin signaling in adipocytes, either by insulin sensitizing agents or through agents mimicking insulin actions [17,18]. The insulin sensitizing drugs thiazolidinediones (TZDs) enhance adipocyte differentiation. This increases lipid partitioning into adipocytes and decreases circulating, hepatic, and intramuscular triglycerides thus enhances insulin sensitivity [19].

Use of pharmacological drugs for controlling different complications of metabolic syndrome is associated with significant risk of side-effects especially when lifelong therapy is required [20]. Not surprisingly, there is growing interest in developing naturally based products to attenuate insulin resistance as safer alternatives. Food derived products are valuable sources of novel therapeutic agents which are generally perceived as safer options compared to synthetic pharmacological drugs [21]. Several food proteins derived hydrolysates and peptides have undergone evaluation for therapeutic usage in metabolic disorders [22, 23].

Egg is a valuable source of dietary proteins. In addition to the nutritional value, egg proteins are also a source for peptides with myriad bioactive properties [24], including ACE inhibition [25–27]. Previous work from our research group has demonstrated the potential of egg white protein ovotransferrin derived bioactive peptides against hypertension, oxidative stress and inflammation *in vitro* and *in vivo* [25,28,29]. Moreover, we have recently reported the effects of egg white hydrolysate (EWH) on reducing blood pressure in hypertensive rats [30]. EWH significantly reduced blood pressure through modulating RAS components, reducing nitrosative stress and enhancing vascular relaxation [30]. While some features of metabolic syndrome such as inflammation and hypertension appear amenable to treatment with egg white protein derivatives [25,28–31], their actions on adipocyte functions have remained largely unknown.

Given the interplay among hypertension, inflammation, and metabolic syndrome, the objective of the study was to test the effect of EWH on differentiation, insulin signaling and inflammatory responses in 3T3-F442A pre-adipocytes. The findings of this study indicate insulin mimetic and insulin sensitizing as well as anti-inflammatory actions of EWH in adipocytes which may potentially prevent or alleviate the complications of metabolic syndrome.

## Material and methods

### Reagents

Pasteurized liquid egg white was purchased from Egg Processing Innovation Cooperative (Lethbridge, Alberta, Canada). Dulbecco's phosphate buffered saline (PBS), LipidTox dye and dithiothreitol (DTT) were all bought from Sigma Aldrich (St Louis, MO, USA). Dulbecco's modified Eagle medium (DMEM) and fetal bovine serum (FBS) were from Gibco/Invitrogen (Carlsbad, CA, USA). The murine tumor necrosis factor alpha (TNF- $\alpha$ ) was obtained from Peprotech (Rocky Hill, NJ, USA). Triton-X-100 was from VWR International (West Chester, PA, USA). Type 1 Collagenase used for cell splitting was from Worthington Biochemical Corporation (Lakewood, NJ, USA). Thermoase PC10F (from *Bacillus thermoproteolyticus* Var. Rokko) was purchased from Amano Enzyme Inc. (Nagoya, Japan). Pepsin (from porcine stomach, 10000 units / mg) was purchased from American Laboratories Inc. (Omaha, NE, USA).

### Preparation of egg white hydrolysate (EWH)

Hydrolysis of egg white was carried out according to our previous method with slight modifications [32]. Briefly, liquid egg white was diluted with water at a ratio of 1:1 (v/v) to obtain a solution with 5% protein solid. After adjusting the pH to 8.0 with 2 M NaOH solution, and the temperature to 65°C, thermoase (0.1%, w/w) was added and protein digestion was carried out for 90 min. The enzyme was then inactivated by adjusting pH to 2.5 for pepsin digestion. The mixture was further hydrolyzed at 55°C by 1% pepsin for 180 min. The reaction was terminated by heating the solution at 95°C for 15 min and the hydrolysate was centrifuged and then condensed to obtain approximately 10% solid. The hydrolysate was then spray dried and the powder was collected and stored -20°C for further experiments. EWH was desalted with 50% acetonitrile/deionized water using Sep-Pak C18 cartridges (product #: WAT043345, Waters, Ontario, Canada) to remove salts in the hydrolysate for using in cell experiments.

### Cell culture & differentiation

The murine pre-adipocyte cell line 3T3-F442A (Sigma Aldrich; Cat# 00070654) was used. Cell culture method is similar to our previous study [33]. The cells were obtained in passage 8, thawed and expanded in culture using DMEM supplemented with 10% FBS (heat-inactivated) and antibiotics. The cells were grown in T-25 flasks to confluence prior to sub-culture in gelatin-coated 48 well plates. All studies were performed using cells in passages 11–37.

To determine the ability of EWH to induce adipogenic differentiation, the cells (grown in 48 well plates) were incubated in standard culture medium (DMEM + 10% FBS + antibiotics) in the presence of EWH or insulin for 72 h without changing the medium. Adipogenic changes were determined by the appearance of intracellular lipid droplets (as shown by LipidTox staining), upregulation of PPAR $\gamma$  and CCAAT/enhancer binding protein alpha (C/EBP- $\alpha$ ) (determined by western blot) and release of adiponectin (measured by ELISA). Insulin (10  $\mu$ g/mL) was used only as a positive control for inducing differentiation.

S961/Insulin Receptor Antagonist (cat#051–86, Phoenix pharmaceuticals Inc. USA) was used at the concentration of 200 nM to investigate the involvement of insulin receptor for the potential effects of EWH on insulin signaling.

For inflammation studies, confluent monolayers of cells were treated with/ without EWH for 48 h followed by administration of murine TNF- $\alpha$  (24 hrs for COX-2, 15 minutes for cell signaling experiment).

## Intracellular lipid staining

Intracellular lipid accumulation, a marker for adipogenic differentiation, was determined by LipidTox staining as described in our previous study [33]. Briefly, the cells were treated for 72 h with EWH or insulin (positive control), fixed and stained with LipidTox (1:250 in phosphate buffered saline) and counter-stained with the nuclear dye Hoechst 33342. The cells were then visualized under an Olympus IX81 fluorescent microscope (Carson Scientific Imaging Group; Markham, Ontario, Canada). Images were obtained and analyzed using the Metamorph imaging software (Molecular Devices, Sunnyvale, CA) and presented at (200X) magnification. A control image from a group of cells without LipidTox was used to detect any nonspecific fluorescence. The images were then quantified by subtracting the background fluorescence of the control image, so only fluorescence from the lipid-specific staining was visible. The cell nuclei were stained by the DNA stain Hoechst3342. The fluorescence intensity was then measured for quantitative analysis and quantified as mean intensity per cell (MFI/cell) and expressed as % of untreated cells.

## Adiponectin measurement

The culture media from untreated and EWH (or insulin) treated cells were centrifuged (10,000 g for 10 min at 4°C) to yield cell-free supernatants which were stored at -80°C until time of the assay. These supernatants were thawed and used in the Mouse Adiponectin DuoSet ELISA kit (R&D Systems; Minneapolis, MN, USA) following the manufacturer's instructions. Data were normalized to supernatants from the untreated cells.

## Western blotting

Western blotting was done on 3T3-F442A cell lysates prepared at the end of experimental procedures as described in our previous studies [33,34]. Protein bands for C/EBP- $\alpha$  (rabbit polyclonal antibody from Cell Signaling Technology, Boston, MA, USA, cat# 2295), PPAR $\gamma$  (rabbit polyclonal antibody from Cell Signaling Technology, cat# 2430), phospho-Akt (rabbit polyclonal antibody from Cell Signaling Technology, cat #9271), Akt (mouse monoclonal from Santa Cruz, cat#sc-81434), phospho-ERK1/2 (rabbit polyclonal antibody from Cell Signaling Technology, cat#9101), ERK1/2 (mouse monoclonal antibody from Cell Signaling Technology, cat#4696), phospho-IRS-1 (rabbit polyclonal antibody from Cell Signaling Technology, cat#3070), IRS-1 (mouse monoclonal antibody from Santa Cruz Biotechnology, Santa Cruz, CA, USA, cat# sc-8038), phospho-p65 (rabbit polyclonal antibody from Santa Cruz Biotechnology, cat# sc-3033) and p65 (mouse monoclonal antibody from Santa Cruz Biotechnology, cat# sc-8008) were normalized to  $\alpha$ -tubulin (rabbit polyclonal antibody from Abcam, Cambridge, MA, cat# ab15246). Anti-tubulin was used at 0.4  $\mu$ g/ml, while all other antibodies were used at 0.5–1  $\mu$ g/ml. Goat anti-rabbit and Donkey anti-mouse conjugated secondary antibodies were purchased from Li-cor Biosciences (Lincoln, NB). The protein bands were detected by a Li-cor Odyssey BioImager and quantified by densitometry using corresponding software Odyssey v3.0 (Li-cor). Cell lysates from untreated cells were loaded on every gel and all data were expressed as % of the corresponding untreated control.

## Statistical analysis

All data are expressed as mean $\pm$ SEM (standard error of mean) of 4–8 independent experiments. Data were analyzed by one-way analysis of variance (ANOVA) with an appropriate post-hoc test (Dunnett's test for comparison to control group; Tukey's test for multiple comparisons). For studying interactions between 2 independent variables (e.g. EWH and insulin),

two-way ANOVA was used. A repeated measures test was used when applicable. The PRISM 6 statistical software (GraphPad Software, San Diego, CA) was used for the analyses. A value of  $p < 0.05$  was considered significant.

## Results

### EWH treatment induces adipogenic differentiation in 3T3-F442A cells

Adipogenic differentiation in pre-adipocytes is characterized by increases in intracellular lipid droplets and release of adiponectin [35,36]. Treating 3T3-F442A cells with EWH for 72 hrs resulted in increased intracellular lipid accumulation as determined by LipidTox staining (Fig 1A). These changes were accompanied by higher levels of adiponectin released into the culture medium, further demonstrating the pro-differentiation properties of EWH (Fig 1B). Interestingly, both effects induced by EWH were similar in magnitude to those of insulin, suggesting that the beneficial effects of EWH could be comparable to the physiological effects of insulin.

### EWH upregulates markers of adipocyte differentiation in 3T3-F442A cells

In addition to lipid accumulation and adiponectin release, adipocyte differentiation is also accompanied by increased expression of a number of proteins involved in different stages of this process. For example, PPAR $\gamma$ , an anti-inflammatory metabolic modulator is highly expressed in differentiated adipocytes and contributes to insulin sensitizing actions [37]. Similarly, C/EBP- $\alpha$ , a transcriptional regulator, is upregulated during differentiation where it coordinates the expression of downstream proteins involved in adipogenesis [35]. Indeed, 72 hrs incubation of 3T3-F442A cells with EWH upregulated both PPAR $\gamma$  (Fig 2A) and C/EBP- $\alpha$  (Fig 2B), demonstrating the successful induction of adipogenic differentiation event at the molecular level. Both effects were also comparable to those induced by insulin, the physiological agonist of adipocyte differentiation.



### EWH upregulates PPAR $\gamma$ expression dose-dependently in 3T3-F442A cells

The effects of different concentrations of EWH on PPAR $\gamma$  expression was also investigated in 3T3-F442A cells. Fig 3 illustrates that EWH enhanced PPAR $\gamma$  expression in a dose-dependent manner. EWH at concentrations of 2.5, 5 and 10 mg/mL enhanced PPAR $\gamma$  expression significantly as compared to untreated cells at  $P < 0.01$ ,  $P < 0.001$  and  $P < 0.0001$  respectively.

### EWH exerts both insulin mimetic and insulin sensitizing effects

Given the similarity in EWH responses to insulin effects, we then investigated the effect of EWH on key insulin signaling pathways in these cells. The mitogen activated protein kinase ERK1/2 is an important downstream signaling target of insulin, which is phosphorylated (and hence, activated) by insulin treatment [38]. Treatment with EWH alone showed increased ERK1/2 phosphorylation in pre-adipocytes, while insulin-induced ERK1/2 activation in EWH-treated cells was comparable to that observed in EWH-free cells (Fig 4A), suggesting a potential insulin mimetic action of EWH.

Another major signaling target of insulin is protein kinase B (PKB)/Akt. Akt regulates many cellular processes including metabolism, proliferation, cell survival, growth and angiogenesis [39]. Akt phosphorylation is a key event involved in mediating the beneficial actions of insulin in glucose transport in adipose tissues [40]. Interestingly, EWH alone had no effects on Akt phosphorylation; while insulin actions on Akt phosphorylation were enhanced in EWH-



**Fig 1. EWH treatment induces adipogenic differentiation in 3T3-F442A cells.** 3T3-F442A cells were incubated with EWH (5 mg/mL) or insulin (positive control; 10  $\mu$ g/mL) for 72 hrs. (a) Following incubation, the cells were fixed and stained with the neutral lipid-specific dye LipidTox (green), the nuclear stain Hoechst33342

(blue) and visualized under fluorescence microscopy. A set of representative images are shown. (b) The cell-free culture supernatants were collected and analyzed by ELISA to determine adiponectin levels. Data are presented as mean $\pm$ SEM of 4–5 independent experiments. \* and \*\* indicate  $p < 0.05$  and  $p < 0.01$  respectively compared to the untreated control (Untr).

<https://doi.org/10.1371/journal.pone.0185653.g001>

treated cells over and above the response seen in control cells (Fig 4B). Indeed, a 2-way ANOVA showed a significant interaction between EWH and insulin effects on Akt phosphorylation indicating a novel insulin sensitizing action for EWH in these cells.

## EWH appears to involve insulin receptor signaling in adipocytes

Next, we examined if the observed EWH actions were dependent on signaling through the insulin receptor. Insulin binding to its receptor leads to phosphorylation of an associated protein, insulin receptor substrate 1 (IRS-1), which is widely used as a marker for insulin receptor mediated functions [41–43]. Interestingly, EWH treatment alone enhanced phosphorylation of insulin receptor  $\beta$  (IR $\beta$ ) (Fig 5A,  $P < 0.05$ ), while the protein expression level of IR $\beta$  was not affected by the treatment (Fig 5B,  $P > 0.05$ ). Furthermore, EWH treatment alone also enhanced IRS-1 phosphorylation (Fig 5C) significantly ( $P < 0.01$ ) compared to untreated cells indicating potential insulin mimetic actions of EWH in these cells. However, EWH did not further enhance the phosphorylation of IR $\beta$  and IRS-1 in the presence of exogenous insulin. Only insulin was able to induce significant phosphorylation of IR $\beta$  and IRS-1, which remained unaffected by concomitant presence of EWH (Fig 5A and 5C).


## Insulin mimetic effect of EWH on ERK phosphorylation is mediated through insulin receptor in adipocytes

Since EWH exerted insulin mimetic effects in 3T3-F442A cells (Fig 4A) and was mediated through insulin receptor (Fig 5A), we aimed to further explore this possibility by using S961, an insulin receptor antagonist. As indicated in Fig 6, EWH enhanced ERK phosphorylation significantly compared to untreated cells ( $P < 0.001$ ) (similar to insulin,  $P < 0.05$ ), whereas, incubating in the presence of the insulin receptor antagonist (S961) blocked the observed effects of both insulin and EWH on the levels comparable to the untreated control. This data suggests that, the insulin mimetic effects of EWH on ERK phosphorylation is potentially mediated through insulin receptor.


## EWH modulates inflammatory response in 3T3-F442A cells

Finally, we investigated the effects of EWH on inflammatory changes in these cells. Adipocyte inflammation releases harmful cytokines which leads to the loss of protective adipokines, increases insulin resistance and contributes to the pathogenesis of metabolic syndrome [44,45]. We used TNF- $\alpha$ , a pro-inflammatory cytokine involved in various inflammatory, atherosclerotic and metabolic disorders, to induce inflammation in these cells. Treatment with TNF- $\alpha$  for 24 hrs upregulated cyclooxygenase -2 (COX-2) levels in 3T3-F442A cells ( $1.64 \pm 0.21$ ), while a 48 hrs pre-treatment with EWH abolished this response ( $0.84 \pm 0.19$ ) (Fig 7A), indicating potentially beneficial anti-inflammatory capabilities of EWH. Further examination of underlying pro-inflammatory signaling/transcriptional pathways revealed a reduction of TNF- $\alpha$ -mediated c-Jun phosphorylation in EWH-treated cells. TNF- $\alpha$  increased c-Jun phosphorylation to  $159.20 \pm 12.17\%$  in adipocytes, while EWH treatment restored it to the basal level ( $98.15 \pm 15.08\%$ ) (Fig 7B). This may account for the mechanisms underlying the anti-inflammatory actions of EWH.

**a**



**b**




**Fig 2. EWH upregulates markers of adipocyte differentiation.** 3T3-F442A cells were incubated with EWH (5 mg/mL) or insulin (positive control; 10 µg/mL) for 72 hrs. The cells were then lysed and western blotting of the lysates was performed with antibodies against PPAR $\gamma$  (a), C/EBP- $\alpha$  (b) and  $\alpha$ -tubulin (loading control; both a and b). A representative set of images are shown. Bands were quantified by densitometric analysis. Data are presented as mean $\pm$ SEM of 4–5 independent experiments. \*, \*\* and \*\*\* indicate  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$  respectively, compared to the untreated control (Untr).

<https://doi.org/10.1371/journal.pone.0185653.g002>



## Discussion

Adipose tissue playing a vital role in energy homeostasis, is one of the critical target organs for insulin actions [11]. Adipose tissue secretes a number of adipokines interacting with central and peripheral organs in the body [10]. This insulin sensitive tissue influences diverse metabolic processes including carbohydrate metabolism, lipid metabolism, inflammation, blood pressure, energy expenditure, and feeding behavior [46,47]. In metabolically normal conditions adipocytes are small in size, sensitive to insulin and secrete insulin sensitizing hormones



**Fig 3. EWH upregulates PPAR $\gamma$  expression in a dose-dependent manner.** 3T3-F442A cells were incubated with different dosages of EWH (0.63–10 mg/mL) or insulin (10 µg/mL) for 72 hrs. The cells were then lysed and western blotting of the lysates was performed with antibodies against PPAR $\gamma$  and  $\alpha$ -tubulin (loading control). A representative set of images is shown. Bands were quantified by densitometric analysis. Data are presented as mean $\pm$ SEM of 3–4 independent experiments. \*\*, \*\*\* and \*\*\*\* indicate  $p < 0.01$ ,  $p < 0.001$  and  $p < 0.0001$  respectively, compared to the untreated control (Untr).

<https://doi.org/10.1371/journal.pone.0185653.g003>



**Fig 4. EWH differentially modulates insulin-mediated phosphorylation of ERK and Akt.** 3T3-F442A cells were incubated with EWH (5 mg/mL) for 48 hrs prior to stimulation with insulin (10  $\mu$ g/mL) for 20 or 40 min. The cells were then lysed and western blotting of the lysates was performed with antibodies against the total and phosphorylated forms of ERK (a) and Akt (b). Representative sets of images are shown. Bands were quantified by densitometric analysis. Data are presented as mean $\pm$ SEM of 5 independent experiments. \*, \*\* and \*\*\* indicate  $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$  respectively compared to untreated control by a two-way ANOVA. 'ns' indicates: not significant.


<https://doi.org/10.1371/journal.pone.0185653.g004>

such as adiponectin [35]. In metabolic disorders on the contrary, adipocytes become larger in size, inflamed, insulin resistant, and increasingly express harmful adipokines leading to adipose tissue dysfunction, insulin resistance and associated diseases [14]. In fact loss of insulin sensitivity in adipose tissue adversely affects glucose utilization and lipid storage leading to ectopic deposition of fat in insulin sensitive tissues which contributes towards the development of insulin resistance and pathogenesis of both type II diabetes and metabolic syndrome [9,48]. Enhanced differentiation of fibroblast-like pre-adipocytes into mature adipocytes by the use of compounds mimicking insulin functions or enhancing insulin sensitivity provides a novel strategy for controlling the complications of metabolic syndrome [49–51].

PPAR $\gamma$  and C/EBP- $\alpha$  are the two key molecules involved in adipocyte differentiation and regulation of the adipogenic network [52]. Over expression of PPAR $\gamma$  can induce adipogenesis in mouse embryonic fibroblasts lacking C/EBP $\alpha$ , but C/EBP $\alpha$  cannot rescue adipogenesis when PPAR $\gamma$  is not expressed, showing that PPAR $\gamma$  is the master regulator of adipogenesis [37,53,54]. Thiazolidinediones with insulin sensitizing effects promote pre-adipocytes differentiation by PPAR $\gamma$  activation [55,56]. Insulin also promotes adipogenic effects by upregulating both PPAR $\gamma$  and C/EBP- $\alpha$  in adipocytes [37,57]. Our study revealed that EWH treatment also exerted insulin-like differentiating effects on pre-adipocytes. In accordance with observed insulin-like effects of EWH on upregulation of pre-adipocyte differentiation markers, this treatment also enhanced PPAR $\gamma$  and C/EBP- $\alpha$  to the same extent as insulin in adipocytes. So, it is plausible to propose that the observed insulin-like properties of EWH on adipogenic response may at least be in part due to involvement of these 2 regulators in 3T3-F442A cells. Indeed, several groups have identified plant derived novel compounds that promote adipogenic effects at least partially through upregulation of PPAR $\gamma$  [58,59].


Binding of insulin to insulin receptor triggers the phosphorylation of IR $\beta$  and consequently, insulin receptor substrate (IRS) proteins providing the basis for the subsequent association with downstream signaling through different pathways mediating metabolic and mitogenic responses of insulin [42,60]. While phosphorylation and activation of Akt is responsible for most of the known metabolic effects of insulin, ERK phosphorylation mediates mitogenic and transcriptional effects of insulin in adipocytes [61]. When investigating the effects of EWH on insulin signaling, we also observed an insulin mimetic effect of EWH on ERK1/2 phosphorylation in these cells. In addition to the observed insulin mimetic effects on 3T3-F442A cells, EWH also exhibited insulin sensitizing effects by enhancing insulin-mediated Akt phosphorylation. Akt acts not only as a regulator of glucose transport but also involves in several other metabolic actions including glycolysis, protein synthesis, lipogenesis, glycogen synthesis, suppression of gluconeogenesis, cell survival, determination of cell size and cell-cycle progression [62]. The fact that EWH affected both ERK and Akt phosphorylation in adipocytes indicates the potential effects of this treatment on both pathways of insulin signaling.

Moreover, since phosphorylation of IR $\beta$  and IRS-1 was significantly enhanced in EWH treated cells while no further increase was observed in the presence of exogenous insulin suggesting that EWH exerts its insulin mimetic effects through insulin receptor which was further supported by the study of an insulin receptor antagonist (Fig 6). Therefore, the insulin




**Fig 5. EWH actions are partly mediated through insulin receptor.** 3T3-F442A cells were incubated with EWH (5 mg/mL) for 48 hrs prior to stimulation with insulin (10 µg/mL) for 20 or 40 min. The cells were then lysed and western blotting of the lysates was performed with antibodies against the: total and phosphorylated forms of IRβ (a), total IRβ and α-tubulin (loading control) (b), and total and phosphorylated forms of IRS-1 (c). Representative sets of images are shown. Bands were quantified by densitometric analysis. Data are presented as mean±SEM of 4–7 independent experiments. \*, \*\* and \*\*\*\* indicate p<0.05, p<0.01, and p<0.0001 respectively.

<https://doi.org/10.1371/journal.pone.0185653.g005>



**Fig 6. EWH mediates its insulin mimetic effects through insulin receptor.** 3T3-F442A cells were incubated with EWH (5 mg/mL) or insulin (positive control; 10 µg/mL) in the presence/absence of S961 (insulin receptor antagonist; 200 nM) for 72 hrs. The cells were then lysed and western blotting of the lysates was performed with antibodies against the total and phosphorylated forms of ERK. A representative image is shown. Bands were quantified by densitometric analysis. Data are presented as mean±SEM of 3 independent experiments. \*, and \*\*\* indicate p<0.05 and p<0.001 compared to untreated cells respectively.

<https://doi.org/10.1371/journal.pone.0185653.g006>



**Fig 7. EWH exerts inhibitory effects on adipocyte inflammation.** The 3T3-F442A cells were incubated with EWH (5 mg/mL) for 48 hrs prior to stimulation with TNF- $\alpha$  for either (a) 24 hrs (for inflammatory marker expression) or (b) 15 min (for transcription pathway experiment). The cells were then lysed and western blotting of the lysates was performed with antibodies against COX-2 and the loading control  $\alpha$ -tubulin (a) or the total and phosphorylated forms of c-Jun (b). A representative set of images are shown. Bands from the COX-2 study were quantified by densitometric analysis. Data is mean $\pm$ SEM of 4–6 independent experiments. \* indicates  $p < 0.05$  compared to the untreated (Untr) group, while ## indicates  $p < 0.01$  compared to TNF- $\alpha$  alone.

<https://doi.org/10.1371/journal.pone.0185653.g007>

sensitizing effect of EWH might be via targets downstream to IRS-1 such as phosphoinositide-3 kinase (PI3-kinase) and phosphoinositide-dependent kinase 1 (PDK1).

There is an interest to assess the potential of established antihypertensive compounds for protection against insulin resistance and other complications of metabolic syndrome due to the role of RAS impairment in the pathogenesis of such diseases [63]. RAS blockade has been reported to inhibit the body fat mass increase [64,65], and improve insulin resistance and glucose tolerance in type-2 diabetic rodents [66,67]. Captopril, the pharmacological ACE inhibitor, has been reported to enhance adipocyte differentiation and reduce inflammation in various tissues [68]. Insulin sensitizing effects of RAS blockade have also been reported in clinical studies in patients with risk factors [69–71] suggesting additional benefits of these drugs in a complex condition like metabolic syndrome. Interestingly, the anti-hypertensive EWH with RAS modulating properties (reducing vascular ACE and angiotensin II type 1 receptor expression) also enhanced pre-adipocyte differentiation, and induced insulin mimetic and sensitizing effects in 3T3-F442A cells. Similarly, milk derived peptides IPP and VPP, with ACE-inhibitory and anti-inflammatory properties exerted insulin mimetic adipogenic effects by promoting the differentiating of pre-adipocytes in 3T3-F442A cells [33]. The flaxseed protein hydrolysate contains peptide fractions with anti-hypertensive [72] as well as anti-diabetic properties [73]. Since hypertension, inflammation and insulin resistance present concomitantly in many cases of metabolic syndrome, EWH as a novel naturally based compound with multiple benefits against hypertension, inflammation and insulin functions may serve as an effective option for the management of complications of this disease.

Adipose tissue inflammation with dysregulated adipokine secretion plays a critical role in the development of a variety of cardiometabolic disorders including metabolic syndrome, type 2 diabetes, inflammatory and vascular disorders and eventually development of coronary heart disease [9]. EWH upregulated the expression of anti-inflammatory molecules such as PPAR $\gamma$  and adiponectin, demonstrating the potential benefits of EWH on adipocyte function and metabolic syndrome. EWH also prevented the TNF- $\alpha$ -mediated induction of the pro-inflammatory enzyme COX-2, a molecule that contributes to the pathologic complications of metabolic syndrome [74,75]. This anti-inflammatory effect is likely due to its interference with the AP-1 transcription factor pathway which is involved in COX-2 expression and can be modulated by the inhibition of c-Jun phosphorylation [76–78]. TNF- $\alpha$  stimulates the pro-inflammatory phenotype in adipose tissues leading to the development of insulin resistance and metabolic syndrome [11]. Indeed, suppression of TNF- $\alpha$  has been suggested as a potential therapy against metabolic syndrome [79]. VPP and IPP prevented inflammatory changes in 3T3-F442A cells [33]. In another study, Sawada et al. have reported that VPP inhibited adipose inflammation *in vitro* and *in vivo* [80]. VPP also enhanced insulin sensitivity in obese mice and inhibited macrophage accumulation and activation in fat tissues [80]. Moreover, beta-mercaptoethanol (BME), the pharmacological redox regulator and radical scavenger, has also been reported to down-regulate the expression of inflammatory cytokines and promote adipocyte differentiation [81]. Our data with EWH further supports its role as a novel regulator of adipose functions with additional anti-inflammatory benefits.

Bioactive peptides in the EWH are potentially responsible for the observed effects of EWH on adipogenic differentiation, insulin signaling and anti-inflammatory responses in adipocytes. Purification and fractionation of EWH to identify its responsible peptides with beneficial effects on adipocyte differentiation is essential in understanding the structure requirements of food-derived bioactive peptides with beneficial effects on adipose tissue function. We have fractionated EWH using stepwise chromatographic methods and have characterized the peptides responsible for adipogenic responses in adipocytes (PPAR $\gamma$  expression). Among total 42 peptides identified from EWH, of four peptides (ERYPIL, VFKGL, WEKAFKDED, and QAMPFRVTEQE) significantly enhanced PPAR $\gamma$  expression, compared to untreated cells (*unpublished data*).

In conclusion, our study demonstrated that EWH, with RAS modulating properties promoted adipocyte differentiation through a combination of insulin mimetic and insulin sensitizing actions on 3T3-F442A cells. In addition, EWH also increased expression of the anti-inflammatory hormone adiponectin and suppressed cytokine mediated inflammatory response in these cells. Considering the fundamental role of adipose tissue dysfunction in the pathogenesis of hypertension, inflammation, insulin resistance, and metabolic syndrome, EWH may have potential benefits in the prevention and management of metabolic syndrome.

## Supporting information

**S1 File.** The relevant data used for generating figures and statistical analysis in this study. (XLSX)

## Acknowledgments

The authors would like to thank Qi Yi Li for the preparation of EWH for this study.

## Author Contributions

**Conceptualization:** Forough Jahandideh, Subhadeep Chakrabarti, Sandra T. Davidge, Jianping Wu.

**Data curation:** Forough Jahandideh, Subhadeep Chakrabarti.

**Funding acquisition:** Jianping Wu.

**Investigation:** Forough Jahandideh, Jianping Wu.

**Methodology:** Forough Jahandideh, Subhadeep Chakrabarti, Jianping Wu.

**Project administration:** Jianping Wu.

**Resources:** Jianping Wu.

**Supervision:** Sandra T. Davidge, Jianping Wu.

**Writing – original draft:** Forough Jahandideh.

**Writing – review & editing:** Subhadeep Chakrabarti, Sandra T. Davidge, Jianping Wu.

## References

1. Salazar MR, Carbajal HA, Espeche WG, Aizpurua M, Leiva Sisniegues CE, March CE, et al. (2013) Identifying cardiovascular disease risk and outcome: use of the plasma triglyceride/high-density lipoprotein cholesterol concentration ratio versus metabolic syndrome criteria. *J Intern Med* 273: 595–601. <https://doi.org/10.1111/joim.12036> PMID: 23331522

2. Potenza MV, Mechanick JI (2009) The Metabolic Syndrome: Definition, Global Impact, and Pathophysiology. *Nutr Clin Pract* 24: 560–577. <https://doi.org/10.1177/0884533609342436> PMID: 19841245
3. Wong ND (2007) Metabolic syndrome: cardiovascular risk assessment and management. *American journal of cardiovascular drugs: drugs, devices, and other interventions* 7: 259–272. PMID: 17696567
4. Grundy SM (2008) Metabolic syndrome pandemic. *Arterioscl Throm Vas* 28: 629–636.
5. Yvan-Charvet L, Quignard-Boulangé A (2011) Role of adipose tissue renin-angiotensin system in metabolic and inflammatory diseases associated with obesity. *Kidney international* 79: 162–168. <https://doi.org/10.1038/ki.2010.391> PMID: 20944545
6. Yusuf S, Gerstein H, Hoogwerf B, Pogue J, Bosch J, Wolfenbutter BHR, et al. (2001) Ramipril and the development of diabetes. *Jama-J Am Med Assoc* 286: 1882–1885.
7. Braga MFB, Leiter LA (2009) Role of Renin-Angiotensin System Blockade in Patients With Diabetes Mellitus. *Am J Cardiol* 104: 835–839. <https://doi.org/10.1016/j.amjcard.2009.05.015> PMID: 19733720
8. Zarich SW (2005) Cardiovascular risk factors in the metabolic syndrome: impact of insulin resistance on lipids, hypertension, and the development of diabetes and cardiac events. *Reviews in cardiovascular medicine* 6: 194–205. PMID: 16379015
9. Coppack S, Mohamed-Ali V, Karpe F (2005) Metabolic Syndrome: Insulin Resistance, Obesity, Diabetes Mellitus, Hypertension, Physical Activity and Genetic Factors. *Cardiovascular Disease: Diet, Nutrition and Emerging Risk Factors*: 22–49.
10. Havel PJ (2002) Control of energy homeostasis and insulin action by adipocyte hormones: leptin, acylation stimulating protein, and adiponectin. *Curr Opin Lipidol* 13: 51–59. PMID: 11790963
11. Ruan H, Lodish HF (2003) Insulin resistance in adipose tissue: direct and indirect effects of tumor necrosis factor- $\alpha$ . *Cytokine Growth F R* 14: 447–455.
12. Lu HL, Wang HW, Wen Y, Zhang MX, Lin HH (2006) Roles of adipocyte derived hormone adiponectin and resistin in insulin resistance of type 2 diabetes. *World J Gastroenterol* 12: 1747–1751.
13. Harwood HJ Jr. (2012) The adipocyte as an endocrine organ in the regulation of metabolic homeostasis. *Neuropharmacology* 63: 57–75. <https://doi.org/10.1016/j.neuropharm.2011.12.010> PMID: 22200617
14. Kadowaki T, Hara K, Yamauchi T, Terauchi Y, Tobe K, Nagai R (2003) Molecular mechanism of insulin resistance and obesity. *Exp Biol Med (Maywood)* 228: 1111–1117.
15. Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM (2001) C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *Jama* 286: 327–334. PMID: 11466099
16. Winkler G, Kiss S, Keszthelyi L, Sapi Z, Ory I, Salamon F, et al. (2003) Expression of tumor necrosis factor (TNF)- $\alpha$  protein in the subcutaneous and visceral adipose tissue in correlation with adipocyte cell volume, serum TNF- $\alpha$ , soluble serum TNF-receptor-2 concentrations and C-peptide level. *European journal of endocrinology / European Federation of Endocrine Societies* 149: 129–135.
17. Majumder K, Wu J (2009) Angiotensin I converting enzyme inhibitory peptides from simulated in vitro gastrointestinal digestion of cooked eggs. *Journal of agricultural and food chemistry* 57: 471–477. <https://doi.org/10.1021/jf8028557> PMID: 19154160
18. Majumder K, Panahi S, Kaufman S, Wu JP (2013) Fried egg digest decreases blood pressure in spontaneous hypertensive rats. *J Funct Foods* 5: 187–194.
19. de Souza CJ, Eckhardt M, Gagen K, Dong M, Chen W, Laurent D, et al. (2001) Effects of pioglitazone on adipose tissue remodeling within the setting of obesity and insulin resistance. *Diabetes* 50: 1863–1871. PMID: 11473050
20. Khan HMS, Murtaza G, Usman M, Rasool F, Akhtar M, Qureshi MIM, et al. (2012) Evidence based study of side effects of drugs used in the treatment of diabetes mellitus. *Afr J Pharm Pharmacol* 6: 1805–1808.
21. Chakrabarti S, Jahandideh F, Wu J (2014) Food-derived bioactive peptides on inflammation and oxidative stress. *BioMed research international* 2014: 608979. <https://doi.org/10.1155/2014/608979> PMID: 24527452
22. Shahidi F, Zhong Y (2008) Bioactive peptides. *J AOAC Int* 91: 914–931. PMID: 18727554
23. Erdmann K, Cheung BW, Schroder H (2008) The possible roles of food-derived bioactive peptides in reducing the risk of cardiovascular disease. *The Journal of nutritional biochemistry* 19: 643–654. <https://doi.org/10.1016/j.jnutbio.2007.11.010> PMID: 18495464
24. Sumner DA, Gow H, Hayes D, Matthews W, Norwood B, Rosen-Molina JT, et al. (2011) Economic and market issues on the sustainability of egg production in the United States: analysis of alternative production systems. *Poult Sci* 90: 241–250. <https://doi.org/10.3382/ps.2010-00822> PMID: 21177466
25. Majumder K, Chakrabarti S, Morton JS, Panahi S, Kaufman S, Davidge ST, et al. (2013) Egg-derived tri-peptide IRW exerts antihypertensive effects in spontaneously hypertensive rats. *PloS one* 8: e82829. <https://doi.org/10.1371/journal.pone.0082829> PMID: 24312436

26. Jahandideh F, Majumder K, Chakrabarti S, Morton JS, Panahi S, Kaufman S, et al. (2014) Beneficial effects of simulated gastro-intestinal digests of fried egg and its fractions on blood pressure, plasma lipids and oxidative stress in spontaneously hypertensive rats. *PloS one* 9: e115006. <https://doi.org/10.1371/journal.pone.0115006> PMID: 25502445
27. Majumder K, Chakrabarti S, Morton JS, Panahi S, Kaufman S, Davidge ST, et al. (2015) Egg-derived ACE-inhibitory peptides IQW and LKP reduce blood pressure in spontaneously hypertensive rats. *J Funct Foods* 13: 50–60.
28. Huang W, Shen S, Nimalaratne C, Li S, Majumder K, Wu J (2012) Effects of addition of egg ovotransferrin-derived peptides on the oxygen radical absorbance capacity of different teas. *Food Chem* 135: 1600–1607. <https://doi.org/10.1016/j.foodchem.2012.05.093> PMID: 22953899
29. Huang W, Chakrabarti S, Majumder K, Jiang Y, Davidge ST, Wu J (2010) Egg-derived peptide IRW inhibits TNF-alpha-induced inflammatory response and oxidative stress in endothelial cells. *Journal of agricultural and food chemistry* 58: 10840–10846. <https://doi.org/10.1021/jf102120c> PMID: 20886881
30. Jahandideh F, Chakrabarti S, Majumder K, Li Q, Panahi S, Morton JS, Davidge ST, Wu J (2016) Egg white protein hydrolysate reduces blood pressure, improves vascular relaxation and modifies aortic angiotensin II receptors expression in spontaneously hypertensive rats. *J Funct Foods*.
31. Moon SH, Lee JH, Lee YJ, Chang KH, Paik JY, Ahn DU, et al. (2013) Screening for cytotoxic activity of ovotransferrin and its enzyme hydrolysates. *Poult Sci* 92: 424–434. <https://doi.org/10.3382/ps.2012-02680> PMID: 23300310
32. Jahandideh F, Chakrabarti S, Majumder K, Li QY, Panahi S, Morton JS, et al. (2016) Egg white protein hydrolysate reduces blood pressure, improves vascular relaxation and modifies aortic angiotensin II receptors expression in spontaneously hypertensive rats. *J Funct Foods* 27: 667–673.
33. Chakrabarti S, Wu J (2015) Milk-derived tripeptides IPP (Ile-Pro-Pro) and VPP (Val-Pro-Pro) promote adipocyte differentiation and inhibit inflammation in 3T3-F442A cells. *PloS one* 10: e0117492. <https://doi.org/10.1371/journal.pone.0117492> PMID: 25714093
34. Chakrabarti S, Davidge ST (2009) High glucose-induced oxidative stress alters estrogen effects on ERalpha and ERbeta in human endothelial cells: Reversal by AMPK activator. *J Steroid Biochem Mol Biol*.
35. Fu YC, Luo NL, Klein RL, Garvey WT (2005) Adiponectin promotes adipocyte differentiation, insulin sensitivity, and lipid accumulation. *J Lipid Res* 46: 1369–1379. <https://doi.org/10.1194/jlr.M400373-JLR200> PMID: 15834118
36. Christensen KB, Minet A, Svenstrup H, Grevsen K, Zhang HB, Schrader E, et al. (2009) Identification of Plant Extracts with Potential Antidiabetic Properties: Effect on Human Peroxisome Proliferator-activated Receptor (PPAR), Adipocyte Differentiation and Insulin-stimulated Glucose Uptake. *Phytotherapy Research* 23: 1316–1325. <https://doi.org/10.1002/ptr.2782> PMID: 19172665
37. Hamm JK, El Jack AK, Pilch PF, Farmer SR (1999) Role of PPAR gamma in regulating adipocyte differentiation and insulin-responsive glucose uptake. *The Metabolic Syndrome X* 892: 134–145.
38. Eduardsen K, Larsen SL, Novak I, Lambert IH, Hoffmann EK, Pedersen SF (2011) Cell volume regulation and signaling in 3T3-L1 pre-adipocytes and adipocytes: on the possible roles of caveolae, insulin receptors, FAK and ERK1/2. *Cell Physiol Biochem* 28: 1231–1246. <https://doi.org/10.1159/000335855> PMID: 22179011
39. Hers I, Vincent EE, Tavaré JM (2011) Akt signalling in health and disease. *Cell Signal* 23: 1515–1527. <https://doi.org/10.1016/j.cellsig.2011.05.004> PMID: 21620960
40. Kleiman E, Carter G, Ghansah T, Patel NA, Cooper DR (2009) Developmentally spliced PKCbeta11 provides a possible link between mTORC2 and Akt kinase to regulate 3T3-L1 adipocyte insulin-stimulated glucose transport. *Biochem Biophys Res Commun* 388: 554–559. <https://doi.org/10.1016/j.bbrc.2009.08.063> PMID: 19686698
41. Velloso LA, Carneiro EM, Crepaldi SC, Boschero AC, Saad MJ (1995) Glucose- and insulin-induced phosphorylation of the insulin receptor and its primary substrates IRS-1 and IRS-2 in rat pancreatic islets. *FEBS Lett* 377: 353–357. [https://doi.org/10.1016/0014-5793\(95\)01370-9](https://doi.org/10.1016/0014-5793(95)01370-9) PMID: 8549754
42. Zhang J, Nakatsu Y, Shinjo T, Guo Y, Sakoda H, Yamamotoya T, et al. (2013) Par14 protein associates with insulin receptor substrate 1 (IRS-1), thereby enhancing insulin-induced IRS-1 phosphorylation and metabolic actions. *J Biol Chem* 288: 20692–20701. <https://doi.org/10.1074/jbc.M113.485730> PMID: 23720771
43. Zolotnik IA, Figueroa TY, Yaspelkis BB 3rd (2012) Insulin receptor and IRS-1 co-immunoprecipitation with SOCS-3, and IKKalpha/beta phosphorylation are increased in obese Zucker rat skeletal muscle. *Life Sci* 91: 816–822. <https://doi.org/10.1016/j.lfs.2012.08.038> PMID: 22982470
44. Rodrigues CE, Vendramini MB, Bueno C, Bonfa E, de Carvalho JF (2012) Adipocytokines in primary antiphospholipid syndrome: potential markers of low-grade inflammation, insulin resistance and metabolic syndrome. *Clin Exp Rheumatol* 30: 871–878. PMID: 22935544

45. Bell DS (2000) Inflammation, insulin resistance, infection, diabetes, and atherosclerosis. *Endocr Pract* 6: 272–276. <https://doi.org/10.4158/EP.6.3.272> PMID: 11421545
46. Smorlesi A, Frontini A, Giordano A, Cinti S (2012) The adipose organ: white-brown adipocyte plasticity and metabolic inflammation. *Obes Rev* 13: 83–96. <https://doi.org/10.1111/j.1467-789X.2012.01039.x> PMID: 23107262
47. Bou M, Todorovic M, Rodriguez J, Capilla E, Gutierrez J, Navarro I (2014) Interplay of adiponectin, TNFalpha and insulin on gene expression, glucose uptake and PPARgamma, AKT and TOR pathways in rainbow trout cultured adipocytes. *Gen Comp Endocrinol* 205: 218–225. <https://doi.org/10.1016/j.ygcen.2014.05.005> PMID: 24846393
48. Muller-Wieland D, Kotzka J, Knebel B, Krone W (1998) Metabolic syndrome and hypertension: pathophysiology and molecular basis of insulin resistance. *Basic Res Cardiol* 93: 131–134. PMID: 9833175
49. Rekovets O, Sirenko YU, Savitskiy S, Pavlyuk E, Vaschylko A, Primak G (2010) Influence Different Type of Antihypertensive Therapy in Patients with Arterial Hypertension (Ah) and Metabolic Syndrome (Ms) on Insulin Resistance Depends on Sex. *J Hypertens* 28: E569–E569.
50. Salazar B, Bellon L, Jean S, Maranchi M, El-Yazidi C, Orsiere T, et al. (2004) Low-level arsenite activates the transcription of genes involved in adipose differentiation. *Cell biology and toxicology* 20: 375–385. <https://doi.org/10.1007/s10565-004-1471-1> PMID: 15868481
51. Strowski MZ, Li ZH, Szalkowski D, Shen XL, Guan XM, Juttner S, et al. (2004) Small-molecule insulin mimetic reduces hyperglycemia and obesity in a nongenetic mouse model of type 2 diabetes. *Endocrinology* 145: 5259–5268. <https://doi.org/10.1210/en.2004-0610> PMID: 15297448
52. Kadowaki T, Hara K, Kubota N, Tobe K, Terauchi Y, Yamauchi T, et al. (2002) The role of PPARgamma in high-fat diet-induced obesity and insulin resistance. *J Diabetes Complications* 16: 41–45. PMID: 11872365
53. Chinetti G, Fruchart JC, Staels B (2000) Peroxisome proliferator-activated receptors (PPARs): Nuclear receptors at the crossroads between lipid metabolism and inflammation. *Inflammation Research* 49: 497–505. <https://doi.org/10.1007/s000110050622> PMID: 11089900
54. Tsuchida A, Yamauchi T, Takekawa S, Hada Y, Ito Y, Maki T, et al. (2005) Peroxisome proliferator-activated receptor (PPAR)alpha activation increases adiponectin receptors and reduces obesity-related inflammation in adipose tissue: comparison of activation of PPARalpha, PPARgamma, and their combination. *Diabetes* 54: 3358–3370. PMID: 16306350
55. Rikimaru K, Wakabayashi T, Abe H, Imoto H, Maekawa T, Ujikawa O, et al. (2012) A new class of non-thiazolidinedione, non-carboxylic-acid-based highly selective peroxisome proliferator-activated receptor (PPAR) gamma agonists: design and synthesis of benzylpyrazole acylsulfonamides. *Bioorg Med Chem* 20: 714–733. <https://doi.org/10.1016/j.bmc.2011.12.008> PMID: 22209730
56. Spiegelman BM (1998) PPAR-gamma: adipogenic regulator and thiazolidinedione receptor. *Diabetes* 47: 507–514. PMID: 9568680
57. MacDougald OA, Cornelius P, Liu R, Lane MD (1995) Insulin regulates transcription of the CCAAT/enhancer binding protein (C/EBP) alpha, beta, and delta genes in fully-differentiated 3T3-L1 adipocytes. *J Biol Chem* 270: 647–654. PMID: 7822291
58. Beh JE, Khoo LT, Latip J, Abdullah MP, Alitheen NBM, Adam Z, et al. (2013) SDF7, a group of Scoparia dulcis Linn. derived flavonoid compounds, stimulates glucose uptake and regulates adipocytokines in 3T3-F442a adipocytes. *J Ethnopharmacol* 150: 339–352. <https://doi.org/10.1016/j.jep.2013.09.001> PMID: 24029250
59. Christensen KB, Minet A, Svenstrup H, Grevsen K, Zhang H, Schrader E, et al. (2009) Identification of plant extracts with potential antidiabetic properties: effect on human peroxisome proliferator-activated receptor (PPAR), adipocyte differentiation and insulin-stimulated glucose uptake. *Phytother Res* 23: 1316–1325. <https://doi.org/10.1002/ptr.2782> PMID: 19172665
60. Valverde AM, Kahn CR, Benito M (1999) Insulin signaling in insulin receptor substrate (IRS)-1-deficient brown adipocytes: requirement of IRS-1 for lipid synthesis. *Diabetes* 48: 2122–2131. PMID: 10535444
61. Laviola L, Perrini S, Cignarelli A, Giorgino F (2006) Insulin signalling in human adipose tissue. *Arch Physiol Biochem* 112: 82–88. <https://doi.org/10.1080/13813450600736174> PMID: 16931450
62. Whiteman EL, Cho H, Birnbaum MJ (2002) Role of Akt/protein kinase B in metabolism. *Trends Endocrinol Metab* 13: 444–451. PMID: 12431841
63. Medvedev IN, Kumova TA (2007) Angiotensin II receptor inhibitors: role and place in arterial hypertension and metabolic syndrome treatment. *Russ J Cardiol*: 97–99.
64. Mathai ML, Naik S, Sinclair AJ, Weisinger HS, Weisinger RS (2008) Selective reduction in body fat mass and plasma leptin induced by angiotensin-converting enzyme inhibition in rats. *Int J Obes (Lond)* 32: 1576–1584.

65. de Kloet AD, Krause EG, Kim DH, Sakai RR, Seeley RJ, Woods SC (2009) The effect of angiotensin-converting enzyme inhibition using captopril on energy balance and glucose homeostasis. *Endocrinology* 150: 4114–4123. <https://doi.org/10.1210/en.2009-0065> PMID: 19497971
66. Lee MH, Song HK, Ko GJ, Kang YS, Han SY, Han KH, et al. (2008) Angiotensin receptor blockers improve insulin resistance in type 2 diabetic rats by modulating adipose tissue. *Kidney international* 74: 890–900. <https://doi.org/10.1038/ki.2008.313> PMID: 18596725
67. Weisinger RS, Stanley TK, Begg DP, Weisinger HS, Spark KJ, Jois M (2009) Angiotensin converting enzyme inhibition lowers body weight and improves glucose tolerance in C57BL/6J mice maintained on a high fat diet. *Physiol Behav* 98: 192–197. <https://doi.org/10.1016/j.physbeh.2009.05.009> PMID: 19465040
68. Roncal CA, Reungjui S, Sanchez-Lozada LG, Mu W, Sautin YY, Nakagawa T, et al. (2009) Combination of captopril and allopurinol retards fructose-induced metabolic syndrome. *Am J Nephrol* 30: 399–404. <https://doi.org/10.1159/000235731> PMID: 19696478
69. Henriksen EJ (2007) Improvement of insulin sensitivity by antagonism of the renin-angiotensin system. *American journal of physiology Regulatory, integrative and comparative physiology* 293: R974–980. <https://doi.org/10.1152/ajpregu.00147.2007> PMID: 17581838
70. Pershadsingh HA, Kurtz TW (2004) Insulin-sensitizing effects of telmisartan. *Diabetes care* 27: 1015–1015. PMID: 15047668
71. Jandeleit-Dahm KAM, Tikellis C, Reid CM, Johnston CI, Cooper ME (2005) Why blockade of the renin-angiotensin system reduces the incidence of new-onset diabetes. *Journal of hypertension* 23: 463–473. PMID: 15716683
72. Udenigwe CC, Adebisi AP, Doyen A, Li H, Bazinet L, Aluko RE (2012) Low molecular weight flaxseed protein-derived arginine-containing peptides reduced blood pressure of spontaneously hypertensive rats faster than amino acid form of arginine and native flaxseed protein. *Food chemistry* 132: 468–475. <https://doi.org/10.1016/j.foodchem.2011.11.024> PMID: 26434317
73. Doyen A, Udenigwe CC, Mitchell PL, Murette A, Aluko RE, Bazinet L (2014) Anti-diabetic and antihypertensive activities of two flaxseed protein hydrolysate fractions revealed following their simultaneous separation by electro dialysis with ultrafiltration membranes. *Food chemistry* 145: 66–76. <https://doi.org/10.1016/j.foodchem.2013.07.108> PMID: 24128450
74. Renna NF, Diez ER, Lembo C, Miatello RM (2013) Role of Cox-2 in Vascular Inflammation: An Experimental Model of Metabolic Syndrome. *Mediat Inflamm*.
75. Murasaki KM, Omori H, Sato K, Hagiwara N, Kasanuki H (2006) Metabolic syndrome contributes to aspirin-resistant thromboxane biosynthesis via inducing platelets COX-2 expression in patients with stable angina. *Circulation* 114: 71–71.
76. Hsu CK, Lee IT, Lin CC, Hsiao LD, Yang CM (2015) Sphingosine-1-phosphate mediates COX-2 expression and PGE2/IL-6 secretion via c-Src-dependent AP-1 activation. *J Cell Physiol* 230: 702–715. <https://doi.org/10.1002/jcp.24795> PMID: 25201048
77. Zhang D, Li J, Song L, Ouyang W, Gao J, Huang C (2008) A JNK1/AP-1-dependent, COX-2 induction is implicated in 12-O-tetradecanoylphorbol-13-acetate-induced cell transformation through regulating cell cycle progression. *Mol Cancer Res* 6: 165–174. <https://doi.org/10.1158/1541-7786.MCR-07-0181> PMID: 18234971
78. Park SA, Kim EH, Na HK, Surh YJ (2007) KG-135 inhibits COX-2 expression by blocking the activation of JNK and AP-1 in phorbol ester-stimulated human breast epithelial cells. *Ann N Y Acad Sci* 1095: 545–553. <https://doi.org/10.1196/annals.1397.059> PMID: 17404068
79. Sonnenberg GE, Krakower GR, Kissebah AH (2004) A novel pathway to the manifestations of metabolic syndrome. *Obes Res* 12: 180–186. <https://doi.org/10.1038/oby.2004.24> PMID: 14981209
80. Sawada Y, Sakamoto Y, Toh M, Ohara N, Hatanaka Y, Naka A, et al. (2015) Milk-derived peptide Val-Pro-Pro (VPP) inhibits obesity-induced adipose inflammation via an angiotensin-converting enzyme (ACE) dependent cascade. *Mol Nutr Food Res* 59: 2502–2510. <https://doi.org/10.1002/mnfr.201500324> PMID: 26346532
81. Guo W, Li Y, Liang W, Wong S, Apovian C, Kirkland JL, et al. (2012) Beta-mecaptoethanol suppresses inflammation and induces adipogenic differentiation in 3T3-F442A murine preadipocytes. *PLoS one* 7: e40958. <https://doi.org/10.1371/journal.pone.0040958> PMID: 22911724