

RESEARCH ARTICLE

High fat diet sensitizes fibromyalgia-like pain behaviors in mice via tumor necrosis factor alpha

Dan Tian¹, Miao Tian², Leilei Zhang¹, Peng Zhao¹, Yunfeng Cui¹, Jinlong Li^{3*}

1 Department of Anesthesiology, Second hospital of Jilin University, Changchun, China, **2** Department of Gynecology, Second hospital of Jilin University, Changchun, China, **3** Department of Gastrointestinal Surgery, Second hospital of Jilin University, Changchun, China

* jinlonglucky@gmail.com



Abstract

Fibromyalgia (FM) and obesity are closely related. However, little is known about how obesity contributes to FM. Importantly, adequate evidence has shown that tumor necrosis factor alpha (TNF- α) plays a critical role in obesity. Thus, we hypothesized that obesity-induced TNF- α release may potentiate FM-associated pain. To test this hypothesis, we investigated the role of TNF- α in the development of FM-like pain in a mouse model of acid saline injection-induced FM. Consistent with previous reports, we showed that repeated acid saline injections induced bilateral mechanical hyperalgesia, and this effect lasted for at least 4 days after acid saline injections. This phenomenon was associated with increased levels of TNF- α in plasma, muscles, and spinal cord. Furthermore, we found that 24 weeks of high fat diet treatment significantly potentiated acid saline-induced bilateral mechanical hyperalgesia. High fat diet-treated mice exhibited robustly increased levels of TNF- α in plasma, muscles, and spinal cord after acid saline injections compared with low fat diet-treated mice. Additionally, using immunofluorescence staining, we found that the number of TNF- α positive cells in dorsal root ganglion (DRG) was increased after acid saline injections, and high fat diet treatment further sensitized this increase. Finally, we reported that acid saline-induced FM-like pain behaviors were abolished in TNFRp55^{-/-} mice, confirming the critical role of TNF- α in the development of FM-like pain. Taken together, our results suggested that high fat diet treatment may sensitize acid saline-induced FM-like pain via increasing TNF- α levels in plasma, muscles, and DRG.

OPEN ACCESS

Citation: Tian D, Tian M, Zhang L, Zhao P, Cui Y, Li J (2018) High fat diet sensitizes fibromyalgia-like pain behaviors in mice via tumor necrosis factor alpha. *PLoS ONE* 13(2): e0190861. <https://doi.org/10.1371/journal.pone.0190861>

Editor: Aamir Ahmad, University of South Alabama Mitchell Cancer Institute, UNITED STATES

Received: November 22, 2017

Accepted: December 21, 2017

Published: February 14, 2018

Copyright: © 2018 Tian et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This research was funded by intramural grant (JUI2016SC5) from Second Hospital of Jilin University. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Fibromyalgia (FM) is a complex syndrome characterized by widespread chronic musculoskeletal pain which is accompanied by fatigue, sleep, memory and mood issues. FM may affect up to 8% of population worldwide [1, 2]. While the underlying disease mechanisms are still unclear, FM associated pain may be contributed by multiple factors, including abnormal regulation of central pain modulation [3], dysregulated hypothalamic-pituitary-adrenal axis (HPA) [4], and immunological vulnerability [5].

Remarkably, these aforementioned factors are often seen in obesity. Specifically, obese people typically exhibit abnormalities in the regulation of the HPA axis [6, 7]. Obese humans [8–12] and rodents [13, 14] exhibit higher sensitivity to experimentally induced nociceptive stimuli. Obese people often report to have chronic pain disorders [15–17]. Numerous studies have suggested that FM and obesity are closely related. Studies showed that up to 50% of FM patients are obese and additional 28% are overweight [18–20]. In addition, FM patients have higher body mass index (BMI) than pain-free individuals [21, 22]. These results suggested that obesity or overweight is related to greater pain/tender sensitivity in FM [18–20]. However, little is known about how obesity contributes to FM.

Recently, studies have reported increased levels of proinflammatory cytokine tumor necrosis factor alpha (TNF- α), in the serum and biopsies of FM patients [23, 24], which might be associated with disease symptoms [25–27]. Intramuscular injection of TNF- α induces muscle hyperalgesia in rats [28]. Importantly, adequate evidence has shown that TNF- α plays a critical role in obesity [29–31]. Increased levels of TNF- α in plasma and muscles were seen in obese patients and rodents [32, 33]. Thus, it is likely that obesity-induced TNF- α release may potentiate FM associated pain. To test this hypothesis in the present study, we examined the effects of high fat diet treatment on FM-like pain in a mouse model of FM. We then further assessed the role of TNF- α in the development of FM.

Materials and methods

Animals

Female TNFRp55^{-/-} (TNFR1 KO) mice were obtained from The Jackson Laboratory (ME USA). Female C57BL/6J mice were obtained from Shanghai Laboratory Animal Center (Shanghai, China). All experiments were started when female mice were 7–8 weeks old. Mice were housed in their home cages with a 12:12 light cycle with 21.0°C–23.0°C and 50%–60% humidity, and were given food and water *ad libitum*. All animal experiments were approved by the Institutional Animal Care and Use Committee of Second Hospital of Jilin University. The housing and treatment of the rats followed the guidelines of the “Guide for the Care and Use of Laboratory Rats” (Institute of Laboratory Animal Resources, Commission on Life Sciences 2011).

High fat diet treatment

Experiments were performed on C57BL/6J mice. After arrival in the animal facility, mice initially received a low-fat diet (LFD) containing 10% fat (D12450B, Research Diets Inc., NJ, USA) for 1 week until the experiment was initiated. Mice were randomly divided into control receiving a low-fat diet (LFD) and experimental group receiving a high-fat diet (HFD) containing 45% fat (D12451, Research Diets Inc., NJ, USA). Mice were then fed for 24 weeks. This treatment duration was decided based on literature [34–36]. Weight and blood glucose measurements (glucose diagnostic reagents; Sigma, St. Louis, MO, USA) were collected at the beginning of high fat diet treatment and at 24 weeks (Table 1). Mice were fasted for 3 hours prior to blood collection from the tail.

Induction of Fibromyalgia (FM)

Following the high fat diet treatment, a mouse model of fibromyalgia was induced and adapted based on the methods described in literature [37]. Acid saline (20 μ L, pH 4.0) was injected into gastrocnemius muscle when mice were anesthetized with isoflurane (1%). Acid saline injection was repeated once at day 5 after the first injection to induce FM in mice. Control mice received

Table 1. Body weights, one-day food intake amounts, and blood glucose concentrations.

	Weeks post-HFD	Low Fat Diet Group	High Fat Diet Group
24	Body weight (g)	38.4±2.2	58.7±1.9*
	Food intake (g)	3.6±0.3	7.1±0.3*
	Blood glucose (mmol/l)	7.2±0.3	9.0±0.5*

Data are expressed as Mean ± SEM, $n = 8$ per group. HFD: high fat diet. Asterisks represent the significant difference, relative to Low fat diet group. $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0190861.t001>

saline injections under the same administration schedule. After treatment, mice were subjected to behavioral tests on the next day for 4 consecutive days.

Analysis of TNF- α in plasma, muscles, and spinal cord

Mice were sacrificed after the last behavioral test, and samples of gastrocnemius muscles and spinal cord were collected. The tissue samples were then homogenized in 500 μ l of buffer solution with protease inhibitors. The concentrations of TNF- α in plasma, muscles, and spinal cord were determined by enzyme linked immunosorbent assay (ELISA), using the Mouse TNF alpha ELISA Kit (ab100747, Abcam, Shanghai, China), based on the protocol recommended by the manufacturer. The limit of detection for TNF- α was 0.1 pg/mL. Experiment was repeated three times.

Immunofluorescence staining

Mice were perfused transcardially with 4% paraformaldehyde under isoflurane anesthesia, and L3-L5 DRG tissues were collected. After being cryoprotected with 30% sucrose, the tissues were then sectioned to a thickness of 15 μ m and were post-fixed with 4% paraformaldehyde. DRG sections were then incubated with blocking solution containing 0.1% Triton X-100, 3% BSA, and 0.02% sodium azide in phosphate-buffered saline for 2 h at room temperature, followed by incubation at with the anti-TNF- α antibody (1:300; ab1793, Abcam, Shanghai, China) prepared in blocking solution at 4°C overnight. The fluorescent-labeled secondary antibodies (Abcam, 1:200) specific to the IgG species were then used. All the fluorescent imaging was performed with the same laser power and exposure time by a CKX41 microscope with an Olympus U-RFLT50 Power Supply Unit (Olympus, Tokyo, Japan). The images were quantitatively analyzed by NIH ImageJ.

Behavioral tests

Mechanical hyperalgesia was measured by testing the force of responses to stimulation with an electronic von Frey anesthesiometer (IITC, Shanghai, China). Briefly, mice were placed on a mesh floor 30 min before testing. Probes were perpendicularly applied to the central area of the hind paw with a gradual increase in pressure. After the withdrawal response, the intensity of the pressure was recorded automatically, and the results (an average of three measurements) were expressed as withdrawal threshold (g). The basal mechanical withdrawal threshold was 3.7 ± 0.1 g (mean ± SEM) before the beginning of experiment in TNFRp55 $^{-/-}$ mice and C57BL/6J mice. There was no difference of basal mechanical withdrawal thresholds between groups in the same experiment.

Statistical analyses

Data were analyzed using SPSS software (IBM, Inc., USA) and are presented as mean ± SEM. Data were analyzed by one-way analysis of variance (ANOVA) or multi-factorial ANOVA, followed by Bonferroni post hoc test, as appropriate. $P < 0.05$ was considered to be significant.

Results

High fat diet treatment increased mechanical sensitivity in a mouse model of FM

Consistent with previous reports [37–39], normal saline injections in mice (i.e., control group) did not induce ipsilateral mechanical hyperalgesia (Fig 1, n = 8). Intramuscular injections of an acidic saline (pH = 4.0) into the gastrocnemius muscle of mice induced bilateral mechanical hyperalgesia (Fig 1, n = 8). These effects lasted for at least 4 days (i.e., day 6–9). Furthermore, previous high fat diet treatment for 24 weeks significantly enhanced mechanical hyperalgesia (Fig 1, n = 8).

High fat diet treatment increased the levels of TNF- α in plasma cord and muscles in a mouse model of FM

To determine whether high fat diet altered the levels of TNF- α in plasma and muscles, we collected blood and muscle tissues after the last behavioral test. We found that intramuscular injections of an acidic saline (pH = 4.0) into the gastrocnemius muscle of mice increased the




Fig 1. Effects of high fat diet treatment on acid saline injection-induced mechanical hypersensitivity. Following 24 weeks of high fat diet treatment, mice received repeated acid saline injections into the gastrocnemius muscle. Mechanical withdrawal thresholds were then measured using an electronic von Frey device for 4 consecutive days. Data are presented as mean \pm SEM. Bonferroni correction for multiple comparisons. * represent the significant difference ($p < 0.05$) to saline control mice. # represent the significant difference ($p < 0.05$) to low fat diet mice. N = 8/group.

<https://doi.org/10.1371/journal.pone.0190861.g001>




Fig 2. Levels of TNF- α in muscles and plasma of mice. Following 24 weeks of high fat diet treatment, mice received repeated acid saline injections into the gastrocnemius muscle. Samples of plasma and gastrocnemius muscles were collected. The concentrations of TNF- α in (A) muscles and (B) plasma were determined by ELISA. The limit of detection for TNF- α was 0.1 pg/mL. *represent the significant difference ($p < 0.05$) to saline control mice. #represent the significant difference ($p < 0.05$) to low fat diet mice. $N = 8$ /group.

<https://doi.org/10.1371/journal.pone.0190861.g002>

levels of TNF- α in both plasma and muscles (Fig 2A and 2B, $n = 8$) compared with normal saline control mice. Furthermore, previous high fat diet treatment for 24 weeks significantly potentiated the levels of TNF- α in both plasma and muscles (Fig 2A and 2B, $n = 8$).

High fat diet treatment increased the levels of TNF- α in spinal cord and DRG in a mouse model of FM

To determine whether high fat diet altered the levels of TNF- α in spinal cord and DRG, we collected spinal cord and L3-L5 DRG tissues after the last behavioral test. We found that intramuscular injections of an acidic saline ($pH = 4.0$) into the gastrocnemius muscle of mice increased the levels of TNF- α in spinal cord (Fig 3, $n = 8$) compared with normal saline control mice. Furthermore, previous high fat diet treatment for 24 weeks significantly potentiated the levels of TNF- α in spinal cord (Fig 3, $n = 8$). Additionally, we conducted immunofluorescence staining to assess the levels of TNF- α in the DRG. Similarly, immunofluorescent labeling




Fig 3. Levels of TNF- α in spinal cord and DRG. Mice were sacrificed after the last behavioral test, and samples of spinal cord and L3-L5 DRG tissues were collected in separate groups of mice. The concentrations of TNF- α in spinal cord were determined by ELISA. L3-L5 DRG tissues were sectioned to a thickness of 15 μ m and underwent immunofluorescent staining. The images were quantitatively analyzed by NIH ImageJ. *represent the significant difference ($p < 0.05$) to saline control mice. #represents the significant difference ($p < 0.05$) to low fat diet mice. $N = 8$ /group.

<https://doi.org/10.1371/journal.pone.0190861.g003>




Fig 4. Effects of high fat diet treatment on acid saline injection-induced mechanical hypersensitivity in female TNFRp55^{-/-} mice. Following 24 weeks of high fat diet treatment, mice received repeated acid saline injections into the gastrocnemius muscle. Mechanical withdrawal thresholds were then measure using an electronic von Frey device for 4 consecutive days. Data are presented as mean \pm SEM. N = 8/group.

<https://doi.org/10.1371/journal.pone.0190861.g004>

visualized by green fluorescence indicated that TNF- α levels in the DRG were increased 4 days after FM induction (Fig 3, n = 8), and were further potentiated by previous high-fat diet treatment (Fig 3, n = 8).

Acid saline injections failed to induce FM-like pain behavior in TNFRp55^{-/-} mice

To explore the putative mechanisms underlying the role of TNF- α in the development of FM, we examined the mechanical hyperalgesia in TNFRp55^{-/-} mice after intramuscular injections of an acidic saline (pH = 4.0). We found that intramuscular injections of an acidic saline (pH = 4.0) into the gastrocnemius muscle of mice did not induce bilateral mechanical hyperalgesia (Fig 4, n = 8).

Discussion

In the present study, we investigated the role of TNF- α in the development of FM-like pain in a mouse model of FM. Based on published literature, we first established FM-like pain

behaviors in mice using acid saline injections. Consistent with previous reports, we showed that repeated acid saline injections induced bilateral mechanical hyperalgesia, and this effect lasted for at least 4 days after acid saline injections. This phenomenon was associated with increased levels of TNF- α in plasma, muscles, and spinal cord, suggesting TNF- α may contribute to the widespread FM-like pain behaviors. Using this mouse model of FM-like pain, we further evaluated the effects of high fat diet treatments on the status of FM-like pain. We found that 24 weeks of high fat diet treatment significantly potentiated acid saline-induced bilateral mechanical hyperalgesia. Furthermore, we found that high fat diet-treated mice exhibited robustly increased levels of TNF- α in plasma, muscles, and spinal cord after acid saline injections compared with low fat diet-treated mice. Additionally, using immunofluorescence staining, we found that the number of TNF- α positive cells in DRG was increased after acid saline injections, and high fat diet treatment further sensitized this increase. Finally, we reported that acid saline-induced FM-like pain behaviors were abolished in TNFRp55 $^{-/-}$ mice, confirming the critical role of TNF- α in the development of FM-like pain. Taken together, our results suggested that high fat diet treatment may sensitize acid saline-induced FM-like pain via increasing TNF- α levels in plasma, muscles, and DRG.

To study the disease mechanisms of FM, various rodent models have been established [40, 41]. For instance, previous studies have shown that injections of acidic saline into the muscles of mice can reliably induce pain symptoms similar to those of FM patients with minimal histopathological changes [40]. This FM mouse model exhibits long-lasting and widespread mechanical hyperalgesia without muscle damage, fatigue, sympathetic predominance or altered central sensitization [42–44]. However, unlike animal models for other types of nociceptive and neuropathic pain, it is relatively difficult to mimic FM etiologies in animal models because it is still unknown about its etiology. Therefore, the acid saline injection model of FM-like pain only mimics the symptomology and management profile of the disease rather than the mechanisms [40]. Nonetheless, this model mimics one important feature in FM that is the lack of tissue injury [40]. Furthermore, there is a high correlation between pain and comorbidities, including fatigue, depression, and anxiety in FM. Thus, ideal animal models of FM should also simulate the development of these symptoms. However, no studies have reported that depression and anxiety-like behaviors were seen in acid saline injection model of FM [40]. In fact, other FM models such as biogenic amine depletion and sound stress models have been used to simulate depression and anxiety-like behaviors in rodents [40, 41]. Therefore, future studies will be important to validate our findings in multiple animal models of FM.

Our study focused on female mice, because clinical evidence shows that FM predominantly occurs in women [45–47]. However, obesity is similarly prevalent in males and females [48]. Importantly, animal studies have indicated that TNF- α production in obese male and female may be substantially different. For instance, there is a general lack of TNF- α autoamplification in obese female mice [49], suggesting that obese male mice may even have a higher response in TNF- α production after acid saline injection. While clinical evidence and previous animal studies all suggest that studies need to be performed in both male and female animals, because sex differences are likely to be important in the development and maintenance of FM-like symptoms. However, few animal models have been tested in both males and females. Thus, it will be necessary to test the effects of high fat diet on the development of FM-like pain in male mice in the future.

Previous studies indicated that FM-like syndrome may result from activation of vanilloid type 1 receptors (TRPV1s) or acid sensing ion channels (ASICs), which are voltage-insensitive cationic channels gated by extracellular protons [50]. Importantly, pH levels are often reduced in peripheral tissues of FM patients [51]. Furthermore, it has been shown that protons can activate the terminals of nociceptors. Electrophysiology recordings of DRG neurons have

demonstrated that acid can activate TRPV1, leading to an up-regulation of CGRP expression in DRG via CaMK-CREB cascade, a series of events that may be associated with inflammatory pain [52]. As decreases in pH are associated with inflammation, it is likely that inflammatory mediators, such as TNF α , could sensitize the pH response in muscles or DRG. In fact, TRPV1 shows an increase in expression after prior treatment with TNF- α in cultured fibroblast-like synoviocytes [53], suggesting a direct interaction of TNF- α and TRPV1. Interestingly, in contrast to TRPV1, application of TNF- α failed to alter the mRNA levels of ASIC1, ASIC2 and ASIC3 in the DRG, suggesting TNF- α may not modulate expression of ASICs in DRG [54, 55]. However, similar to TRPV1, TNF- α mediated inflammation may reduce the pH and trigger the activation of ASICs leading to enhanced muscle pain. Thus, given that our results clearly showed an increase in TNF- α level in DRG, future experiments should investigate an interaction between inflammatory mediators and the pH response in DRG neurons.

Adding to existing literature, our study first reported that TNFRp55 $^{-/-}$ mice is resistant to the development of FM-like pain behaviors after repeated acid saline injections. In fact, previous studies have demonstrated that multiple molecules are involved in FM-like pain. Besides ASICs and TRPV1, these molecules including N-methyl-D-aspartate receptors (NMDARs), voltage-gated calcium channels and voltage-activated sodium channels, as well as substance P have been implicated in mouse models of FM pain [37, 42, 56–58]. Importantly, TNF- α has been shown to interact with these molecules. Specifically, TNF- α can increase NMDAR function in spinal cord neurons [59]. In nociceptive DRG neurons, TNF- α decreases voltage-gated calcium channel currents, and increases voltage-activated sodium channels currents [60]. Furthermore, substance P release is enhanced in DRG cultures after chronic TNF- α treatment [61]. These results together suggest that TNF- α may play a central role in mouse models of FM pain.

In summary, our study was the first to report the role of TNF- α in the development of FM-like pain. Consistent with a lot of previous studies that inhibition of TNF- α -mediate neuroinflammation signaling can improve pain-like behavior in various mouse models of pain, these findings together suggested that TNF- α may have overlapping mechanisms that contribute to attenuation of distinct types of pain. In fact, peripheral synaptic transmission from DRG neurons to the spinal cord dorsal horn is crucial for pain signaling [62, 63], which is then transferred into the brain for processing the pain sensation and responses [64]. In our study, we reported that obesity sensitized acid saline-induced TNF- α production in the DRG. Thus, it is likely that increased DRG TNF- α production in obesity may contribute to the maintenance of FM-like pain behaviors. Future studies will be also important to explore the effects of TNF- α inhibition on the development of acid saline-induced FM-like pain behaviors, and dissect the putative role of TNF- α subpopulations (muscle, plasma, or spinal cord) in modulations of this phenomenon. This line of research will not only be important in advancing our knowledge in the neuropathological mechanisms of FM, but may provide an effective pharmacotherapeutic intervention in treating FM patients.

Acknowledgments

This research was funded by intramural grant (JUI2016SC5) from Second Hospital of Jilin University.

Author Contributions

Conceptualization: Jinlong Li.

Data curation: Miao Tian, Leilei Zhang.

Formal analysis: Miao Tian, Leilei Zhang, Yunfeng Cui.

Funding acquisition: Jinlong Li.

Investigation: Dan Tian, Peng Zhao.

Methodology: Dan Tian, Peng Zhao, Jinlong Li.

Project administration: Jinlong Li.

Resources: Jinlong Li.

Supervision: Jinlong Li.

Validation: Leilei Zhang.

Writing – original draft: Dan Tian, Yunfeng Cui.

Writing – review & editing: Jinlong Li.

References

1. English B. Neural and psychosocial mechanisms of pain sensitivity in fibromyalgia. *Pain Management Nursing*. 2014; 15(2):530–8. <https://doi.org/10.1016/j.pmn.2012.07.009> PMID: 24882030
2. Clauw DJ. Fibromyalgia: a clinical review. *Jama*. 2014; 311(15):1547–55. <https://doi.org/10.1001/jama.2014.3266> PMID: 24737367
3. Arendt-Nielsen L, Graven-Nielsen T. Central sensitization in fibromyalgia and other musculoskeletal disorders. *Curr Pain Headache Rep*. 2003; 7(5):355–61. Epub 2003/08/30. PMID: 12946288.
4. Okifuji A, Turk DC. Stress and psychophysiological dysregulation in patients with fibromyalgia syndrome. *Appl Psychophysiol Biofeedback*. 2002; 27(2):129–41. Epub 2002/09/11. PMID: 12206047.
5. Wallace DJ. Is there a role for cytokine based therapies in fibromyalgia. *Current pharmaceutical design*. 2006; 12(1):17–22. PMID: 16454720
6. Bjorntorp P, Holm G, Rosmond R. Hypothalamic arousal, insulin resistance and Type 2 diabetes mellitus. *Diabet Med*. 1999; 16(5):373–83. Epub 1999/05/26. PMID: 10342336.
7. Rosmond R, Dallman MF, Bjorntorp P. Stress-related cortisol secretion in men: relationships with abdominal obesity and endocrine, metabolic and hemodynamic abnormalities. *The Journal of Clinical Endocrinology & Metabolism*. 1998; 83(6):1853–9.
8. Torensma B, Oudejans L, van Velzen M, Swank D, Niesters M, Dahan A. Pain sensitivity and pain scoring in patients with morbid obesity. *Surg Obes Relat Dis*. 2017; 13(5):788–95. Epub 2017/02/22. doi: S1550-7289(17)30029-1 [pii] <https://doi.org/10.1016/j.soard.2017.01.015> PMID: 28216116.
9. Eslami V, Katz MJ, White RS, Sundermann E, Jiang JM, Ezzati A, et al. Pain Intensity and Pain Interference in Older Adults: Role of Gender, Obesity and High-Sensitivity C-Reactive Protein. *Gerontology*. 2017; 63(1):3–12. Epub 2016/08/04. doi: 000446651 [pii] <https://doi.org/10.1159/000446651> PMID: 27486843; PubMed Central PMCID: PMC5154794 conflicts of interest to report.
10. Price RC, Asenjo JF, Christou NV, Backman SB, Schweinhardt P. The role of excess subcutaneous fat in pain and sensory sensitivity in obesity. *Eur J Pain*. 2013; 17(9):1316–26. Epub 2013/04/12. <https://doi.org/10.1002/j.1532-2149.2013.00315.x> PMID: 23576531.
11. Khimich S. Level of sensitivity of pain in patients with obesity. *Acta Chir Hung*. 1997; 36(1–4):166–7. Epub 1997/01/01. PMID: 9408331.
12. Zahorska-Markiewicz B, Zych P, Kucio C. Pain sensitivity in obesity. *Acta Physiol Pol*. 1988; 39(3):183–7. Epub 1988/05/01. PMID: 3227867.
13. Ramzan I, Wong BK, Corcoran GB. Pain sensitivity in dietary-induced obese rats. *Physiology & behavior*. 1993; 54(3):433–5.
14. Okifuji A, Hare BD. The association between chronic pain and obesity. *Journal of pain research*. 2015; 8:399. <https://doi.org/10.2147/JPR.S55598> PMID: 26203274
15. Lake JK, Power C, Cole TJ. Back pain and obesity in the 1958 British birth cohort. cause or effect? *J Clin Epidemiol*. 2000; 53(3):245–50. Epub 2000/04/13. doi: S0895-4356(99)00155-9 [pii]. PMID: 10760633.
16. Peres MF, Lerario DD, Garrido AB, Zukerman E. Primary headaches in obese patients. *Arq Neuropsiquiatr*. 2005; 63(4):931–3. Epub 2006/01/10. <https://doi.org/S0004-282X2005000600005> PMID: 16400407.

17. Peltonen M, Lindroos AK, Torgerson JS. Musculoskeletal pain in the obese: a comparison with a general population and long-term changes after conventional and surgical obesity treatment. *Pain*. 2003; 104(3):549–57. PMID: [12927627](#)
18. Neumann L, Lerner E, Glazer Y, Bolotin A, Shefer A, Buskila D. A cross-sectional study of the relationship between body mass index and clinical characteristics, tenderness measures, quality of life, and physical functioning in fibromyalgia patients. *Clinical rheumatology*. 2008; 27(12):1543–7. <https://doi.org/10.1007/s10067-008-0966-1> PMID: [18622575](#)
19. Okifuji A, Bradshaw DH, Olson C. Evaluating obesity in fibromyalgia: neuroendocrine biomarkers, symptoms, and functions. *Clinical rheumatology*. 2009; 28(4):475–8. <https://doi.org/10.1007/s10067-009-1094-2> PMID: [19172342](#)
20. Yunus MB, Arslan S, Aldag JC. Relationship between body mass index and fibromyalgia features. *Scandinavian journal of rheumatology*. 2002; 31(1):27–31. PMID: [11922197](#)
21. Elert J, Kendall SA, Larsson B, Mansson B, Gerdle B. Chronic pain and difficulty in relaxing postural muscles in patients with fibromyalgia and chronic whiplash associated disorders. *J Rheumatol*. 2001; 28(6):1361–8. Epub 2001/06/21. PMID: [11409132](#).
22. Bennett RM, Jones J, Turk DC, Russell IJ, Matallana L. An internet survey of 2,596 people with fibromyalgia. *BMC musculoskeletal disorders*. 2007; 8(1):27.
23. Bennett R. The Fibromyalgia Impact Questionnaire (FIQ): a review of its development, current version, operating characteristics and uses. *Clinical and experimental rheumatology*. 2005; 23(5):S154.
24. Salemi S, Rethage J, Wollina U, Michel BA, Gay RE, Gay S, et al. Detection of interleukin 1beta (IL-1beta), IL-6, and tumor necrosis factor-alpha in skin of patients with fibromyalgia. *The Journal of rheumatology*. 2003; 30(1):146–50. PMID: [12508404](#)
25. Bernstein MS, Costanza MC, Morabia A. Association of physical activity intensity levels with overweight and obesity in a population-based sample of adults. *Preventive medicine*. 2004; 38(1):94–104. PMID: [14672646](#)
26. Mattsson P. Migraine headache and obesity in women aged 40–74 years: a population-based study. *Cephalalgia*. 2007; 27(8):877–80. Epub 2007/07/20. doi: CHA1360 [pii] <https://doi.org/10.1111/j.1468-2982.2007.01360.x> PMID: [17635528](#).
27. Gilbert GJ. Obesity and migraine: a population study. *Neurology*. 2007; 68(3):241; author reply Epub 2007/01/18. PMID: [17228422](#).
28. Schäfers M, Sorkin LS, Sommer C. Intramuscular injection of tumor necrosis factor-alpha induces muscle hyperalgesia in rats. *Pain*. 2003; 104(3):579–88. PMID: [12927630](#)
29. Cawthorn WP, Sethi JK. TNF- α and adipocyte biology. *FEBS letters*. 2008; 582(1):117–31. <https://doi.org/10.1016/j.febslet.2007.11.051> PMID: [18037376](#)
30. Schreyer SA, Chua SC Jr, LeBoeuf RC. Obesity and diabetes in TNF-alpha receptor-deficient mice. *Journal of Clinical Investigation*. 1998; 102(2):402. <https://doi.org/10.1172/JCI2849> PMID: [9664082](#)
31. Tzanavari T, Giannogonas P, Karalis KP. TNF-alpha and obesity. *Curr Dir Autoimmun*. 2010; 11:145–56. Epub 2010/02/23. <https://doi.org/10.1159/000289203> [pii]. PMID: [20173393](#).
32. Plomgaard P, Nielsen A, Fischer C, Mortensen O, Broholm C, Penkowa M, et al. Associations between insulin resistance and TNF- α in plasma, skeletal muscle and adipose tissue in humans with and without type 2 diabetes. *Diabetologia*. 2007; 50(12):2562–71. <https://doi.org/10.1007/s00125-007-0834-6> PMID: [17928988](#)
33. Valerio A, Cardile A, Cozzi V, Bracale R, Tedesco L, Pisconti A, et al. TNF- α downregulates eNOS expression and mitochondrial biogenesis in fat and muscle of obese rodents. *Journal of Clinical Investigation*. 2006; 116(10):2791. <https://doi.org/10.1172/JCI28570> PMID: [16981010](#)
34. Lin S, Thomas T, Storlien L, Huang X. Development of high fat diet-induced obesity and leptin resistance in C57Bl/6J mice. *International journal of obesity*. 2000; 24(5):639. PMID: [10849588](#)
35. Weisberg SP, Hunter D, Huber R, Lemieux J, Slaymaker S, Vaddi K, et al. CCR2 modulates inflammatory and metabolic effects of high-fat feeding. *Journal of Clinical Investigation*. 2006; 116(1):115. <https://doi.org/10.1172/JCI24335> PMID: [16341265](#)
36. Abu-Elheiga L, Oh W, Kordari P, Wakil SJ. Acetyl-CoA carboxylase 2 mutant mice are protected against obesity and diabetes induced by high-fat/high-carbohydrate diets. *Proceedings of the National Academy of Sciences*. 2003; 100(18):10207–12.
37. Chen W-N, Lee C-H, Lin S-H, Wong C-W, Sun W-H, Wood JN, et al. Roles of ASIC3, TRPV1, and Na V 1.8 in the transition from acute to chronic pain in a mouse model of fibromyalgia. *Molecular pain*. 2014; 10(1):40.
38. Yen L-T, Hsieh C-L, Hsu H-C, Lin Y-W. Targeting ASIC3 for Relieving Mice Fibromyalgia Pain: Roles of Electroacupuncture, Opioid, and Adenosine. *Scientific Reports*. 2017; 7.

39. Lin J-G, Hsieh C-L, Lin Y-W. Analgesic effect of electroacupuncture in a mouse fibromyalgia model: roles of TRPV1, TRPV4, and pERK. *PloS one*. 2015; 10(6):e0128037. <https://doi.org/10.1371/journal.pone.0128037> PMID: 26043006
40. DeSantana JM, da Cruz KM, Sluka KA. Animal models of fibromyalgia. *Arthritis research & therapy*. 2013; 15(6):222.
41. McLean SA, Clauw DJ. Biomedical models of fibromyalgia. *Disability and Rehabilitation*. 2005; 27(12):659–65. <https://doi.org/10.1080/09638280400009048> PMID: 16012058
42. Vilin YY, Peters CH, Ruben PC. Acidosis differentially modulates inactivation in Nav1. 2, Nav1. 4, and Nav1. 5 channels. *Frontiers in pharmacology*. 2012; 3.
43. Chen C-C, Zimmer A, Sun W-H, Hall J, Brownstein MJ, Zimmer A. A role for ASIC3 in the modulation of high-intensity pain stimuli. *Proceedings of the National Academy of Sciences*. 2002; 99(13):8992–7.
44. Chen W-H, Hsieh C-L, Huang C-P, Lin T-J, Tzen JT, Ho T-Y, et al. Acid-sensing ion channel 3 mediates peripheral anti-hyperalgesia effects of acupuncture in mice inflammatory pain. *Journal of biomedical science*. 2011; 18(1):82.
45. Rollman GB, Lautenbacher S. Sex differences in musculoskeletal pain. *The Clinical journal of pain*. 2001; 17(1):20–4. PMID: 11289085
46. Fillingim RB. Sex, gender, and pain: women and men really are different. *Current Pain and Headache Reports*. 2000; 4(1):24–30.
47. Fillingim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL. Sex, gender, and pain: a review of recent clinical and experimental findings. *The journal of pain*. 2009; 10(5):447–85. <https://doi.org/10.1016/j.jpain.2008.12.001> PMID: 19411059
48. Kelly T, Yang W, Chen C-S, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *International journal of obesity*. 2008; 32(9):1431–7. <https://doi.org/10.1038/ijo.2008.102> PMID: 18607383
49. Neels JG, Pandey M, Hotamisligil GS, Samad F. Autoamplification of tumor necrosis factor- α : a potential mechanism for the maintenance of elevated tumor necrosis factor- α in male but not female obese mice. *The American journal of pathology*. 2006; 168(2):435–44. <https://doi.org/10.2353/ajpath.2006.050699> PMID: 16436658
50. Wemmie JA, Price MP, Welsh MJ. Acid-sensing ion channels: advances, questions and therapeutic opportunities. *Trends in neurosciences*. 2006; 29(10):578–86. <https://doi.org/10.1016/j.tins.2006.06.014> PMID: 16891000
51. Staud R. Peripheral pain mechanisms in chronic widespread pain. *Best Pract Res Clin Rheumatol*. 2011; 25(2):155–64. Epub 2011/11/19. <https://doi.org/10.1016/j.berh.2010.01.010> S1521-6942(11)00003-9 [pii]. PMID: 22094192; PubMed Central PMCID: PMC3220877.
52. Nakanishi M, Hata K, Nagayama T, Sakurai T, Nishisho T, Wakabayashi H, et al. Acid activation of Trpv1 leads to an up-regulation of calcitonin gene-related peptide expression in dorsal root ganglion neurons via the CaMK-CREB cascade: a potential mechanism of inflammatory pain. *Molecular biology of the cell*. 2010; 21(15):2568–77. <https://doi.org/10.1091/mbc.E10-01-0049> PMID: 20534813
53. Kochukov MY, McNearney TA, Yin H, Zhang L, Ma F, Ponomareva L, et al. Tumor necrosis factor-alpha (TNF- α) enhances functional thermal and chemical responses of TRP cation channels in human synoviocytes. *Molecular pain*. 2009; 5(1):49.
54. Mamet J, Baron A, Lazdunski M, Voilley N. Proinflammatory mediators, stimulators of sensory neuron excitability via the expression of acid-sensing ion channels. *Journal of Neuroscience*. 2002; 22(24):10662–70. PMID: 12486159
55. Voilley N, de Weille J, Mamet J, Lazdunski M. Nonsteroid anti-inflammatory drugs inhibit both the activity and the inflammation-induced expression of acid-sensing ion channels in nociceptors. *Journal of Neuroscience*. 2001; 21(20):8026–33. PMID: 11588175
56. Sluka KA, Price MP, Breese NM, Stucky CL, Wemmie JA, Welsh MJ. Chronic hyperalgesia induced by repeated acid injections in muscle is abolished by the loss of ASIC3, but not ASIC1. *Pain*. 2003; 106(3):229–39. PMID: 14659506
57. Chen W-N, Chen C-C. Acid mediates a prolonged antinociception via substance P signaling in acid-induced chronic widespread pain. *Molecular pain*. 2014; 10(1):30.
58. Chen W-K, Liu IY, Chang Y-T, Chen Y-C, Chen C-C, Yen C-T, et al. Cav3. 2 T-type Ca²⁺ channel-dependent activation of ERK in paraventricular thalamus modulates acid-induced chronic muscle pain. *Journal of Neuroscience*. 2010; 30(31):10360–8. <https://doi.org/10.1523/JNEUROSCI.1041-10.2010> PMID: 20685979
59. Zhang L, Berta T, Xu ZZ, Liu T, Park JY, Ji RR. TNF- α contributes to spinal cord synaptic plasticity and inflammatory pain: distinct role of TNF receptor subtypes 1 and 2. *Pain*. 2011; 152(2):419–27.

Epub 2010/12/17. <https://doi.org/10.1016/j.pain.2010.11.014> S0304-3959(10)00699-8 [pii]. PMID: 21159431; PubMed Central PMCID: PMC3022092.

60. Czeschik JC, Hagenacker T, Schafers M, Busselberg D. TNF-alpha differentially modulates ion channels of nociceptive neurons. *Neurosci Lett*. 2008; 434(3):293–8. Epub 2008/03/04. <https://doi.org/10.1016/j.neulet.2008.01.070> S0304-3940(08)00153-5 [pii]. PMID: 18314270.
61. Lin YT, Ro LS, Wang HL, Chen JC. Up-regulation of dorsal root ganglia BDNF and trkB receptor in inflammatory pain: an in vivo and in vitro study. *J Neuroinflammation*. 2011; 8:126. Epub 2011/10/01. <https://doi.org/10.1186/1742-2094-8-126> 1742-2094-8-126 [pii]. PMID: 21958434; PubMed Central PMCID: PMC3203068.
62. Wood JN. Chapter 5 Molecular mechanisms of nociception and pain. *Handb Clin Neurol*. 2006; 81:49–59. Epub 2008/09/24. [https://doi.org/10.1016/S0072-9752\(06\)80009-6](https://doi.org/10.1016/S0072-9752(06)80009-6) S0072-9752(06)80009-6 [pii]. PMID: 18808827.
63. Julius D, Basbaum AI. Molecular mechanisms of nociception. *Nature*. 2001; 413(6852):203–10. Epub 2001/09/15. <https://doi.org/10.1038/35093019> [pii]. PMID: 11557989.
64. Cregg R, Momin A, Rugiero F, Wood JN, Zhao J. Pain channelopathies. *The Journal of Physiology*. 2010; 588(11):1897–904. <https://doi.org/10.1113/jphysiol.2010.187807> PMID: 20142270