

RESEARCH ARTICLE

Coral reef fishes exhibit beneficial phenotypes inside marine protected areas

Robert Y. Fidler^{1*}, Jessica Carroll², Kristen W. Rynerson², Danielle F. Matthews¹, Ralph G. Turingan¹

1 Department of Biological Sciences, Florida Institute of Technology, Melbourne, Florida, United States of America, **2** Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission, St. Petersburg, Florida, United States of America

* rfidler2011@my.fit.edu



OPEN ACCESS

Citation: Fidler RY, Carroll J, Rynerson KW, Matthews DF, Turingan RG (2018) Coral reef fishes exhibit beneficial phenotypes inside marine protected areas. PLoS ONE 13(2): e0193426. <https://doi.org/10.1371/journal.pone.0193426>

Editor: Heather M. Patterson, Department of Agriculture and Water Resources, AUSTRALIA

Received: August 29, 2017

Accepted: February 9, 2018

Published: February 22, 2018

Copyright: © 2018 Fidler et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This research was funded through a U.S. Student Fulbright Fellowship (<https://us.fulbrightonline.org/fulbright-us-student-program>) in association with the Philippine-American Education Foundation (<https://fulbright.org.ph/>) awarded to RYF (ID #34143036). The funders had no role in study design, data collection, and analysis, decision to publish, or preparation of the manuscript.

Abstract

Human fishing effort is size-selective, preferentially removing the largest individuals from harvested stocks. Intensive, size-specific fishing mortality induces directional shifts in phenotypic frequencies towards the predominance of smaller and earlier-maturing individuals, which are among the primary causes of declining fish biomass. Fish that reproduce at smaller size and younger age produce fewer, smaller, and less viable larvae, severely reducing the reproductive capacity of harvested populations. Marine protected areas (MPAs) are extensively utilized in coral reefs for fisheries management, and are thought to mitigate the impacts of size-selective fishing mortality and supplement fished stocks through larval export. However, empirical evidence of disparities in fitness-relevant phenotypes between MPAs and adjacent fished reefs is necessary to validate this assertion. Here, we compare key life-history traits in three coral-reef fishes (*Acanthurus nigrofuscus*, *Ctenochaetus striatus*, and *Parupeneus multifasciatus*) between MPAs and fished reefs in the Philippines. Results of our analyses support previous hypotheses regarding the impacts of MPAs on phenotypic traits. Asymptotic length (L_{inf}) and growth rates (K) differed between conspecifics in MPAs and fished reefs, with protected populations exhibiting phenotypes that are known to confer higher fecundity. Additionally, populations demonstrated increases in length at 50% maturity (L_{50}) inside MPAs compared to adjacent areas, although age at 50% maturity (A_{50}) did not appear to be impacted by MPA establishment. Shifts toward advantageous phenotypes were most common in the oldest and largest MPAs, but occurred in all of the MPAs examined. These results suggest that MPAs may provide protection against the impacts of size-selective harvest on life-history traits in coral-reef fishes.

Introduction

In addition to driving population declines, increasing evidence suggests that fishing induces widespread alterations to life-history characteristics within harvested stocks. Rapid phenotypic shifts have been driven by the non-random nature of human fishing effort, which preferentially removes the oldest and largest individuals from any given population [1–2]. Intensive

Competing interests: The authors have declared that no competing interests exist.

size-selective mortality generates directional selection pressure that favors individuals with smaller body-sizes [3–9] and those that reach reproductive maturity at smaller body-size and younger age [10–13]. Selective pressures from fishing can be strong enough to produce population-wide declines in asymptotic length, growth rate, and size- and age-at-maturation within contemporary timescales [4].

In many fishes, decreased female body-size in mature females correlates with reduced larval size-at-hatch, growth rate, feeding rate, and viability [14–15]. Additionally, females that reach sexual maturity at a young age breed for shorter periods and produce fewer, smaller eggs compared to conspecifics that mature later in life [16–17]. Declines in the abundance of old, large, highly-fecund individuals are therefore among the primary drivers of reductions in harvestable biomass in exploited stocks [18]. Decreased population size coupled with diminished reproductive capability can result in fishery collapse, best exemplified by the near disappearance of Atlantic cod (*Gadus morhua*) in southern Labrador and eastern Newfoundland, Canada, in the late 1980s; an event preceded by the widespread induction of reduced size- and age-at-maturation [7–8, 19–20]. Highlighting the long-term consequences of human-induced alterations to stock characteristics, *G. morhua* populations in the region have struggled to recover despite a fishing moratorium enacted in 1992 [21]. Only recently, after 25 years of fishing regulation and protection, have some *G. morhua* stocks displayed evidence of a significant rebound in population sizes [22], although they remain well below historical averages. Considering the rapid and devastating impacts of size-selective harvesting on the long-term viability of harvested populations, it is critical that life-history shifts be included in the design and assessment of programs aimed at managing fishery resources [23–24].

At present, a common component of coral-reef fishery management efforts are no-take marine reserves or marine protected areas (MPAs); spatially explicit zones in which fishing and other exploitative activities are prohibited. Between ~6,000–9,000 MPAs have been established worldwide, covering ~14.2 million km² (3.9%) of the total ocean surface, with ~1.93 million km² (0.53% of ocean surface) being fully no-take areas [25]. As no-fishing zones, MPAs eliminate size-selective fishing effort and the resulting directional selection pressure, theoretically arresting further proliferation of fishing-induced phenotypes within their borders [26–28]. Deleterious shifts in phenotypic frequencies may even be reversed by MPAs, as ecological forces generate selection gradients that favor larger body-size and greater age- and size-at-maturation [26–27, 29–30]. Controlled selection experiments have demonstrated that populations are able to restore historical body-size frequency distributions in 6–12 generations after the cessation of size-selective harvest [31]. It is unknown, however, if the processes observed in laboratory trials can occur in wild fisheries, or if they will be mitigated by natural predation and external influences, including habitat degradation and fishing pressure, outside of protected areas [7, 32].

The ability of MPAs to promote increased reproductive output per unit biomass is critical to their ability to meet their stated biological and economic objectives. Among the most important proposed benefits of MPAs is recruitment subsidy, the process by which excess larvae produced by healthy populations inside MPAs are transported to adjacent areas, increasing ecological resilience and improving fishery yields [33]. Consequently, MPAs must not only increase fish density, but also the relative fecundity of individuals within their borders, as relatively small populations inside MPAs must be able to supplement individuals lost to fishing in much larger spatial areas. Assessments of MPAs have demonstrated increased fish density, biomass, species diversity, and average body-size and age within MPAs compared to fished reefs [34–38]. Additionally, beneficial shifts in population structure towards the predominance of large-bodied individuals within MPAs have been observed in a multitude of reefs worldwide [35–36, 39]. However, it is unclear whether these observed trends are primarily the result of

increased longevity in protected populations or are driven by shifts in phenotypic characteristics inside MPAs. Although studies have indicated that average batch fecundity may be increased inside MPAs compared to adjacent areas [40], egg production has also been shown to vary spatially due to a variety of influences unrelated to fishing pressure including water temperature and prey availability [41–42]. As a result, the ubiquity with which MPAs promote shifts in phenotypic frequencies within their borders remains in question.

In light of the growing anthropogenic pressure being placed on marine fisheries and the increasing use of MPAs as a management tool, a more thorough understanding of the effects of MPAs on fishery resources is essential. This study was designed to assess disparities in fitness-relevant phenotypes between protected and fished populations of three exploited reef-fish species in the Philippines. To do this, we use empirical data to compare the average asymptotic body size, growth rate, and length and age at 50% maturity of conspecific reef fish populations inside and outside of MPA borders.

Materials and methods

To test for disparities in fitness-relevant phenotypes of reef fishes between protected and harvested populations, individuals of three coral-resident fish species were collected in three MPAs and adjacent, fished reefs. Experimental fishes consisted of species directly targeted by fishers for food in the region: *Acanthurus nigrofuscus* (brown surgeonfish), *Ctenochaetus striatus* (lined bristletooth), and *Parupeneus multifasciatus* (manybar goatfish). Collections were conducted in Masinloc, Zambales Province, Luzon, Philippines (15° 31' 40"N, 119° 57' 38"E) between March and June of 2015. Collections were conducted in fished reefs and in three MPAs of varying size and age: Bani MPA (est. 2006; 50 ha), San Salvador MPA (est. 1989; 127 ha), Taklobo Farm MPA (est. 1989; 2 ha), (Fig 1). All MPAs are fully no-take zones and strongly enforced, with guards standing watch 18–24 hours a day with access to boats to confront any illegal fishing activity. Specimens of each species were collected throughout their observed body-size range at each site via scuba and spearfishing equipment. To ensure that individuals collected outside of MPAs were exposed to continuous fishing pressure, all collections in “fished reefs” took place at least 300m outside of MPA boundaries. Fish collection, handling, and otolith extraction and processing were conducted pursuant to the animal-handling protocol included in the Republic of the Philippines, Department of Agriculture Gratuitous Permit GP-0096-15 issued on June 01, 2015 in Quezon City, Philippines.

After collection, fish were sacrificed by immersion in ice while being transported to the laboratory for processing. Species-level identification was then confirmed, and the standard length (mm) of each fish was recorded. Gonads were removed, sexed, and staged using a protocol adapted from Murua et al. [43]. Gonads were classified into three stages: Stage 1 (immature); Stage 2 (sexually mature, but not fully engorged); or Stage 3 (mature and ripe, preparing to spawn). Both sagittal otoliths were removed, cleaned of adhering tissue, and suspended in an araldite epoxy mixture (82% Araldite 502, specific gravity 1.13, Electron Microscopy Sciences Inc.; 12% Hardener HY 956, Vantico Inc.), which was hardened at room temperature overnight. For consistency, the left otolith was used for analysis whenever possible. After hardening, three 0.45–0.70mm sections were cut from the center of each otolith using a Buehler Isomet low-speed saw. Following sectioning, each otolith was secured to a slide using a toluene-based liquid mounting medium, and subsequently aged on a Leica MZ9.5 stereomicroscope under transmitted light. Ageing was conducted by enumerating opaque rings (counted as annuli) along the ventral edge of the sulcus acousticus. Alternating patterns of translucent and opaque rings have been validated as annuli in *A. nigrofuscus* [44] and *C. striatus* [45]. Due to their structure, ageing analyses were conducted on whole otoliths rather than sections for *P.*




Fig 1. Sites of MPAs and fished reefs where collections of three coral-reef resident fishes occurred.

<https://doi.org/10.1371/journal.pone.0193426.g001>

multifasciatus. Otoliths were submerged in water within a black petri dish, and analyzed using a Leica MZ9.5 stereomicroscope under reflected light. Annual deposition is unvalidated for *P. multifasciatus*; however, the seasonality of opaque zone formation was consistent with annual deposition.

To test for disparities in asymptotic lengths and growth rates between populations, Von Bertalanffy growth models were developed for populations collected within each MPA and associated fished reefs. The average length (l) of fish at time (t) was calculated as:

$$l_t = L_{inf}(1 - e^{-k(t-t_0)})$$

where t_0 is the theoretical time at which length = 0, L_{inf} is the mean maximum length or “asymptotic length” where growth = 0, and K is the rate (years⁻¹) at which L_{inf} is attained. Von Bertalanffy growth models were compared using the likelihood-ratio method [46]. This analysis tests for differences in the average asymptotic length (L_{inf}), growth rate (K), and time at length 0 (t_0) between populations. Due to variations in the availability of specimens less than a year old and limited sample sizes within MPAs, Von Bertalanffy growth models occasionally produced unrealistic growth patterns at early ages within individual sites. Consequently, an additional individual of age “0” and size “0” was added to each population in order to properly set growth functions. Statistical analyses were conducted using the *fishmethods* package in R statistical software [47].

To estimate the length (L_{50}) and age (A_{50}) at which 50% of individuals were mature within each population, logistic regression models were fit to data regarding the percentage of

individuals found to be mature within age classes (year) and size classes (every 2mm). In addition, 95% confidence intervals around these estimates were created by bootstrapping with 999 iterations. Analyses were conducted using the *glm* function and *FSA* package in R statistical software [48].

Results

A total of 569 specimens were collected from MPAs and associated fished reefs (S1 Table). Populations of *P. multifasciatus* were not robust enough in the Taklobo Farm MPA to conduct collections, and therefore were only collected from the Bani and San Salvador MPAs. In the following descriptions of statistics, the term “MPA” will be used when referring to protected populations, whereas “FR” will be used when referring to fished reefs.

Von Bertalanffy growth models revealed considerable variation in life-history traits between species and sites. Populations of *A. nigrofuscus*, *C. striatus*, and *P. multifasciatus* all exhibited disparities in growth parameters between at least one MPA compared to fished reefs. Significant differences were concentrated in the two oldest MPAs in the region: San Salvador and Taklobo Farm (both 26 years). Differential life-history traits were, however, occasionally observed between populations inside and outside of the Bani MPA. Statistical results of comparisons of asymptotic lengths (L_{inf}), growth rates (K), and time at length 0 (t_0) for all MPA-fished reef pairs are listed in Table 1.

Significant increases in L_{inf} were observed in populations of *A. nigrofuscus* inside MPAs compared to fished reefs for both the San Salvador MPA and the Taklobo Farm MPA (Fig 2). Asymptotic lengths inside these MPAs were 11mm (14%) and 6mm (8%) greater than populations in fished reefs, respectively. Populations of *C. striatus* exhibited higher L_{inf} values in the Bani and Taklobo Farm MPAs, with asymptotic lengths being 13mm (11%) and 7mm (6%) greater than those in fished reefs (Fig 3). Populations of *P. multifasciatus* demonstrated significantly higher asymptotic length inside the San Salvador MPA, with protected stocks reaching body-sizes 38mm (28%) greater than those in fished reefs (Fig 4). Shifts in the growth rate (K) of conspecifics between MPAs and fished reefs occurred only for *C. striatus* and *P. multifasciatus*, with both species exhibiting lower K values inside the San Salvador MPA compared to fished reefs (Figs 3 and 4). Slower growth rates were rarely associated with significant differences in asymptotic length, with shifts in both life-history traits occurring simultaneously only once (*P. multifasciatus*, San Salvador MPA).

Observed values of age- and length-at-maturation are illustrated in S1–S3 Figs, and the results of logistic regression models for length (L_{50}) and age (A_{50}) at 50% maturity can be

Table 1. Results of statistical comparisons of asymptotic length (L_{inf} , SL, mm), growth rates (K), and time-at-age-0 (t_0) between populations inside MPAs (MPA) and associated fished reefs (FR), using the likelihood ratio method of Kimura [46]. Length values have been rounded to the nearest mm. Values for each variable in MPAs and fished reefs are provided, along with χ^2 statistics and associated p-values. Significant results are indicated by bold font.

Species	Trait	Bani				San Salvador				Taklobo Farm			
		MPA	FR	X2	P	MPA	FR	X2	P	MPA	FR	X2	P
<i>Acanthurus</i>	L_{inf}	80	78	0.48	0.488	89	78	11.52	0.001	84	78	5.60	0.018
<i>nigrofuscus</i>	K	1.17	0.89	1.98	0.159	0.82	0.89	0.13	0.718	0.98	0.89	0.21	0.647
	t_0	-0.05	-0.28	3.36	0.067	-0.18	-0.28	0.35	0.554	-0.11	-0.28	1.32	0.251
<i>Ctenochaetus</i>	L_{inf}	132	119	6.00	0.014	125	119	2.56	0.110	126	119	4.82	0.028
	K	0.66	0.72	0.05	0.823	0.49	0.72	3.97	0.046	0.69	0.72	0.06	0.806
	t_0	-0.16	-0.12	0.68	0.410	-0.36	-0.12	1.68	0.195	-0.05	-0.12	0.28	0.597
<i>Parupeneus</i>	L_{inf}	157	135	2.76	0.097	173	135	5.29	0.021	N/A	135	N/A	N/A
<i>multifasciatus</i>	K	0.22	0.29	0.68	0.410	0.14	0.29	4.83	0.028	N/A	0.29	N/A	N/A
	t_0	-0.42	-0.30	0.06	0.806	-1.09	-0.30	1.56	0.212	N/A	-0.30	N/A	N/A

<https://doi.org/10.1371/journal.pone.0193426.t001>

Masinloc: *A. nigrofuscus* Growth Patterns




Fig 2. Scatterplot of the length-at-age of *A. nigrofuscus* individuals caught inside and outside of the Bani, San Salvador, and Taklobo Farm MPAs in Masinloc, and associated Von Bertalanffy growth curves for protected and fished populations.

<https://doi.org/10.1371/journal.pone.0193426.g002>

found in Tables 2 and 3, respectively. In general, the 95% confidence intervals around estimates of L_{50} and A_{50} for populations in MPAs and associated fished reefs were overlapping. However, the confidence intervals for L_{50} did not overlap for *P. multifasciatus* at either of the two MPAs from which it was collected and fished reefs, and for *A. nigrofuscus* in the Taklobo Farm MPA compared to fished reefs, with L_{50} values being greater inside MPAs in all cases (Table 2). Conversely, the confidence intervals for A_{50} overlapped in all but one instance (*C. striatus*, Bani MPA), which demonstrated A_{50} values lower inside the MPA compared to those in fished reefs (Table 3).

Discussion

A direct comparison of life-history characteristics in three reef fishes revealed considerable variation in fitness-relevant phenotypes of conspecifics, with disparities in asymptotic length (L_{inf}) and/or growth rates (K) being observed in each species within at least one MPA when compared to fished reefs. This result suggests that MPAs can promote shifts in the frequency of large-bodied fish within their borders with relative consistency across a multitude of species. The impact of MPA establishment on maturation schedules in protected populations,

Masinloc: *C. striatus* Growth Patterns




Fig 3. Scatterplot of the length-at-age of *C. striatus* individuals caught inside and outside of the Bani, San Salvador, and Taklobo Farm MPAs in Masinloc, and associated Von Bertalanffy growth curves for both protected and fished populations.

<https://doi.org/10.1371/journal.pone.0193426.g003>

however, was much less consistent. Although estimates of length at 50% maturity (L_{50}) were larger inside two MPAs for *P. multifasciatus* and one MPA for *A. nigrofuscus*, L_{50} values were not significantly different for *C. striatus* in any MPA. In addition, age at 50% maturity (A_{50}) differed only between the Bani MPA and fished reefs for *C. striatus*, but was lower inside the MPA than in fished reefs. That fish inside MPAs did not delay maturation until later ages may mitigate some of the potential reproductive benefits of larger asymptotic length and larger size at maturation. However, the increase of large-bodied, larger-maturing individuals is still likely to result in an overall increase in relative reproductive potential inside MPAs compared to adjacent, fished reefs.

The age of MPAs appeared to play a substantial role in whether phenotypes differed between protected and harvested populations. Significantly larger L_{inf} values were observed predominantly in the two oldest MPAs (San Salvador [26 years; *A. nigrofuscus*, *P. multifasciatus*], and Taklobo Farm [26 years; *A. nigrofuscus*, *C. striatus*]), although *C. striatus* also exhibited higher asymptotic length inside the Bani MPA (9 years). These results support theoretical predictions regarding the temporal scale required for populations inside MPAs to demonstrate phenotypic recovery, with demonstrable changes in life-history frequencies requiring many

Masinloc: *P. multifasciatus* Growth Patterns




Fig 4. Scatterplot of the length-at-age of *P. multifasciatus* individuals caught inside and outside of the Bani and San Salvador MPAs in Masinloc, and associated Von Bertalanffy growth curves for both protected and fished populations.

<https://doi.org/10.1371/journal.pone.0193426.g004>

years to appear [31, 49–50]. The size of MPAs, however, did not seem to be a contributing factor to phenotypic variation between sites. Populations in which life-history traits differed from fished reefs were found in both the largest (San Salvador [127 ha]) and smallest (Taklobo Farm [2 ha]) MPAs in the study, suggesting that long-term protection can promote beneficial changes across a range of MPA sizes. This result is unexpected, as movement into fishing grounds increases the selective force of fishing on protected populations, thereby dampening the benefits of protected areas [28, 51]. Species with small territories such as *C. striatus* (12m²

Table 2. Estimated length at 50% maturity (L₅₀) and associated 95% confidence intervals (CI) for each population. Values have been rounded to the nearest mm. Instances where 95% CIs did not overlap between any MPA and associated fished reef are indicated by bold font.

MPA/ Fished Reef	<i>A. nigrofuscus</i>			<i>C. striatus</i>			<i>P. multifasciatus</i>		
	L ₅₀	CI Lower	CI Upper	L ₅₀	CI Lower	CI Upper	L ₅₀	CI Lower	CI Upper
Bani	57	49	61	84	75	88	105	92	111
San Salvador	58	57	62	70	67	83	102	94	110
Taklobo Farm	64	62	66	79	76	88	N/A	N/A	N/A
Fished Reef	59	57	61	84	76	89	85	73	89

<https://doi.org/10.1371/journal.pone.0193426.t002>

Table 3. Estimated age at 50% maturity (A_{50}) and associated 95% confidence intervals (CI) for each population. Instances where 95% CIs did not overlap between any MPA and associated fished reef are indicated by bold font.

MPA/ Fished Reef	<i>A. nigrofuscus</i>			<i>C. striatus</i>			<i>P. multifasciatus</i>		
	A_{50}	CI Lower	CI Upper	A_{50}	CI Lower	CI Upper	A_{50}	CI Lower	CI Upper
Bani	1.02	0.53	1.49	0.98	0.91	1.03	5.02	4.00	6.48
San Salvador	1.09	1.01	1.53	1.01	0.94	1.49	5.71	4.53	6.63
Taklobo Farm	1.54	1.02	1.97	1.11	1.05	1.53	N/A	N/A	N/A
Fished Reef	1.06	0.99	1.50	1.49	1.15	1.91	4.11	3.37	5.14

<https://doi.org/10.1371/journal.pone.0193426.t003>

[52]), are not likely to traverse MPA boundaries even in the smallest MPAs examined here. In contrast, individuals of *A. nigrofuscus* are known to conduct daily feeding and spawning migrations from sheltered sites that can range from 300m [53] to 1.5km [54], and *P. multifasciatus* roams areas up to 245m² daily [55], making it unlikely that populations of these species are distinct between MPAs and fished reefs. It is also possible that the age and size of MPAs play synergistic roles in promoting phenotypic shifts in exploited populations. For example, increases in L_{inf} for *A. nigrofuscus* were greater in the San Salvador (127 ha; 26 years) MPA than the Taklobo Farm MPA (2 ha; 26 years), and asymptotic length of *C. striatus* was higher in the Bani MPA (50 ha; 9 years) than in the Taklobo Farm MPA. The effects of MPA size and age on the life-history variations will likely vary based on the genetic, plastic, and density-dependent processes controlling phenotypic shifts occurring inside MPAs [6].

There is considerable debate regarding the mechanism underlying phenotypic changes observed in exploited fish stocks. Specifically, the relative contributions of environmentally- or density-dependent phenotypic plasticity and fixed genetic alterations remain in question [56]. Although fish body-size and maturation are plastic in many species [31], the intensive harvest rates and size-selectivity of fisheries [5] and demonstrated heritability of life-history traits in fishes [3] provide the necessary conditions to produce genetic shifts in growth and maturation patterns [57]. Additionally, alterations to life-history traits have been shown to correlate with concurrent shifts in allelic frequencies [52], suggesting that phenotypic shifts in wild fish stocks have some genotypic basis. The distinction between phenotypic plasticity and genetics is critical to the recovery potential of exploited populations. Genetic changes are expected to be slower and more challenging to reverse than plastic responses [58–60], affect fundamental population dynamics [61], and impact resilience to environmental stress [62]. However, genetic shifts will be necessary for beneficial traits to extend to fished reefs through spillover and recruitment subsidy [26, 28]. Consequently, genotypic responses within MPAs would represent long-term benefits throughout interconnected populations, while plastic changes would only manifest inside MPAs, and may be rendered null if protection is ceased.

Given that the populations examined here are interconnected through both pelagic larval stages and adult movement patterns, genetic stratification between MPAs and fished reefs is unlikely. This suggests that observed phenotypic changes inside MPAs are more likely the result of plastic responses to the removal of fishing mortality. Conversely, the prevalence of life-history shifts in older MPAs indicates that multiple generations are required for alterations to growth rates and body sizes to occur, supporting a genotypic basis of growth and maturation phenotypes. It is also possible that extended periods of protection may be necessary for populations to reach densities great enough to induce plastic responses, in which case density-dependent growth may be primarily responsible for changes in life-history characteristics. Density-dependent mechanisms would help to explain life-history shifts in the Taklobo Farm (2 ha) and Bani (50 ha) MPAs, but not in the larger San Salvador MPA (127 ha). It is also conceivable that variations in environmental parameters such as habitat or food availability

between MPAs and fished areas may be responsible for observed shifts in growth patterns [44]. However, no distinct differences in coral cover or algal prevalence were observed between sites during the course of the study. Alternatively, increased growth and egg production inside MPAs may have been driven by differential stress levels between fished and protected populations, as fish flight initiation distance is positively associated with fishing pressure and decreases significantly inside MPAs [63–64]. Populations undergoing constant fishing pressure therefore spend more time avoiding fishing mortality and less time actively foraging, limiting the amount of potential energy that can be allocated to growth and reproduction. This interpretation is consistent with studies that have suggested that fishing pressure actively selects against bold behavioral types, which are often associated with increased reproductive output [65].

Although it appears that *A. nigrofuscus*, *C. striatus*, and *P. multifasciatus* have exhibited positive responses to MPA establishment, species-specific analyses across a wide range of taxa will be critical in determining the overall effect of MPAs on reef-fish communities, as there are many factors that may modulate the phenotypic impacts of MPAs. First, family-level susceptibility to fishing pressure [66], as well as recovery rates of individual fish families [67], differ significantly. Second, harvest intensity varies greatly depending on the species in question and the locality of the study [66, 68]. Differential harvest rates will affect the degree to which fishing-induced phenotypic shifts occur within a population, which will not only determine current population sizes and reproductive potential, but also the rate at which fisheries are likely to recover after the cessation of harvest pressure [15, 69]. Third, responses to protection depend on species-specific movement and behavioral patterns, economic value, body size, habitat depth range, and schooling behavior [51, 70]. Fourth, rates of evolutionary change have been shown to vary across conspecific fishery stocks that exhibit different somatic growth rates [71]. As a result, the rates and magnitude of life-history shifts are likely to differ not only across taxa, but also between disparate populations of any given species. Finally, traditional MPA theory predicts greater biological benefits from older and larger MPAs, necessitating the inclusion of the size and age of MPAs in assessments of their impact on fish life-history traits. Future empirical investigations and modelling studies will be critical in advancing our understanding of how populations will respond to spatially explicit cessation of harvest mortality, and how MPAs might best be employed to maximize their benefit to fishery stocks.

The restoration of healthy fishery populations is not only important for resource management, but is also essential for future ecological sustainability in Philippine coral reefs. Fishing pressure is considered a driving force behind deleterious ecological shifts within marine ecosystems [72], as the preferential removal of commercially targeted species significantly reduces functional diversity in exploited reefs [73]. Size- and age-truncated fishery stocks recover more slowly after fishing moratoria, as high reproductive investment at small size and early age diminish relative population growth under natural mortality scenarios [58], increasing the vulnerability of fisheries to environmental stress. MPAs have demonstrated the ability to act as buffers against mass mortality from isolated events [74], but their ability to combat population collapses against repeated, unpredictable environmental perturbations is less understood. As global climate patterns become increasingly variable, it is essential that MPAs promote the development of robust, highly fecund populations that can better withstand environmental heterogeneity [75]. Here, we have provided evidence of disparities in phenotypic frequencies between conspecific fish populations inside and outside of MPAs. Continued studies are necessary to improve our understanding of how ubiquitous these trends are across various MPAs, and whether phenotypic shifts will be substantial enough to promote long-term population resilience in harvested coral-reef fishes in the face of mounting anthropogenic pressures.

Supporting information

S1 Dataset. Dataset used for all analyses. In the tables, “F” represents females, “M” males, and “U” unknown gender. For gonad stages, “1” represents immature, “2” mature, and “3” mature, engorged and preparing to spawn. The term “STG” refers to individuals that had gonads damaged during collection, making sexing and staging unreliable.

(XLSX)

S1 Table. Total counts of all individuals collected from MPAs and fished reefs. Fish are divided by gender (male [M] or female [F]) and by site. Individuals listed as unknown (U) either had their gonads significantly damaged during collection, or were juveniles, making gender identification unreliable. The term “T” denotes the total number of fish caught in any given category.

(XLSX)

S1 Fig. Relative maturity by age (years) and size (standard length, mm) of collections of *A. nigrofuscus*.

(TIF)

S2 Fig. Relative maturity by age (years) and size (standard length, mm) of collections of *C. striatus*.

(TIF)

S3 Fig. Relative maturity by age (years) and size (standard length, mm) of collections of *P. multifasciatus*.

(TIF)

Acknowledgments

We thank the U.S. Student Fulbright Foundation and the Philippine-American Education Foundation, specifically Executive Director Dr. Esmeralda S. Cunanan, for their support of this research. We also thank the University of the Philippines Marine Science Institute as well as the Local Governmental Unit and citizens of Masinloc for their assistance in this research program. This is contribution #1 of the Junda Lin Professor of Marine Biology series.

Author Contributions

Conceptualization: Robert Y. Fidler, Ralph G. Turingan.

Data curation: Robert Y. Fidler.

Formal analysis: Robert Y. Fidler, Jessica Carroll, Kristen W. Rynerson, Danielle F. Matthews.

Funding acquisition: Robert Y. Fidler, Ralph G. Turingan.

Investigation: Robert Y. Fidler, Jessica Carroll, Kristen W. Rynerson, Danielle F. Matthews.

Methodology: Robert Y. Fidler, Jessica Carroll, Kristen W. Rynerson, Danielle F. Matthews, Ralph G. Turingan.

Project administration: Robert Y. Fidler, Danielle F. Matthews, Ralph G. Turingan.

Resources: Robert Y. Fidler, Jessica Carroll, Danielle F. Matthews.

Supervision: Robert Y. Fidler, Ralph G. Turingan.

Validation: Robert Y. Fidler, Jessica Carroll, Kristen W. Rynerson.

Writing – original draft: Robert Y. Fidler, Ralph G. Turingan.

Writing – review & editing: Robert Y. Fidler, Jessica Carroll, Kristen W. Rynerson, Danielle F. Matthews, Ralph G. Turingan.

References

1. Myers RA, Hoenig JM. Direct estimates of gear selectivity from multiple tagging experiments. *Can J Fish Aquat Sci* 1997; 54(1):1–9.
2. Birkeland C, Dayton PK. The importance in fishery management of leaving the big ones. *Trends Ecol Evol* 2005; 20(7):356–8. <https://doi.org/10.1016/j.tree.2005.03.015> PMID: 16701393
3. Law R. Fishing, selection, and phenotypic evolution. *ICES J Mar Sci* 2000; 57(3):659–69.
4. Law R. Fisheries-induced evolution: present status and future directions. *Mar Ecol Prog Ser* 2007; 335:271–7.
5. Heino M, Godø OR. Fisheries-induced selection pressures in the context of sustainable fisheries. *Bull Mar Sci* 2002; 70(2):639–56.
6. Kuparinen A, Merilä J. Detecting and managing fisheries-induced evolution. *Trends Ecol Evol* 2007; 22(12):652–9. <https://doi.org/10.1016/j.tree.2007.08.011> PMID: 17981361
7. Swain DP, Sinclair AF, Hanson MJ. Evolutionary response to size-selective mortality in an exploited fish population. *Proc Biol Sci* 2007; 274(1613):1015–22. <https://doi.org/10.1098/rspb.2006.0275> PMID: 17264058
8. Swain DP. Life-history evolution and elevated natural mortality in a population of Atlantic cod (*Gadus morhua*). *Evol Appl* 2011; 4(1):18–29. <https://doi.org/10.1111/j.1752-4571.2010.00128.x> PMID: 25567950
9. Heino M, Dieckmann U. Fisheries-induced evolution. In: *Encyclopedia of Life Sciences (ELS)*. Chichester, UK: John Wiley & Sons, Ltd. 2009.
10. Heino M, Dieckmann U, Godø OR. Measuring probabilistic reaction norms for age and size at maturation. *Evolution* 2002; 56(4):669–78. PMID: 12038525
11. Grift RE, Rijnsdorp AD, Barot S, Heino M, Dieckmann U. Fisheries-induced trends in reaction norms for maturation in North Sea plaice. *Mar Ecol Prog Ser* 2003; 257:247–57.
12. Taborsky B, Dieckmann U, Heino M. Unexpected discontinuities in life-history evolution under size-dependent mortality. *Proc Biol Sci* 2003; 270(1516):713–21. <https://doi.org/10.1098/rspb.2002.2255> PMID: 12713745
13. Ernande B, Dieckmann U, Heino M. Adaptive changes in harvested populations: plasticity and evolution of age and size at maturation. *Proc Biol Sci* 2004; 271(1537):415–23. <https://doi.org/10.1098/rspb.2003.2519> PMID: 15101701
14. Haugen TO, Vøllestad LA. A century of life-history evolution in grayling. *Genetica* 2001; 112–113:475–91. PMID: 11838784
15. Walsh MR, Munch SB, Chiba S, Conover DO. Maladaptive changes in multiple traits caused by fishing: impediments to population recovery. *Ecol Lett* 2006; 9(2):142–8. <https://doi.org/10.1111/j.1461-0248.2005.00858.x> PMID: 16958879
16. Bobko SJ, Berkeley SA. Maturity, ovarian cycle, fecundity, and age-specific parturition of black rockfish (*Sebastes melanops*). *Fish Bull* 2003; 102(3):418–29.
17. Berkeley SA, Chapman C, Sogard SM. Maternal age as a determinant of larval growth and survival in a marine fish, *Sebastes melanops*. *Ecology* 2004; 85(5):1258–64.
18. Conover DO, Munch SB. Sustaining fisheries yields over evolutionary time scales. *Science* 2002; 297(5578):94–6. <https://doi.org/10.1126/science.1074085> PMID: 12098697
19. Olsen EM, Heino M, Lilly GR, Morgan MJ, Brattey J, Ernande B, et al. Maturation trends indicative of rapid evolution preceded the collapse of northern cod. *Nature* 2004; 428(6986):932–5. <https://doi.org/10.1038/nature02430> PMID: 15118724
20. Olsen EM, Lilly GR, Heino M, Morgan MJ, Brattey J, Dieckmann U. Assessing changes in age and size at maturation in collapsing populations of Atlantic cod (*Gadus morhua*). *Can J Fish Aquat Sci* 2005; 62(4):811–23.
21. Hutchings JA, Rangeley RW. 2011. Correlates of recovery for Canadian Atlantic cod (*Gadus morhua*). *Can J Zool* 2011; 89(5):386–400.
22. Rose GA, Row S. Northern cod comeback. *Can J Fish Aquat Sci* 2015; 72:1789–98.
23. Stokes K, Law R. Fishing as an evolutionary force. *Mar Ecol Prog Ser* 2000; 208:307–9.

24. Stockwell CA, Hendry AP, Kinnison MT. Contemporary evolution meets conservation biology. *Evolution* 2003; 18(2):94–101.
25. Boonzaier L, Pauly D. Marine protection targets: an updated assessment of global progress. *Oryx* 2016; 50(1): 27–35.
26. Baskett ML, Levin SA, Gaines SD, Dushoff J. Marine reserve design and the evolution of size at maturation in harvested fish. *Ecol Appl* 2005; 15(3):882–901.
27. Dunlop ES, Baskett ML, Heino M, Dieckmann U. Propensity of MPAs to reduce the evolutionary effects of fishing in a migratory species. *Evol Appl* 2009; 2(3):371–93. <https://doi.org/10.1111/j.1752-4571.2009.00089.x> PMID: 25567887
28. Miethe T, Dytham C, Dieckmann U, Pitchford JW. MPAs and the evolutionary effects of fishing on size at maturation. *ICES J Mar Sci* 2010; 67(3):412–25.
29. Trexler JC, Travis J. Can marine protected areas restore and restore stock attributes of reef fishes? *Bull Mar Sci* 2000; 66(3):853–73.
30. Edeline E, Carlson SM, Stige LC. Trait changes in a harvested population are driven by a dynamic tug-of-war between natural and harvest selection. *Proc Natl Acad Sci USA* 2007; 104(40):15799–804. <https://doi.org/10.1073/pnas.0705908104> PMID: 17898170
31. Conover DO, Munch SB, Arnott SA. Reversal of evolutionary downsizing caused by selective harvest of large fish. *Proc Biol Sci* 2009; 276:2015–20. <https://doi.org/10.1098/rspb.2009.0003> PMID: 19324761
32. Babcock RC, Shears NT, Alcala AC, Barrett NS, Edgar GJ, Lafferty KD, et al. 2010. Decadal trends in MPAs reveal differential rates of change in direct and indirect effects. *Proc Natl Acad Sci USA* 2010; 107(43):18256–61. <https://doi.org/10.1073/pnas.0908012107> PMID: 20176941
33. Russ GR. MPAs as reef fisheries management tools: yet another review. In: Sale PF, (editor). *Coral Reef Fishes. Dynamics and Diversity in a Complex Ecosystem*. 2002. San Diego, CA (USA): Academic Press. p 421–4.
34. Mosqueira I, Côté IM, Jennings S, Reynolds JD. Conservation benefits of MPAs for fish populations. *Anim Conserv* 2000; 4:321–32.
35. Lester SE, Halpern BS, Grorud-Colvert K, Lubchenco J, Ruttenberg BI, Gaines SD, et al. Biological effects within no-take marine reserves: a global synthesis. *Mar Ecol Prog Ser* 2009; 384:33–46.
36. Fidler RY, Maypa A, Apistar D, White AT, Turingan RG. Body size shifts in Philippine reef fishes: interfamilial variation in responses to protection. *Biology* 2014; 3(2):264–80. <https://doi.org/10.3390/biology3020264> PMID: 24833509
37. Soykan CU, Lewison RL. Using community-level metrics to monitor the effects of marine protected areas on biodiversity. *Conserv Biol* 2015; 29(3):775–83. <https://doi.org/10.1111/cobi.12445> PMID: 25572325
38. Adams S, Mapstone BD, Russ GR, Davies CR. Geographic variation in the sex ratio, sex specific size, and age structure of *Plectropomus leopardus* (Serranidae) between reefs open and closed to fishing on the Great Barrier Reef. *Can J Fish Aquat Sci* 2000; 57:1448–58.
39. Fidler RY, Turingan RG, White AT, Alava M. Reef-wide beneficial shifts in fish population structure following establishment of marine protected areas (MPAs) in Philippine coral reefs. *Mar Ecol Prog Ser* 2017; 570:187–202.
40. Evans RD, Russ GR, Krizter JP. Batch fecundity of *Lutjanus carponotatus* (Lutjanidae) and implications of no-take marine reserves on the Great Barrier Reef, Australia. *Coral Reefs* 2008; 27: 179–89.
41. Carter AB, Davies CR, Mapstone BD, Russ GR, Tobin AJ, Williams AJ. Effects of region, demography, and protection from fishing on batch fecundity of common coral trout (*Plectropomus leopardus*). *Coral Reefs* 2014; 33:751–63.
42. Carter AB, Davies CR, Emslie MJ, Mapstone BD, Russ GR, Tobin AJ, et al. Reproductive benefits of no-take marine reserves vary within region for an exploited coral reef fish. *Scientific Reports* 2017; 7:9693. <https://doi.org/10.1038/s41598-017-10180-w> PMID: 28852089
43. Murua H, Kraus G, Saborido-Rey F, Witthames PR, Thorsen A, Junquera S. Procedures to estimate fecundity of marine fish species in relation to their reproductive strategy. *J Northwest Atl Fish Sci* 2003; 33: 33–54.
44. Hart AM, Russ GR. Response of herbivorous fishes to crown-of-thorns starfish *Acanthaster planci* outbreaks. III. Age, growth, mortality and maturity indices of *Acanthurus nigrofuscus*. *Mar Ecol Prog Ser* 1996; 136:25–35.
45. Choat JH, Axe LM. Growth and longevity in acanthurid fishes; an analysis of otolith increments. *Mar Ecol Prog Ser* 1996; 134:15–26.
46. Kimura DK. Likelihood methods for the Von Bertalanffy growth curve. *Fish Bull* 1980; 77(4):765–76.
47. Nelson GA. *Fishmethods: Fishery Science Methods and Models in R*. R package version 1.10–4. 2017.

48. Ogle D.H. FSA: Fisheries Stock Analysis. R package version 0.8.17. 2017.
49. McClanahan TR, Humphries A. Differential and slow life-histories responses of fishes to coral reef closures. *Mar Ecol Prog Ser* 2012; 469:121–31.
50. McClanahan TR, Graham NAJ. Marine reserve recovery rates towards a baseline are slower for reef fish community life histories than biomass. *Proc Biol Sci* 2015; 282(1821):20151938. <https://doi.org/10.1098/rspb.2015.1938> PMID: 26702040
51. Grüss A, Kaplan DM, Guénette S, Roberts CM, Botsford LW. Consequences of adult and juvenile movement for marine protected areas. *Biol Conserv* 2011; 144(2):692–702.
52. Krone R, Bshary R, Paster M, Eisinger M, van Treeck P, Schuhmacher H. 2008. Defecation behavior of the lined bristletooth surgeonfish *Ctenochaetus striatus* (Acanthuridae). *Coral Reefs* 2008; 27(3):619–22.
53. Robertson DR. On the spawning behavior and spawning cycles of eight surgeonfishes (Acanthuridae) from the Indo-Pacific. *Environ Biol Fishes* 1983; 9(3):193–223.
54. Mazeroll AI, Montgomery WL. Structure and organization of local migrations in brown surgeonfish (*Acanthurus nigrofuscus*). *Ethology* 1995; 99(1–2):89–106.
55. McCormick MI, Makey LJ. Post-settlement transition in coral reef fishes: overlooked complexity in niche shifts. *Mar Ecol Prog Ser* 1997; 153:247–57.
56. Eikeset AM, Dunlop ES, Heino M, Størvik G, Stenseth NC, Dieckmann U. Roles of density-dependent growth and life history evolution in accounting for fisheries-induced trait changes. *Proc Natl Acad Sci USA* 2016; 113(52):15030–5. <https://doi.org/10.1073/pnas.1525749113> PMID: 27940913
57. Sharpe DM, Hendry AP. Life history change in commercially exploited fish stocks: an analysis of trends across studies. *Evol Appl* 2009; 2(3):260–75. <https://doi.org/10.1111/j.1752-4571.2009.00080.x> PMID: 25567879
58. Uusi-Heikkilä S, Whiteley AR, Kuparinen A, Matsumura S, Venturelli PA, Wolter C, et al. The evolutionary legacy of size-selective harvesting extends from genes to populations. *Evol Appl* 2015; 8(6):597–620. <https://doi.org/10.1111/eva.12268> PMID: 26136825
59. de Roos AM, Boukal DS, Persson L. Evolutionary regime shifts in age and size at maturation of exploited fish stocks. *Proc Biol Sci*. 2006; 273(1596):1873–80. <https://doi.org/10.1098/rspb.2006.3518> PMID: 16822746
60. Enberg K, Jørgensen C, Dunlop ES, Heino M, Dieckmann U. Implications of fisheries-induced evolution for stock rebuilding and recovery. *Evol Appl* 2009; 2(3):394–414. <https://doi.org/10.1111/j.1752-4571.2009.00077.x> PMID: 25567888
61. Dunlop ES, Eikeset AM, Stenseth NC. From genes to populations: how fisheries-induced evolution alters stock productivity. *Ecol Appl* 2015; 25(7):1860–8. PMID: 26591452
62. Hsieh C, Yamauchi A, Nakazawa T, Wang W. Fishing effects on age and spatial structures undermine population stability of fishes. *Aquat Sci* 2010; 72(2):165–78.
63. Januchowski-Hartley FA, Graham NAJ, Feary DA, Morove T, Cinner JE. Fear of fishers: human predation explains behavioral changes in coral reef fishes. *PLoS ONE* 2011; 6: e22761. <https://doi.org/10.1371/journal.pone.0022761> PMID: 21853046
64. Januchowski-Hartley FA, Nash KL, Lawton RJ. Influence of spear guns, dive gear and observers on estimating fish flight initiation distance on coral reefs. *Mar Ecol Prog Ser* 2012; 469:113–9.
65. Mittlebach GG, Ballew NG, Kjølvik MK. Fish behavioral types and their ecological consequences. *Can J Fish Aquat Sci* 2014; 71:927–44.
66. Russ GR, Alcala AC. Natural fishing experiments in MPAs 1983–1993: roles of life history and fishing intensity in family responses. *Coral Reefs* 1998; 17(4):399–416.
67. Jennings S. Patterns and prediction of population recovery in marine reserves. *Rev Fish Biol Fish* 2001; 10(2):209–31.
68. Russ GR, Alcala AC. Effects of intense fishing pressure on an assemblage of coral reef fishes. *Mar Ecol Prog Ser* 1989; 56:13–27.
69. Kuparinen A, Hutchings JA. Consequences of fisheries-induced evolution for population productivity and recovery potential. *Proc Biol Sci* 2012; 279(1738):2571–79. <https://doi.org/10.1098/rspb.2012.0120> PMID: 22398166
70. Claudet J, Osenberg CW, Domenici P, Badalamenti F, Milazzo M, Falcón JM, et al. 2010. Marine reserves: fish life history and ecological traits matter. *Ecol Appl* 2010; 20(3):830–9. PMID: 20437967
71. Devine JA, Wright PJ, Pardoe HI, Heino M. Comparing rates of contemporary evolution in life-history traits for exploited fish stocks. *Can J Fish Aquat Sci* 2012; 69(6):1105–20.

72. Palkovacs EP, Kinnison MT, Correa C, Dalton CM, Hendry AP. Fates beyond traits: ecological consequences of human-induced trait change. *Evol Appl* 2011; 5(2):183–91. <https://doi.org/10.1111/j.1752-4571.2011.00212.x> PMID: 25568040
73. Martins GM, Arenas F, Neto AI, Jenkins SR. Effects of fishing and regional species pool on the functional diversity of fish communities. *PLoS ONE* 2012; 7(8):e44297. <https://doi.org/10.1371/journal.pone.0044297> PMID: 22952950
74. Micheli F, Saenz-Arroyo A, Greenley A, Vazquez L, Montes JAE, Rossetto M, et al. Evidence that marine reserves enhance resilience to climatic impacts. *PLoS ONE* 2012; 7(7):e40832. <https://doi.org/10.1371/journal.pone.0040832> PMID: 22855690
75. McLeod E, Salm R, Green A, Almany J. Designing marine protected area networks to address the impacts of climate change. *Front Ecol Environ* 2009; 7(7):362–70.