

## CORRECTION

# Correction: The Ketamine Analogue Methoxetamine and 3- and 4-Methoxy Analogues of Phencyclidine Are High Affinity and Selective Ligands for the Glutamate NMDA Receptor

Bryan L. Roth, Simon Gibbons, Warunya Arunotayanun, Xi-Ping Huang, Vincent Setola, Ric Treble, Les Iversen


Part of the data are not included in the Supporting Information file [S1 Table](#). Please see the complete, correct [S1 Table](#) here.

## Supporting information

**S1 Table. Ki values for ketamine, methoxetamine, phencyclidine and analogues.**  
(XLSX)

## Reference

1. Roth BL, Gibbons S, Arunotayanun W, Huang X-P, Setola V, Treble R, et al. (2013) The Ketamine Analogue Methoxetamine and 3- and 4-Methoxy Analogues of Phencyclidine Are High Affinity and Selective Ligands for the Glutamate NMDA Receptor. PLoS ONE 8(3): e59334. <https://doi.org/10.1371/journal.pone.0059334> PMID: 23527166



## OPEN ACCESS

**Citation:** Roth BL, Gibbons S, Arunotayanun W, Huang X-P, Setola V, Treble R, et al. (2018) Correction: The Ketamine Analogue Methoxetamine and 3- and 4-Methoxy Analogues of Phencyclidine Are High Affinity and Selective Ligands for the Glutamate NMDA Receptor. PLoS ONE 13(3): e0194984. <https://doi.org/10.1371/journal.pone.0194984>

**Published:** March 22, 2018

**Copyright:** © 2018 Roth et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.